

# Type 2 diabetes mellitus in Canada's First Nations: status of an epidemic in progress

T. Kue Young, Jeff Reading, Brenda Elias, John D. O'Neil

## Abstract

THIS REVIEW PROVIDES A STATUS REPORT ON THE EPIDEMIC of type 2 diabetes mellitus that is affecting many of Canada's First Nations. We focus on the published literature, especially reports published in the past 2 decades, and incorporate data from the Aboriginal Peoples Survey and the First Nations and Inuit Regional Health Survey. We look at the extent and magnitude of the problem, the causes and risk factors, primary prevention and screening, clinical care and education, and cultural concepts and traditional knowledge. The epidemic of type 2 diabetes is still on the upswing, with a trend toward earlier age at onset. Genetic-environmental interactions are the likely cause. Scattered intervention projects have been implemented and evaluated, and some show promise. The current health and social repercussions of the disease are considerable, and the long-term outlook remains guarded. A national Aboriginal diabetes strategy is urgently needed.

It is widely recognized that type 2 diabetes mellitus has become a serious health problem among many Aboriginal populations in North America. In the 1970s West<sup>1</sup> provided a global survey and hinted at the emergence of an epidemic. The Pima study in Arizona contributed substantially to our understanding of the causes, pathogenesis and epidemiological features of the disease.<sup>2</sup>

In Canada a few clinical studies on carbohydrate metabolism were conducted in the 1970s among Inuit hospital patients.<sup>3,4</sup> During the 1980s prevalence studies were performed in various regions, using a variety of survey methods and diagnostic criteria.<sup>5-11</sup> Table 1 identifies recent developments that signal the increased awareness of the problem among academic researchers, service providers, and representatives of governments and Aboriginal organizations.

Diabetes can be considered to be indicative of the rapid sociocultural changes experienced by Aboriginal people in the past several decades.<sup>12</sup> In this article we update an earlier review<sup>13</sup> by including new data collected from several community studies, the 1991 Aboriginal Peoples Survey (APS)<sup>14</sup> and the 1997 First Nations and Inuit Regional Health Survey (FNIRHS).<sup>15</sup> We cover the epidemiological features and causes of the disease and issues relevant to providing health care to Aboriginal people with diabetes.

We focus on First Nations people in Canada and exclude Inuit and Metis, the 2 other constitutionally defined Aboriginal groups. Diabetes is not yet an important health problem among the Inuit,<sup>16</sup> although the situation may change.<sup>17</sup> Little data are available for the Metis, although they indicate that this group is at an increased risk relative to the overall Canadian population.<sup>18</sup> We also do not review the massive literature on American Indians in the United States.<sup>19</sup>

## Extent and magnitude of the problem

There are several ways to determine how much of a health problem diabetes is in the population, including vital statistics derived from death certificates, registries of diagnosed cases, health interview surveys and screening surveys involving measurement of plasma glucose levels (fasting or 2 hours after oral glucose challenge, or both).

Using data from the Canadian Mortality Database, Mao and colleagues<sup>20</sup> found a greater than 5-fold risk of death from diabetes among women resident on reserves compared with Canadians nationally.

## Review

## Synthèse

**Dr. Young is Professor and Head, Dr. Reading is Assistant Professor, Ms. Elias is Research Associate and Dr. O'Neil is Professor and Director, Northern Health Research Unit, Department of Community Health Sciences, Faculty of Medicine, University of Manitoba, Winnipeg, Man.**

*This article has been peer reviewed.*

CMAJ 2000;163(5):561-6

[Return to September 5, 2000 Table of Contents](#)

Based on case registries maintained by Health Canada's Medical Services Branch, the federal agency responsible for the health care of most First Nations, the prevalence of diabetes has been found to vary according to language group, culture area, geographic location and degree of isolation, both nationally<sup>21</sup> and regionally.<sup>22</sup> Of the main language groups in Canada, the Dene (Athapaskans) have the lowest reported rate.<sup>16,21,22</sup> Substantial variation can exist even between communities of the same tribal background in close proximity.<sup>23,24</sup>

Health interview surveys provide an estimate of self-reported prevalence. Both the APS<sup>14</sup> and the FNIRHS<sup>15</sup> included questions on diabetes and its treatment. The APS was a survey of all self-identified First Nations, Inuit and Metis across Canada conducted by Statistics Canada after the 1991 census. The FNIRHS had a sample size of 9870 people aged 15 years or more; when weights derived from the age and sex strata of each regional target population were applied, it provided prevalence estimates for a national target population of 199 782 people. It covered First Nations people living on reserves in all provinces (except Newfoundland) as well as the Inuit of Labrador. The age- and sex-specific prevalence rates of self-reported diabetes from these 2 sources are shown in Fig. 1, together with all-Canadian data from the 1994 National Population Health Survey.<sup>25</sup> The crude prevalence of diabetes in the FNIRHS was 8% among men and 13% among women. When age-adjusted to the Canadian population, the prevalence was 3.6 and 5.3 times higher among First Nations men and women

respectively than among Canadian men and women.

Oral glucose tolerance surveys have been conducted in a few localities. In 2 Algonquin reserves in northeastern Quebec<sup>23,26</sup> and among the Oji-Cree of Sandy Lake, in northwestern Ontario,<sup>27</sup> the prevalence reached as high as 25% among all adults and 80% among women aged 50–64 years. New cases detected by oral glucose tolerance screening account for 40% of all diabetes cases,<sup>27</sup> an indication that diagnosed cases do not represent the total burden of disease.

Increasingly, type 2 diabetes is observed in children and adolescents.<sup>28–30</sup> In the Island Lake region of northeastern Manitoba, new cases in children as young as 8 years have been identified through screening of fasting plasma glucose levels.<sup>31</sup> The prevalence among females aged 10–19 years has been reported to be 3.6%.<sup>31</sup>

Before the 1950s type 2 diabetes was rare in Aboriginal populations.<sup>32</sup> Within the past 2 decades, a rapid increase in prevalence has been documented in a few regions where there had been continuous surveillance. In the Sioux Lookout Zone of northwestern Ontario the prevalence increased by 45% over a 10-year period.<sup>33</sup> In Saskatchewan the rate doubled between 1980 and 1990.<sup>22</sup>

The public health significance of diabetes is manifested in the various long-term complications affecting the cardiovascular system, eyes, kidneys and nerves, resulting in premature death, disability and a compromised quality of life.

In the FNIRHS, 50% of the respondents with diabetes also reported having hypertension, and 26% reported having heart disease, prevalence rates that were 3.3 and 3.9, respectively, times those among the respondents without diabetes<sup>15</sup> (Fig. 2).

Analyses of data from the Manitoba database on use of health care services indicate that, compared with non-Aboriginal patients, Aboriginal patients with diabetes have higher rates of hospital separations and readmissions and longer stays, in both Winnipeg and nonurban areas.<sup>34</sup>

In Kahnawake, Que., over 60% of patients with diabetes were found to have at least one major complication.<sup>35</sup> As the duration of illness is a strong predictor of complications,<sup>6</sup> the rate of complications can be expected to rise as the age at onset decreases.<sup>28–31</sup> Among the James Bay Cree, factors associated with the presence of renal or retinal microvascular complications have included poor glycemic control and hypertriglyceridemia, in addition to duration of illness greater than 5 years.<sup>36</sup>

It has been shown nationally<sup>37</sup> and regionally<sup>38,39</sup> that Aboriginal people are at increased risk for end-stage renal disease, a substantial proportion of cases of which can be attributed to diabetes.

A high prevalence of microalbuminuria and macroalbuminuria, indicative of eventual renal failure, has been found among both insulin-using and non-insulin-using patients in southern Alberta First Nations.<sup>40</sup> They also have a high prevalence of serious and untreated diabetic retinopathy.<sup>40</sup>

Diabetes increases the susceptibility to infection. In Manitoba, Aboriginal women are admitted to hospital 5 to

**Table 1: Diabetes mellitus in the Canadian Aboriginal population: a chronology of significant recent developments**

1985	Canadian Diabetes Association produced videotape <i>Walking in Balance</i> and resource kit "Diabetes and Diabetes Education among the Native Population"
1986	National Workshop on Diabetes Research in the Canadian Native Population convened in Toronto, funded by the National Health Research and Development Program (NHRDP) of Health and Welfare Canada
1987	Medical Services Branch of Health and Welfare Canada established an education working group, which produced the "Duncan Declaration on Standards of Care and Education for Native Peoples with Diabetes"
1990	First International Conference on Diabetes and Indigenous Peoples held in Minneapolis, with strong Canadian participation; subsequent conferences held in Honolulu (1993), Winnipeg (1995) and San Diego (1997)
1991	NHRDP launched Special Competition on Diabetes in the Native Population and funded 6 research projects across the country
1992	Canadian Diabetes Advisory Board produced clinical practice guidelines with special section on Aboriginal people
1996	National Aboriginal Diabetes Association formed
1997	Health Canada initiated consultations with Aboriginal organizations to develop a national Aboriginal diabetes strategy
1999	Federal government allocated significant resources to the control of diabetes in Aboriginal people

20 times more frequently for acute pyelonephritis than non-Aboriginal women.<sup>41</sup> The higher rates are partially attributable to the greater frequency of pregnancy and diabetes among Aboriginal women.

Diabetes affects the psychosocial well-being of affected people. Those with diabetes are more likely to regard their health as poor and to experience some limitation in their daily activity<sup>15</sup> (Fig. 2).

### Causes and risk factors

Diabetes is a chronic disease with multifactorial causes involving the interactions of genetic susceptibility and environmental factors.

The Sandy Lake Study has yielded a wealth of new genetic data.<sup>42-44</sup> Systematic genome-wide scanning for diabetes susceptibility among affected sibling pairs revealed 4 markers suggestive of association with diabetes.<sup>45</sup> A new variant of the gene encoding the hepatic nuclear factor-1 $\alpha$ , believed to be unique to the Oji-Cree, was also discovered.<sup>46</sup> The S319 allele is present in 20% of diabetic subjects, twice the frequency among nondiabetic subjects. Diabetic subjects who are homozygous for S319 tend to have an earlier age at onset, a lower body mass index and a higher postchallenge plasma glucose level.<sup>46</sup>

Plasma insulin levels have not been studied extensively in Canadian populations. Among the Dogrib, who are at relatively low risk for diabetes, the mean plasma insulin level is comparable to that of people of European ancestry.<sup>47</sup> Fasting insulin levels vary according to the Gc genotype, the group-specific component that binds vitamin D, involved in the regulation of plasma insulin.<sup>48</sup>

Among Algonquins in Quebec, gestational diabetes is associated with higher serum triglyceride levels, measures of obesity and fasting insulin level.<sup>49</sup> Gestational diabetes is a strong predictor of high birth weight in the newborn,<sup>50-53</sup>

which predisposes to future diabetes. Surveys in northern Ontario and Quebec have shown that gestational diabetes may affect as many as 13% of pregnancies among Aboriginal women.<sup>51-53</sup> Among women over age 35, just under half of all pregnancies are associated with pre-existing type 2 diabetes or gestational diabetes;<sup>51</sup> this substantially increases the risk of diabetes in the next generation.

Cross-sectional surveys have tended to show an association between diabetes and hyperlipidemia and between diabetes and obesity.<sup>23,27,54</sup> Unlike half a century ago, obesity is now prevalent in many Aboriginal populations.<sup>55</sup> The obesity among Aboriginal people is predominantly of the central (or abdominal, upper-body) type, characterized by a high waist-to-hip ratio,<sup>5,23,27,56,57</sup> which is associ-




Fig. 1: Prevalence of self-reported diabetes mellitus among all Canadians (1994<sup>25</sup>) and First Nations people (1991<sup>14</sup> and 1997<sup>15</sup>).




Fig. 2: Self-rated health, concomitant conditions and disability of respondents with and without diabetes in the First Nations and Inuit Regional Health Survey.<sup>15</sup>

ated with a higher risk of diabetes. Abdominal obesity among the Salishan was found to be associated with hyperinsulinemia in younger people and with elevated glycated hemoglobin concentration in older people.<sup>58</sup> The Sandy Lake Study indicated a positive association between levels of serum leptin, the genetically regulated hormone produced by adipose tissue, and measures of body fat and plasma insulin.<sup>59</sup>

An association between diabetes and dietary factors has not been demonstrated consistently. Two studies from northern Ontario and Manitoba have shown that diabetic subjects consume a higher level of proteins (adjusting for total caloric intake) than nondiabetic subjects but differ in terms of the protective effect of fibre.<sup>54,60</sup> In Sandy Lake, diabetes was found to be associated with the consumption of “junk foods” and a “fatty” method of food preparation.<sup>61</sup> Although type 1 diabetes has been reported to be associated with an impaired metabolic availability of vitamin A (retinol), a similar relation with type 2 diabetes was not found in a study among the Plains Cree of central Alberta.<sup>62</sup>

Dietary acculturation, particularly the substitution of modern for traditional food items, has been observed in many Aboriginal communities.<sup>12,63</sup> Nationally, only a minority (15%) of Aboriginal people still obtain most of their meat and fish from hunting and fishing.<sup>14</sup>

Lack of physical activity is an important risk factor.<sup>64</sup> The adoption of a more sedentary lifestyle often accompanies the transition to a modern diet. The APS showed that 54% of Aboriginal adults nationally participate in leisure-time activity.<sup>14</sup>

The “thrifty genotype” is often invoked to explain why diabetes is so prevalent today in many Aboriginal populations.<sup>65</sup> In times of food shortage, it enables the rapid production of insulin in response to rising blood glucose levels, which facilitates the storage of glucose in the form of triglycerides in fat cells. With the assurance of a continuous and ample food supply, the quick insulin trigger results in hyperinsulinemia, hyperglycemia, obesity and diabetes. Critics of the theory note that it assumes a nutritional environment in which carbohydrate intake exceeds daily energy requirements.<sup>66-68</sup> The early occupants of North America lived in an arctic or subarctic environment on a protein- and fat-based diet with little carbohydrates. Alternative pathways to provide energy (e.g., through enhanced gluconeogenesis and release of free fatty acids) may have been favoured by natural selection.

## Primary prevention and screening

The primary prevention of diabetes involves the promotion of healthy behaviours, especially achieving and maintaining a healthy body weight through increased physical activity and a balanced diet. Equally important is a supportive environment conducive to behavioural change. A variety of community-based diabetes prevention projects have begun in a few locations across Canada, most notably in

Kahnawake,<sup>69,70</sup> Sandy Lake,<sup>71,72</sup> and Okanagan First Nation in British Columbia.<sup>73,74</sup> These participatory research projects, with intervention sites in schools, stores and the community at large, respect Aboriginal culture, traditions and learning styles, and have achieved a high degree of community support and awareness. A project focusing on physical activity and directed at gestational diabetes was initiated in Saskatoon.<sup>64</sup>

Although national expert committees such as the Canadian Task Force on the Periodic Health Examination<sup>75</sup> (now the Canadian Task Force on Preventive Health Care) do not recommend screening for diabetes in the general population, the more recent Canadian clinical practice guidelines for the management of diabetes recommend testing fasting plasma glucose levels in all Canadians over age 45 once every 3 years and more frequently in high-risk groups,<sup>76</sup> which would include Aboriginal people.

## Clinical care and education

Health care professionals who serve Aboriginal people need to adapt their treatment plans and education programs to the culture and social environment of their patients. Furthermore, many First Nations are geographically remote, with little access to needed specialized services. The cultural dimensions of diabetes care were highlighted in the submission from Alethea Kewayosh, coordinator of diabetes education with the Assembly of First Nations, to the Royal Commission on Aboriginal Peoples.<sup>77</sup>

The APS revealed a high use of health care services: over 90% of diabetic people reported having seen a health care professional in the year before the survey, as compared with 75% of those without diabetes.<sup>78</sup> Over 80% of the respondents with diabetes in the FNIRHS considered their health care to be in need of improvement, requiring more staff, chronic care facilities, home care, teaching about medications, education about prevention and mental health counselling.<sup>15</sup> Diabetic respondents were less likely than nondiabetic respondents to be current smokers (50% v. 64%).<sup>15</sup> This finding suggests some success in lifestyle modification, although a prevalence of smoking of 50% is still too high.

The 1998 Canadian clinical practice guidelines make 3 recommendations specifically directed at the Aboriginal population.<sup>76</sup> These relate to the need for community-based screening and primary prevention programs and urge respect for, and sensitivity to, language and cultural issues.

The effects of a traditional lifestyle on metabolic control in patients with diabetes was investigated among the James Bay Cree in northern Quebec.<sup>79</sup> Diabetic subjects who spent 3 months in the bush did not differ from those who stayed in the village in body weight, plasma glucose level, glycated hemoglobin concentration and blood pressure. Although bush dwellers were more active physically than those who remained in the village, they also brought along large quantities of store-bought foods for sustenance.

## Cultural concepts and traditional knowledge

It is important for non-Aboriginal health care professionals to understand how Aboriginal people interpret their illness experience and respond to treatment regimens, and to respect the logic and rationality of another system of thought.

Many Aboriginal people consider diabetes an example of "white man's illness," a new, introduced disease similar to smallpox and tuberculosis in the past. The adoption of modern foods and the decline of hunting and fishing are widely believed to be the underlying causes of the epidemic.<sup>80,81</sup>

Balance is central to the Aboriginal understanding of diabetes. The concept can be used effectively in explaining how diabetes works and suggesting means to cope with the illness. Aboriginal legends can also be incorporated into teaching materials.<sup>82</sup>

The concept of obesity is culture specific.<sup>83</sup> In Sandy Lake, researchers found that 16% of the community members were satisfied with their current body shape.<sup>84</sup> Those with a higher body mass index were less satisfied with their bodies and believed themselves to be less healthy than people with a lower body mass index.

New research on Aboriginal understanding and explanation of diabetes, using qualitative methods, has been conducted in such communities as the James Bay Cree<sup>85</sup> and the Cape Breton Mi'kmaq.<sup>86</sup>

Aboriginal peoples have a long tradition of using the healing properties of plants in the environment.<sup>87</sup> Ethnobotanical research may yet reveal effective remedies. A single-blind, placebo-controlled trial of Aboriginal herbal tea in the treatment of diabetes failed to show a significant reduction in blood glucose levels.<sup>88</sup> Traditional plant beverages, regardless of their medicinal value, may offer healthful substitutes to sweetened soft drinks. Much research needs to be done to test and document the uses of indigenous plants.

## Conclusion

The epidemic of diabetes in many First Nations communities in Canada is still on the upswing, and its causes remain to be elucidated. Scattered intervention projects have been implemented, and some show promise. The health and social repercussions of the disease are considerable, and the long-term outlook remains guarded.

Diabetes is not simply a metabolic disorder that can be adequately dealt with on an individual basis in a clinic or hospital ward. Its prevention and control require community action and collaboration among Aboriginal organizations, governments, voluntary agencies and health care professionals.

We thank the National Steering Committee of the First Nations and Inuit Regional Health Survey for its cooperation in releasing data relating to the survey for use in this paper.

Dr. Young is supported by a Medical Research Council of Canada Senior Scientist Award.

Competing interests: None declared.

## References

- West KM. Diabetes in American Indians and other native populations of the New World. *Diabetes* 1974;23:841-55.
- Knowler WC, Pettitt DJ, Saad MF, Bennett PH. Diabetes mellitus in Pima Indians: incidence, risk factors and pathogenesis. *Diabetes Metab Rev* 1990;6:1-27.
- Schaefer O. Glucose tolerance testing in Canadian Eskimos: a preliminary report and a hypothesis. *CMAJ* 1968;99(6):252-62.
- Schaefer O, Crockford PM, Romanowski B. Normalization effect of preceding protein meals on "diabetic" oral glucose tolerance in Eskimos. *CMAJ* 1972;107(8):733-8.
- Szathmary EJ, Holt N. Hyperglycemia in Dogrib Indians of the Northwest Territories, Canada: association with age and a centripetal distribution of body fat. *Hum Biol* 1983;55:493-515.
- Young TK, McIntyre LL, Dooley J, Rodriguez J. Epidemiologic features of diabetes mellitus among Indians in northwestern Ontario and northeastern Manitoba. *CMAJ* 1985;132(7):793-97.
- Montour LT, Macaulay AC, Adelson N. Diabetes mellitus in Mohawks of Kahnawake, PQ: a clinical and epidemiologic description. *CMAJ* 1989;141(6):549-52.
- Evers S, McCracken E, Antone I, Deagle G. Prevalence of diabetes in Indians and Caucasians living in southwestern Ontario. *Can J Public Health* 1987;78:240-3.
- Young TK, Krahn J. Comparison of screening methods in a diabetes prevalence survey among northern Indians. *Clin Invest Med* 1988;11:380-5.
- Brassard P, Robinson E, Lavallée C. Prevalence of diabetes mellitus among the James Bay Cree of northern Quebec. *CMAJ* 1993;149(3):303-7.
- Young TK, editor. *Diabetes in the Canadian Native population: biocultural perspectives*. Toronto: Canadian Diabetes Association; 1987.
- Young TK. *The health of Native Americans: toward a biocultural epidemiology*. New York: Oxford University Press; 1994. p. 145-68.
- Young TK. Diabetes mellitus among Native Americans in Canada and the United States: an epidemiological review. *Am J Hum Biol* 1993;5:399-413.
- 1991 Aboriginal peoples survey: language, tradition, health, lifestyle and social issues. Ottawa: Statistics Canada; 1993. Cat no 89-533.
- National Steering Committee for the First Nations and Inuit Regional Health Survey. *Final report*. 1999. [Copies can be purchased from Ms. Gail McDonald, Akwesasane Mohawk Territory, McDonald Rd., St. Regis QC H0M 1A0; fax 613 574-1443; gmcd\_akw@glen-net.ca]
- Young TK, Schrager CD, Shubnikoff EV, Szathmary EJ, Nikitin YP. Prevalence of diagnosed diabetes in circumpolar indigenous populations. *Int J Epidemiol* 1992;21:730-6.
- Bjerregaard P, Young TK. *The circumpolar Inuit: health of a population in transition*. Copenhagen: Munksgaard; 1998. p. 127-9.
- Bruce S. The prevalence and determinants of diabetes mellitus among the Metis of western Canada. *Am J Hum Biol* 2000;12:542-51.
- Ghodes D. Diabetes in North American Indians and Alaska Natives. In: National Diabetes Data Group. *Diabetes in America*. 2nd ed. Bethesda (MD): National Institutes of Health; 1995. p. 683-701. NIH publ no 95-1468.
- Mao Y, Moloughney BW, Semenciw RM, Morrison HI. Indian Reserve and registered Indian mortality in Canada. *Can J Public Health* 1992;83:350-3.
- Young TK, Szathmary EJ, Evers S, Wheatley B. Geographical distribution of diabetes among the Native population of Canada: a national survey. *Soc Sci Med* 1990;31:129-39.
- Piolo M, Dyck RF, Gillis DC. Diabetes prevalence rates among First Nations adults on Saskatchewan reserves in 1990: comparison by tribal grouping, geography and with non-First Nation people. *Can J Public Health* 1996;87:325-8.
- Delisle HF, Rovard M, Ekoe JM. Prevalence estimates of diabetes and of other cardiovascular risk factors in the two largest Algonquin communities in Québec. *Diabetes Care* 1995;18:1255-9.
- Szathmary EJE, Ferrell RE. Glucose level, acculturation and glycosylated hemoglobin: an example of biocultural interaction. *Med Anthropol Q* 1990;4:315-41.
- James R, Young TK, Mustard CA. The health of Canadians with diabetes. *Health Rep* 1997;9(3):47-52.
- Delisle HF, Ekoé JM. Prevalence of non-insulin-dependent diabetes mellitus and impaired glucose tolerance in two Algonquin communities in Québec. *CMAJ* 1993;148(1):41-7.
- Harris SB, Gittelsohn J, Hanley A, Barnie A, Wolever TMS, Gao J, et al. The prevalence of NIDDM and associated risk factors in Native Canadians. *Diabetes Care* 1997;20:185-7.
- Dean HJ, Mundy RL, Moffatt M. Non-insulin-dependent diabetes mellitus in Indian children in Manitoba. *CMAJ* 1992;147(1):52-7.
- Harris SB, Perkins BA, Whalen-Brough E. Non-insulin-dependent diabetes mellitus among First Nations children. A new entity among First Nations people of north western Ontario. *Can Fam Physician* 1996;42:869-76.
- Dean HJ. NIDDM-Y in First Nations children in Canada. *Clin Pediatr* 1998; 37:898-96.
- Dean HJ, Young TK, Flett B, Wood-Steiman P. Screening for type-2 diabetes in aboriginal children in northern Canada. *Lancet* 1998;352:523-4.
- Chase LA. The trend of diabetes in Saskatchewan, 1905-1934. *CMAJ* 1937;36: 366-9.
- Fox C, Harris SB, Whalen-Brough E. Diabetes among Native Canadians in northwestern Ontario: 10 years later. *Chronic Dis Can* 1994;15:92-6.

34. Dow G. Diabetes mellitus and hospital utilization in the province of Manitoba [dissertation]. Winnipeg: University of Manitoba, 1991-92.
35. Macaulay AC, Montour LT, Adelson N. Prevalence of diabetic and atherosclerotic complications among Mohawk Indians of Kahnawake, PQ. *CMAJ* 1988; 139(3):221-4.
36. Brassard P, Robinson E. Factors associated with glycemia and microvascular complications among James Bay Cree Indian diabetics of Québec. *Arctic Med Res* 1995;54:116-24.
37. Young TK, Kaufert JM, McKenzie JK, Hawkins A, O'Neil JD. Excessive burden of end-stage renal disease among Canadian Indians: a national survey. *Am J Public Health* 1989;79:756-8.
38. Wilson R, Krefling LH, Sutcliffe P, VanBussel L. Incidence and prevalence of end-stage renal disease among Ontario's James Bay Cree. *Can J Public Health* 1992;83:143-6.
39. Dyck RF, Tan L. Rates and outcomes of diabetic endstage renal disease among registered native people in Saskatchewan. *CMAJ* 1994;150(2):203-8.
40. Ross SA, Fick GF. Insulin as a risk factor for diabetes complications [abstract]. *Diabetes* 1991;40(Suppl 1):333A.
41. Nicole LE, Friesen D, Harding GK, Roos LL. Hospitalization for acute pyelonephritis in Manitoba, Canada, during the period from 1989 to 1992; impact of diabetes, pregnancy, and aboriginal origin. *Clin Infect Dis* 1996;22:1051-6.
42. Hegele RA, Connelly PW, Scherer SW, Hanley AJ, Harris SB, Tsui LC, et al. Paraoxonase-2 gene (PON2) G148 variant associated with elevated fasting plasma glucose in noninsulin-dependent diabetes mellitus. *J Endocrinol Metab* 1997;82:3373-7.
43. Hegele RA, Harris SB, Zinman B, Wang J, Cao H, Hanley AJ, et al. Variation in the AU(AT)-rich element within the 3'-untranslated region of PP1R3 is associated with variation in plasma glucose in aboriginal Canadians. *J Endocrinol Metab* 1998;83:3980-3.
44. Hegele RA, Harris SB, Hanley AJ, Azouz H, Connelly PW, Zinman B. Absence of association between genetic variation of the beta 3-adrenergic receptor and metabolic phenotypes in Oji-Cree. *Diabetes Care* 1998;21:851-4.
45. Hegele RA, Sun F, Harris SB, Anderson C, Hanley AJ, Zinman B. Genome-wide scanning for type-2 diabetes susceptibility in Canadian Oji-Cree, using 190 microsatellite markers. *J Hum Genet* 1999;44:10-4.
46. Hegele RA, Cao H, Harris SB, Hanley AJ, Zinman B. The hepatic nuclear factor-1 alpha G319S variant is associated with early onset type-2 diabetes in Canadian Oji-Cree. *J Clin Endocrinol Metab* 1999;84:1077-82.
47. Szathmary EJE. The impact of low carbohydrate consumption on glucose tolerance, insulin concentration and insulin response to glucose challenge in Dogrib Indians. *Med Anthropol* 1989;11:329-50.
48. Szathmary EJE. The effect of Gc genotype on fasting insulin level in Dogrib Indians. *Hum Genet* 1987;75:368-72.
49. Ekoe JM, Thomas F, Balkau B, Eschwege E, Delisle H. Effect of maternal diabetes on the pattern of selected insulin resistance syndrome parameters in normal glucose tolerant subjects of two Algonquin Indian communities in Québec. *Diabetes Care* 1996;19:822-6.
50. Caulfield LE, Harris SB, Whalen EA, Sugamori ME. Maternal nutritional status, diabetes and risk of macrosomia among Native Canadian women. *Early Hum Dev* 1998;50:293-303.
51. Harris SB, Caulfield LE, Sugamori MS, Whalen EA, Henning B. The epidemiology of diabetes in pregnant Native Canadians. A risk profile. *Diabetes Care* 1997;20:1422-5.
52. Rodrigues S, Robinson E, Gray-Donald K. Prevalence of gestational diabetes among James Bay Cree women in northern Quebec. *CMAJ* 1999;160(9):1293-7. Abstract available: [www.cma.ca/cmaj/vol-160/issue-9/1293.htm](http://www.cma.ca/cmaj/vol-160/issue-9/1293.htm)
53. Godwin M, Muirhead M, Huynh J, Helt B, Grimmer J. Prevalence of gestational diabetes mellitus among Swampy Cree women in Moose Factory, James Bay. *CMAJ* 1999;160(9):1299-302. Abstract available: [www.cma.ca/cmaj/vol-160/issue-9/1299.htm](http://www.cma.ca/cmaj/vol-160/issue-9/1299.htm)
54. Young TK, Sevenhuysen GP, Ling N, Moffatt MEK. Determinants of plasma glucose level and diabetic status in a northern Canadian Indian population. *CMAJ* 1990;142(8):821-30.
55. Young TK. Obesity among Aboriginal peoples in North America: epidemiological patterns, risk factors and metabolic consequences. In: Angel A, Anderson H, Bouchard C, Lau D, Leiter L, Mendelson R, editors. *Progress in obesity research* 7. London: John Libby; 1996. p. 337-42.
56. Young TK, Sevenhuysen G. Obesity in northern Canadian Indians: patterns, determinants, and consequences. *Am J Clin Nutr* 1989;49:786-93.
57. Evers S, McCracken E, Antone I, Deagle G. Body fat distribution and non-insulin-dependent diabetes mellitus in North American Indians. *Nutrition Res* 1989;9:977-87.
58. Daniel M, Marion SA, Sheps SB, Hertzman C, Gamble D. Variation by body mass index and age in waist-to-hip ratio associations with glycemic status in an aboriginal population at risk for type 2 diabetes in British Columbia, Canada. *Am J Clin Nutr* 1999;69:455-60.
59. Hanley AJG, Harris SB, Gao XJ, Kwan J, Zinman B. Serum immunoreactive leptin concentrations in a Canadian Aboriginal population with high rates of NIDDM. *Diabetes Care* 1997;20:1408-15.
60. Wolever TMS, Hamad S, Gittelsohn J, Gao J, Hanley AJG, Harris SB, et al. Low dietary fiber and high protein intakes associated with newly diagnosed diabetes in a remote Aboriginal community. *Am J Clin Nutr* 1997;66:1470-4.
61. Gittelsohn J, Wolever TMS, Harris SB, Harris-Giraldo R, Hanley AJG, Zinman B. Specific patterns of food consumption and preparation are associated with diabetes and obesity in a Native Canadian community. *J Nutr* 1998;128:541-7.
62. Basualdo CG, Wein EE, Basu TK. Vitamin A (retinol) status of First Nation adults with non-insulin-dependent diabetes mellitus. *J Am Coll Nutr* 1997; 16:39-45.
63. Szathmary EJ, Ritenbaugh C, Goodby CS. Dietary change and plasma glucose levels in an Amerindian population undergoing cultural transition. *Soc Sci Med* 1987;24:791-804.
64. Dyck RF, Cassidy H. Preventing non-insulin-dependent diabetes among Aboriginal peoples: Is exercise the answer? *Chronic Dis Can* 1995;16:175-7.
65. Neel JV, Weder AB, Julius S. Type II diabetes, essential hypertension, and obesity as "syndromes of impaired genetic homeostasis": the "thrifty genotype" hypothesis enters the 21st century. *Perspect Biol Med* 1998;42:44-74.
66. Szathmary EJ. Diabetes in Amerindian populations: the Dogrib studies. In: Armelagos G, Swedlund A, editors. *Health and disease of populations in transition*. New York: Bergin and Garvey; 1990. p. 75-103.
67. Szathmary EJ. Non-insulin-dependent diabetes mellitus among Aboriginal North Americans. *Annu Rev Anthropol* 1994;23:457-82.
68. Ritenbaugh C, Goodby C. Beyond the thrifty gene: metabolic implications of prehistoric migration into the New World. *Med Anthropol* 1989;11:227-36.
69. Macaulay AC, Hanusaik N, Delisle DD. Diabetes education program in the Mohawk community of Kahnawake, Québec. *Can Fam Physician* 1988;34:1591-3.
70. Macaulay AC, Paradis G, Potvin L, Cross EJ, Saad-Haddad C, McComber A, et al. The Kahnawake Schools Diabetes Prevention Project: intervention, evaluation and baseline results of a diabetes primary prevention program with a Native community in Canada. *Prev Med* 1997;26:779-90.
71. Gittelsohn J, Harris SB, Whitehead S, Wolever TMS, Hanley JG, Barnie A, et al. Developing diabetes interventions in an Ojibwa-Cree community in northern Ontario: Linking qualitative and quantitative data. *Chronic Dis Can* 1995;16:157-64.
72. Gittelsohn J, Harris SB, Burris K, Kakegamic L, Landman LT, Sharma A, et al. Use of ethnographic methods for applied research on diabetes among the Ojibwa-Cree in northern Ontario. *Health Educ Q* 1996;23:365-82.
73. Daniel M, Gamble D, Henderson J, Burgess S. Diabetes prevalence, behavioural and anthropometric risk factors, and psychosocial constructs in three Aboriginal communities in central British Columbia. *Chronic Dis Can* 1995;16:165-74.
74. Daniel M, Green LW, Marion SA, Gamble D, Herbert CP, Hertzman C, et al. Effectiveness of community-directed diabetes prevention and control in a rural Aboriginal population in British Columbia, Canada. *Soc Sci Med* 1999;48:815-32.
75. Canadian Task Force on the Periodic Health Examination. *The Canadian guide to clinical preventive health care*. Ottawa: Health Canada; 1994. p. 602-9.
76. Meltzer S, Leiter L, Daneman D, Gerstein HC, Lau D, Ludwig S, et al, and the Steering and Expert Committees. 1998 clinical practice guidelines for the management of diabetes in Canada. *CMAJ* 1998;159(8 Suppl):S1-29.
77. Royal Commission on Aboriginal People. *Report of the Royal Commission on Aboriginal People*. Vol 3. *Gathering strength*. Ottawa: The Commission; 1996. p. 147.
78. Bobet E. *Diabetes among First Nations people*. Ottawa: Health Canada; 1998. Cat no H34-88-1998E.
79. Robinson EJ, Gebre Y, Pickering J, Petawabano B, Superville B, Lavallée C. Effect of bush living on Aboriginal Canadians of the Eastern James Bay region with non-insulin-dependent diabetes mellitus. *Chronic Dis Can* 1995;16:144-8.
80. Garro LC. Individual or societal responsibility? Explanations of diabetes in an Anishinaabe (Ojibway) community. *Soc Sci Med* 1995;40:37-46.
81. Garro LC. Intracultural variation in causal accounts of diabetes: a comparison of three Canadian Anishinaabe (Ojibway) communities. *Cult Med Psychiatry* 1996;20:381-420.
82. Hagey R. The phenomenon, the explanations and the responses: metaphors surrounding diabetes in urban Canadian Indians. *Soc Sci Med* 1984;18:265-72.
83. Ritenbaugh C. Obesity as a culture-bound syndrome. *Cult Med Psychiatry* 1982; 6:347-61.
84. Gittelsohn J, Harris SB, Thorne-Lyman AL, Hanley AJ, Barnie A, Zinman B. Body image concepts differ by age and sex in an Ojibway-Cree community in Canada. *J Nutr* 1996;126:2990-3000.
85. Boston P, Jordan S, McNamara E, Kozolanka K, Bobbish-Rondeau E, Iserhoff H, et al. Using participatory action research to understand the meanings aboriginal Canadians attribute to the rising incidence of diabetes. *Chronic Dis Can* 1997;18:5-12.
86. Travers KD. Using qualitative research to understand the sociocultural origins of diabetes among Cape Breton Mi'kmaq. *Chronic Dis Can* 1995;16:140-3.
87. Kuhnlein HV, Turner NJ. *Traditional plant foods of Canadian indigenous peoples: nutrition, botany and use*. Philadelphia: Gordon and Beach; 1991. p. 9-23.
88. Imes S, Wallace C, Jones S, Ryan EA. Herbal tea in the treatment of diabetes mellitus [abstract]. *Clin Invest Med* 1996;19:S19.

**Reprint requests to:** Dr. T. Kue Young, Department of Community Health Sciences, University of Manitoba, 750 Bannatyne Ave., Winnipeg MB R3E 0W3; fax 204 789-3905; [tkyoung@cc.umanitoba.ca](mailto:tkyoung@cc.umanitoba.ca)