

Table 2: Daily volumes of breast milk recorded by subjects in the domperidone and placebo groups

Patient	Study day; milk volume, mL						
	Baseline*	2	3	4	5	6	7
Domperidone							
1	13†	3	49	66	51	68	45
2	262	286	291	342	323	337	346
3	6	21	19	35	39	45	45
4	17	49	81	121	146	170	190
5	120	135	167	132	–	–	–
6	51	69	95	96	125	126	125
7	320	350	356	385	378	323	350
Mean volume	112.8	130.4	151.1	168.1	177.0	178.2	183.5
Placebo							
1	201†	152	180	102	133	80	110
2	12†	14	15	16	12	14	14
3	32†	36	36	32	36	40	40
4	14	37	62	55	42	57	58
5	11	28	18	8	4	0	0
6	50	92	82	120	121	150	180
7	89	102	94	111	95	101	107
8	23	21	22	22	–	–	–
9	< 1	10	20	25	27	15	19
Mean volume	48.2	54.7	58.8	54.6	58.8	57.1	66.1

*Defined as the volume of milk produced during the 24 hours before the start of the study medication.

†Baseline volume was unavailable; instead, volume produced within the 24 hours following enrolment was considered as the baseline.

[\[Return to text\]](#)