

Preventive health care, 2001 update: Should women be routinely taught breast self-examination to screen for breast cancer?

Nancy Baxter, with the Canadian Task Force on Preventive Health Care

Abstract

Objectives: To evaluate the evidence relating to the effectiveness of breast self-examination (BSE) to screen for breast cancer and to provide recommendations for routine teaching of BSE to women in various age groups as part of a periodic health examination.

Options: Routine teaching of BSE to women.

Evidence: The electronic databases MEDLINE, PreMEDLINE, CINAHL, HealthSTAR, Current Contents and the Cochrane Library were searched for abstracts and full reports of studies published from 1966 to October 2000 that evaluated the effectiveness of BSE in reducing breast cancer mortality. In addition, references of key papers were searched and experts consulted to ensure that all relevant articles had been identified.

Outcomes: Prevention of death from breast cancer was viewed as the most important outcome; other outcomes examined included the stage of cancer detected, the rate of benign biopsy results, the number of patient visits for breast complaints, and psychological benefits and harms.

Values: The recommendations of this report reflect the commitment of the Canadian Task Force on Preventive Health Care to provide a structured, evidence-based appraisal of whether a manoeuvre should be included in the periodic health examination.

Benefits, harms and costs: Breast cancer is the most frequently diagnosed cancer among Canadian women, accounting for 30% of all new cancer cases each year. In 2000 an estimated 19 200 Canadian women would have been diagnosed with breast cancer, and 5500 would have died from the disease. To date, 2 large randomized controlled trials, a quasi-randomized trial, a large cohort study and several case-control studies have failed to show a benefit for regular performance of BSE or BSE education, compared with no BSE. In contrast, there is good evidence of harm from BSE instruction, including significant increases in the number of physician visits for the evaluation of benign breast lesions and significantly higher rates of benign biopsy results.

Recommendations:

- *Women aged 40–49 years:* Because there is fair evidence of no benefit, and good evidence of harm, there is fair evidence to recommend that routine teaching of BSE be excluded from the periodic health examination of women in this age group (grade D recommendation).
- *Women aged 50–69 years:* Because there is fair evidence of no benefit, and good evidence of harm, there is fair evidence to recommend that routine teaching of BSE be excluded from the periodic health examination of women in this age group (grade D recommendation).
- The lack of sufficient evidence to evaluate the effectiveness of the manoeuvre in women younger than 40 years and those 70 years and older precludes making recommendations for teaching BSE to women in these age groups. The follow-

Research

Recherche

At the time of writing, Dr. Baxter was a resident in general surgery at the University of Toronto, Toronto, Ont.

A list of the task force members appears at the end of the article.

This article has been peer reviewed.

CMAJ 2001;164(13):1837-46

‡ See related article page 1851

[Return to June 26, 2001
Table of Contents](#)

ing issues may be important to consider: *Women younger than 40 years*: There is little evidence for effectiveness specific to this group. Because the incidence of breast cancer is low in this age group, the risk of net harm from BSE and BSE instruction is even more likely. *Women 70 years and older*: Although the incidence of breast cancer is high in this group, there is insufficient evidence to make a recommendation concerning BSE for women 70 years and older.

- **Important note:** Although the evidence indicates no benefit from routine instruction, some women will ask to be taught BSE. The potential benefits and harms should be discussed with the woman, and if BSE is taught, care must be taken to ensure she performs BSE in a proficient manner.

Validation: The findings of this analysis were reviewed through an iterative process by the members of the Canadian Task Force on Preventive Health Care. The task force sent the final review and recommendations to 4 independent experts, and their feedback was incorporated in the final draft of the manuscript.

Sponsors: The Canadian Task Force on Preventive Health Care is funded through a partnership between the Provincial and Territorial Ministries of Health and Health Canada.

In 1994 the Canadian Task Force on the Periodic Health Examination (now the Canadian Task Force on Preventive Health Care) concluded that there was insufficient evidence to recommend for or against breast cancer screening using breast self-examination (BSE) (grade C recommendation).¹ This review examines the evidence published since 1994 and makes recommendations about the value of routine teaching of BSE to women in various age groups.

For breast cancer screening in general, current evidence supports mammography and clinical breast examination for women aged 50–69 years.¹ For women aged 40–49, the evidence is insufficient to recommend for or against routine screening (although upon reaching the age of 40, Canadian women should be assisted in deciding at what age they wish to initiate mammography).² For women over 70, there is limited evidence regarding the benefit of screening.

Many breast tumours are found by women themselves,³ even in highly screened populations.⁴ However, in women regularly performing BSE, many self-detected tumours are found incidentally, not during self-examination.^{3–6} In one study, only 7.6% of women with breast tumours who were practising regular BSE actually detected the tumour by means of self-examination.³ In addition, tumours developing between screens in some age groups (e.g., 50–69) may be inherently more aggressive and thus may not be influenced by slightly earlier detection with BSE.

Burden of suffering

Breast cancer is the most frequently diagnosed cancer among Canadian women, accounting for 30% of all new cancer cases each year. In 2000 an estimated 19 200 Canadian women were diagnosed with the disease, and an estimated 5500 women died from it.⁷ The current age-standardized incidence and mortality rates for Canada are

106 and 27 per 100 000 respectively, with a lifetime risk of death of 1 in 25.8.⁷ Breast cancer is the leading cause of person-years of life lost for women.⁸ The most important risk factor for breast cancer in women is age;⁹ the disease is rare in women under 30.¹⁰ Because most women with breast cancer have no other identifiable risk factors,^{11–13} the effectiveness of teaching BSE should be demonstrated in the general population if it is to affect disease burden.

Methods

With the help of a reference librarian, the electronic databases MEDLINE, PreMEDLINE, CINAHL, HealthSTAR, Current Contents and the Cochrane Library were searched for articles published from 1966 to October 2000 using the following key words: “breast diseases,” “breast self-examination,” “palpation,” “mass screening” and “clinical trials.” The search was limited to English-language abstracts and studies involving humans. Related articles and references of key papers were searched and experts consulted to ensure that no important research was missed. Abstracts of all retrieved papers were read; those relevant to the review were critically appraised.

Prevention of death from breast cancer was viewed as the most important outcome; other outcomes included stage of cancer detected, benign biopsy rate, number of patient visits for breast complaints, and psychological benefits and harms.

The evidence was reviewed systematically using the methodology of the Canadian Task Force on Preventive Health Care.¹⁴ In brief, the principal author rated the quality of the evidence using the methodological hierarchy (Appendix 1). Preliminary drafts of the manuscript were circulated to the task force, comprising expert clinicians and methodologists, before it met in November 1999 and January 2000. A further draft incorporating feedback from 4 independent experts was finalized and endorsed by the task force. Procedures to achieve adequate documentation, consistency, comprehensiveness, objectivity and adherence to the task force’s methodology were maintained at all stages during review development, the consensus process and production of the final manuscript.

Results

Breast self-examination

BSE is a systematic method of self-inspection and palpation of the breast and axilla. There is no conclusive evidence regarding the most effective technique, the best teaching and reinforcement methods or the optimal frequency. Women's ability to detect lumps in silicone breast models has been found to correlate with the number of steps taken in the breast examinations,¹⁵⁻¹⁷ the length of the examination,^{15,18-20} the completeness of the search^{16,21} and the use of the pads of the fingers.^{17,20} Individual aspects of BSE associated with the greatest accuracy and long-term compliance have not been fully elucidated. The currently recommended frequency of once monthly has not been determined with scientific rigour.

In a review of interventions to enhance BSE practice,²² the most effective teaching methods were provision of information, demonstration with individual skills training, and feedback plus prompts such as mailed reminders. Although this review compares studies varying in type of participant and methodology, there is significant evidence that increasing the intervention strategy can improve compliance with BSE.

The overall sensitivity of BSE alone has been estimated to be 26% among screened women, assuming that all interval cancers were detected by BSE.^{23,24} Sensitivity varied with age, from 41% among women aged 35-39 to 21% among those aged 60-74. Specificity was not estimated. In studies of lump detection in silicone breast models, sensitivity ranged from 40% to 89%, and specificity from 66% to 81%.^{25,26} Although sensitivity improved with training, so did the rate of false-positive detection.¹⁹ Sensitivity with the silicone breast model was significantly lower among women over 60.²⁶

Effect of self-examination on breast cancer mortality

Details about studies that have evaluated the effectiveness of BSE in reducing breast cancer mortality are presented in Table 1 (randomized and quasi-randomized controlled trials²⁷⁻³⁴) and Table 2 (cohort and case-control studies³⁵⁻³⁹).

Randomized and quasi-randomized controlled trials

Two randomized controlled trials (RCTs) were conducted in populations with little concomitant screening.

Table 1: Summary of randomized and quasi-randomized controlled trials evaluating the effects of breast self-examination (BSE) on breast cancer outcomes

Trial	Participants	Follow-up/outcomes measured	Results	Strengths/potential biases
Shanghai RCT of BSE training ²⁷	Women aged 31-64; residents of Shanghai and current or former employees of Shanghai Textile Industry Bureau (STIB). Randomly assigned at factory level to BSE training group ($n = 133\ 375$) or control group ($n = 133\ 665$)	Follow-up: 5 yr Outcomes: breast cancer mortality (from STIB tumour and death registry), and follow-up of breast cancer cases, tumour stages and no. of benign lesions detected	No difference between groups in breast cancer mortality or stage of breast cancer; higher rate of benign biopsy results in BSE group than in control group (1.1% v. 0.5%)	Randomized trial, high participation and compliance rates; no concurrent screening programs. Inadequate length of follow-up (further follow-up underway). Political changes in China may affect ability to complete the study
Russian/WHO RCT of BSE training ²⁸⁻³²	Women aged 40-64; residents of Leningrad (St. Petersburg), randomly assigned at medical clinic level to BSE training group ($n = 57\ 712$ at 9 yr) or control group ($n = 64\ 759$ at 9 yr)	Follow-up: 9 yr Outcomes: breast cancer mortality among women diagnosed with breast cancer at oncology referral centre and medical clinic, tumour stages and rate of benign biopsy results	No difference between groups in breast cancer mortality or stage of breast cancer; higher rate of benign biopsy results in BSE group than in control group (at 5 yr: RR = 1.5, 95% CI 1.1-1.9)	Randomized trial, high participation rate; no concurrent screening. Decrease in compliance over time; inadequate power (further follow-up underway). Political changes in Russia may affect ability to complete the study
United Kingdom quasi-randomized controlled trial of breast cancer screening ³³⁻³⁴	Women aged 40-49 from 8 geographic areas assigned to following groups by centre: Screening by CBE and mammography (2 centres) ($n = 45\ 607$) BSE instruction (2 centres) ($n = 63\ 373$) Control group (4 centres) ($n = 127\ 123$)	Follow-up: mean 14.4 yr (98.2% of women traced) Outcome: breast cancer mortality (from tumour registry records), rate of benign biopsy results	No difference between groups in breast cancer mortality (RR = 0.99, 95% CI 0.87-1.12); no significant difference detected in secondary analysis by 5-yr age groups. Rate of benign biopsy results significantly higher in BSE group than in control group (0.91% v. 0.61%)	Differences across centres in demographics, rates of breast cancer, medical services, attendance at BSE instruction sessions (31% and 53% at 2 centres respectively) and breast cancer treatment patterns. No assessment of prevalent cancers, or BSE frequency or technique. Last years of follow-up overlapped with large national breast cancer screening study

Note: RCT = randomized controlled trial, WHO = World Health Organization, CBE = clinical breast examination, RR = relative risk, CI = confidence interval.

One trial involved 267 040 women aged 31–64 recruited from 520 factories in Shanghai, China.²⁷ The 133 375 women in the intervention group received extensive BSE training using silicone breast models and personalized instruction, followed by 2 reinforcement sessions and multiple reminders to practise the technique. Women were followed up for the development of breast cancer and death, confirmed by registry data. If the follow-up of breast cancer deaths continues through a 10-year period, the power

to detect a 30% reduction in breast cancer mortality will be 80%. After the first 5 years of follow-up, the cumulative breast cancer mortality rate was not significantly lower among women who received BSE instruction than among the control subjects (30.9 v. 32.7 per 100 000 woman-years). Similar numbers of breast tumours were detected in both groups (331 in the BSE group and 322 in the control group). No consistent, large or significant differences in tumour size or stage were found between the 2

Table 2: Summary of cohort and case-control studies evaluating BSE as a screening technique for breast cancer

Study	Participants	Exposure	Results	Strengths/potential biases
Holmberg et al ³⁵ (cohort study)	548 000 women from 26 US states recruited by volunteers in 1959 for large population-based study; 93% followed to 1973. For those who died, copies of death certificates were obtained to determine cause of death	Self-reported interview data from 1959 served as baseline. Women divided into those performing BSE ($n = 177\ 602$) and those not ($n = 272\ 554$)	No difference in breast cancer mortality between groups based on BSE performance reported in 1959, even after adjustments for covariates	Large, prospective study. Poor measure of BSE performance; no objective assessment or evaluation of technique. Cause of death on death certificate accepted as cause of death
Harvey et al ³⁶ (case-control study nested in Canadian NBSS RCT of breast cancer screening)	Women aged 40–59 without breast cancer recruited in 1980–1985. <i>Cases:</i> 163 women who died of breast cancer and 57 with distant metastases; cases had to have been diagnosed 2 yr after study entry <i>Controls:</i> 10 per case randomly selected from NBSS and matched by 5-yr age group, screening centre and year of enrolment, and by randomized group in NBSS study	All participants given BSE instruction at entry. Annual measures of BSE frequency. Objective measure of BSE technique for subgroup analysis	Similar proportions of cases and controls practised BSE before study entry ($p = 0.10$). Frequency of BSE not associated with case status. Omission of specific components of BSE technique associated with increased risk of death from breast cancer among cases: for those who omitted 3 key manoeuvres at 2 yr before diagnosis, the OR was 2.95 (95% CI 1.19–7.30; $p = 0.02$); no difference between groups at 1 or 3 yr	Study designed to reduce recall, lead-time and length biases. Controls selected from same population as larger study. Objective measures of BSE performance and frequency. Multiple comparisons in post hoc analysis without statistical adjustment. High compliance with BSE in trial; thus, compliers with BSE may differ from noncompliers in other ways that affect breast cancer mortality
Muscat et al ³⁷	<i>Cases:</i> 453 women with newly diagnosed advanced breast cancer (axillary or distant metastases, or tumour > 2 cm in diameter) <i>Controls:</i> 887 age-matched women randomly selected from population	Self-reported BSE frequency before study entry	No difference in BSE frequency between groups	Study designed to avoid lead-time bias. Retrospective interview
Newcomb et al ³⁸	<i>Cases:</i> 209 women with advanced breast cancer (TNM stage III or IV) <i>Controls:</i> 433 age-matched women without advanced breast cancer	Self-reported BSE practice	No difference in BSE frequency between groups. Compared with cases, controls were more proficient at BSE (OR 0.54, $p = 0.003$) and more likely to have had clinical breast examination within 5 yr of reference date*	Study designed to avoid lead-time bias. Retrospective interview, self-reported BSE practice without confirmation, no standardized measure of proficiency. Multiple comparisons with no statistical adjustment. No adjustment in analysis for CBE
Gastrin et al ³⁹	56 177 Finnish women; 29 018 women returned BSE compliance calendars and were linked to the Finnish Cancer Registry	All women were enrolled in BSE information session and mammography self-referral system and asked to complete BSE compliance calendars annually	Lower breast cancer mortality (RR = 0.75) among women returning calendars than among women in general female population (excluding women diagnosed with breast cancer within the first 2 yr of the program)	Population-controlled cohort study. Designed to avoid lead-time bias, but not length or over-diagnosis bias. Appeared to be significant selection bias (twice as many subjects in cohort as in general population attended college)

Note: NBSS = National Breast Screening Study, OR = odds ratio, TNM = tumour-node-metastasis.

*Date of breast cancer diagnosis in a case subject was used as the reference date for both the case and matched control subject.

groups. A random sample of women from each group was tested for their ability to detect lumps in silicone breast models. Women from the BSE group found more lumps and demonstrated greater specificity than did those from the control group. Competency of BSE was assessed in subgroups of the intervention group and was judged to be good, with more than 90% of the women correctly palpating most of the breast even 1 year after reinforcement of BSE technique. The study's large size and its high participation and compliance rates are strengths. Although the methods used to classify tumours by size and stage may have been nonstandardized, any misclassification would not bias mortality outcomes; however, they could obscure any real differences between the groups for other comparisons.

The other RCT was conducted in St. Petersburg, Russia, and involved women aged 40–64 randomly assigned to BSE educational programs or to a control group.^{28–32} Five-year and 9-year follow-up data were published for 122 471 women;^{30–32} the 9-year follow-up report was published in Russian, and an English translation was obtained from the authors. Women in the BSE arm attended small-group sessions, run by trained nurses or physicians, that involved a demonstration of the BSE technique. Among 400 randomly selected women, the self-reported compliance rate for BSE 5 or more times per year was 82% at 1 year but decreased to 55.8% by 5 years. Medical personnel assessed the quality of BSE performed by this subset of women as good. After 9 years of follow-up 493 of the 57 712 women in the BSE group and 446 of the 64 759 women in the control group had breast cancer. There was no difference in tumour stage between the BSE and control groups, nor was there a difference in breast cancer mortality (0.27% [$n = 157$] and 0.26% [$n = 167$] respectively). Although a well-developed cancer registry was lacking in St. Petersburg, there is no evidence that any failure to find all cases would have biased the study in either direction. Inadequate information about the control group could theoretically mask a high rate of BSE (although unlikely in this population) and decrease the ability of the study to detect a benefit in the intervention group. Because of lower-than-expected compliance and cancer incidence,⁴⁰ there was inadequate power to detect a 20% difference in mortality between the 2 groups, and the study was extended until the year 2001. Although this study is unable to answer the question of the effect of mass teaching of BSE on mortality at this time, the failure to find any difference between the 2 groups in breast cancer stage decreases the likelihood of a positive study.

A quasi-randomized trial in the United Kingdom in a large population of women aged 45–64 years at 8 centres with little concomitant screening^{33,34} showed no reduction in breast cancer mortality between the BSE and control centres after 16 years of follow-up. However, significant differences existed between the centres in methods of recruitment for BSE teaching, breast cancer care and the pre-

trial breast cancer mortality rate, all of which may have confounded the results of the study. A case–control study at one of the centres demonstrated a significant reduction in deaths from breast cancer in those attending BSE classes;⁴¹ no adjustment for confounders was attempted. Because the same data were reported in the quasi-randomized controlled trial, a study of stronger design, the case–control study was not considered separately.

Cohort and case–control studies

A prospective cohort study in the United States involving 450 156 women over age 30 demonstrated no benefit of self-reported BSE practice in preventing death from breast cancer, regardless of age.³⁵ Self-reporting of BSE and unverified data from death certificates were potential limitations. The strengths of the study included its large sample, and the other breast cancer risk factors measured in the study significantly influenced risk of breast cancer mortality in the expected direction.

Three well-performed case–control studies evaluated the effectiveness of BSE.^{36–38} No difference in self-reported frequency was found between the cases and controls in any of these studies. One study showed that, although women who reported examining their underarms were at low risk of advanced breast cancer (odds ratio 0.48, 95% confidence interval [CI] 0.33–0.69), women who reported examining their nipples appeared to be at increased risk (relative risk 1.48, 95% CI 1.05–2.09). The fact that more women in the control group than in the case group had clinical breast examinations in the 5 years before the reference date was not controlled for in the analysis.³⁸ (The date of cancer diagnosis in a case subject was used as the reference date for both the case and matched control subject.)

The case–control study nested within the Canadian National Breast Screening Study eliminated lead-time, length and recall bias through prospective data collection.³⁶ No difference was found in self-reported BSE frequency between the case and control subjects. In a subgroup analysis, the performance of some components of BSE was found to differ between case and control subjects; however, these components were not selected a priori but based on a regression analysis, which increased the chance that the results may have been confounded.³⁶ Of note, one component of BSE in this study, the use of circular palpation, was associated with an increased risk of death from breast cancer or metastatic disease at 1 year preceding diagnosis. Women in this study were highly compliant with BSE, with over 74% usually performing visual examination, 71% using 3 digits for examination and 70% using the finger pads by the fifth year of the study.³² The possibility that compliers with these components of BSE differed significantly from noncompliers in ways other than BSE performance must therefore be entertained and may explain the results found.

Other studies

A Finnish cohort study³⁹ involving 56 177 women reported a breast cancer mortality ratio of 0.75 in the BSE group, lower than that in the general female population. However, methodological problems with this study, particularly selection bias, make it impossible to draw firm conclusions. A large number of cross-sectional studies examined the effect of BSE on various measures of cancer stage and survival.^{39,43-62} Because these have shown mixed results and are subject to lead-time, length and selection bias, they were not included in this analysis.

Effect of proficiency on breast cancer mortality

In 2 of the case-control studies described in Table 2^{36,38} and in an observational study,⁴⁹ secondary analyses were used to evaluate the effect of BSE technique. In the study by Newcomb and associates,³⁸ although BSE performance was not effective overall, in subgroup analysis more thorough BSE was associated with a decreased risk of advanced breast cancer.³⁸ However, no statistical adjustment was made for multiple comparisons. Also, subjects highly compliant with thorough BSE may differ significantly from noncompliant subjects in other ways (e.g., age,^{16,20,21,63} health motivation,⁶⁴ marital status,¹⁷ educational level,¹⁷ perceived barriers to BSE⁶⁴ and perceived susceptibility or family history of breast cancer⁶⁴), which may influence breast cancer mortality. In the secondary analyses, the effectiveness of some individual components of BSE was also evaluated, with conflicting results. Because of potential selection and other biases, and susceptibility to type I error, it is impossible to draw conclusions about BSE proficiency based on results of such studies.

Potential harms

Although considered a relatively minor procedure, breast biopsy causes permanent scarring and may cause significant breast deformity and emotional distress.⁶⁵⁻⁶⁷ In the UK trial the rate of benign breast biopsy results was significantly higher in the centres assigned to BSE than in the control centres (0.91% v. 0.61%).⁶⁸ Differences in benign biopsy rate were reported between the 2 BSE sites (0.71% and 1.28%); however, both sites had higher rates than those in the control sites.⁶⁸

The Russian trial reported a significant difference in the proportion of women who presented for assessment of signs or symptoms of breast disease (7.5% in the BSE group v. 3.8% in the control group at 9 years).³² After 5 years, there was a significantly higher rate of benign biopsy results among the BSE-trained women than among the control subjects (0.21% v. 0.14% respectively; relative risk 1.5; 95% CI 1.1-1.9). In addition, the benign needle biopsy rate was higher in the BSE group than in the control group (0.57% v. 0.32%). No significant dif-

ference in the rate of malignant breast biopsy results was found between the groups.²⁹ In the Shanghai trial, although the rate of benign breast biopsy results was not provided, many more benign lesions were detected by women in the BSE group than in the control group (1.09% v. 0.47%), with a nearly equal number of breast tumours diagnosed.²⁷

Negative psychological impacts related to BSE training include increased levels of worrying, anxiety and depression,^{15,69} and increased numbers of follow-up investigations because of false-positive findings.^{65,70,71}

No published studies provided a clear breakdown by age of the outcomes of investigating positive BSE findings. However, because the incidence of breast cancer is low among women under 30 and breast lumps are commonly benign in that age group, it would be expected that the false-positive rate would be much higher among younger than among older women.⁷²

Costs

Teaching and reinforcing BSE are costly activities and potentially divert resources from other preventive strategies. Although no Canadian data exist, a US cost-effectiveness analysis of 2 nursing-led BSE education programs estimated the cost to be between US\$574 and US\$848 per competent frequent self-examiner added.⁷³ A well-conducted trial showing no evidence of efficacy would justify utilization of limited funds and resources for other purposes. Some evidence exists about the effect of BSE programs on health care utilization: one study indicated that a hospital management organization (HMO)-based BSE program may increase utilization of health services by all women,⁷⁴ and another study showed that, 1 year after training, women who entered a trial of BSE education had similar health care use to those in the control groups.¹⁵

Summary

There were 7 studies that evaluated the effectiveness of BSE in preventing death from breast cancer. Although neither trial has reached maturity, 2 large RCTs have shown to date no difference in breast cancer mortality, or stage at diagnosis, between women taught BSE and control subjects.^{27,32} Although it is important that both trials continue to completion, these current findings make the possibility of a large impact of BSE on mortality unlikely. Political and social changes in China and Russia may jeopardize the completion of both studies.

After 16 years of follow-up in the UK quasi-randomized trial, no reduction in breast cancer mortality was found between the BSE and control centres. However, the significant differences between the centres in methods of recruitment for BSE teaching, breast cancer care and pretrial mortality may have confounded the results.³⁴

A large US cohort study of women over 30 demonstrated no effect of self-reported BSE practice on breast cancer mortality. No confirmation of BSE performance or adequacy was attempted, and this may have biased the study toward the null hypothesis.³⁵ Three case-control studies were properly performed to evaluate the effectiveness of BSE.³⁶⁻³⁸ Although they may have shown a benefit from more thorough BSE, no statistical adjustment was made for multiple comparisons. Also, because subjects highly compliant with thorough BSE may differ significantly from noncompliant subjects in other ways that might affect breast cancer mortality, this result should not be given undue weight. In summary, no appropriately designed study has shown a benefit from regular performance of BSE or BSE education over no BSE.

In contrast, there is good evidence of harm from BSE instruction. Both RCTs demonstrated a significant increase in the number of physician visits for the evaluation of benign breast lesions in the groups assigned to BSE education. The Russian study reported a benign breast biopsy rate that was significantly higher in the group assigned to BSE instruction than in the control group, whereas the number of biopsies showing cancer was virtually identical between the 2 groups. In the UK trial, the benign biopsy rate was also significantly higher in the centres assigned to BSE education than in the control centres.

To conclude, many breast tumours are discovered by women themselves, and although the evidence does not support routinely teaching BSE, women should be instructed to promptly report any breast changes or concerns. Any woman who wishes to practise BSE and who requests instruction should be counselled regarding the risks and benefits, and the health care professional should ensure that BSE instruction is thorough and that the woman is performing BSE in a proficient manner.

Recommendations

By the Canadian Task Force on Preventive Health Care

The recommendations for routine teaching of BSE to women in various age groups are summarized in Table 3.

Women aged 40-69 years

Women aged 40-49 years: In this age group, mammography and clinical breast examination are not of clear benefit, and studies evaluating the effectiveness of BSE have shown no benefit. In addition, the evidence for increased risk of benign biopsy and the lower incidence of breast cancer in this younger group indicates an increased potential for net harm. Thus, there is fair evidence to recommend that routine teaching of BSE be excluded from the periodic health examination of women aged 40-49 (grade D recommendation).

Women aged 50-69: There is fair evidence to conclude that screening with BSE is not effective in reducing breast cancer mortality in populations not receiving mammography or routine clinical breast examinations. Because mammography and clinical breast examinations reduce breast cancer mortality, they are currently recommended in Canada for this age group. It is highly unlikely that screening with BSE would confer additional benefits to screening of proven efficacy. In addition, there is good evidence that BSE education increases the number of physician visits for assessment of breast complaints and the rate of biopsies of benign breast lesions. Because there is good evidence of harm and fair evidence of no benefit, there is fair evidence to recommend that routine teaching of BSE be excluded from the periodic health examination of women aged 50-69 (grade D recommendation).

Table 3: Summary table of recommendations for the routine teaching of BSE to women

Manoeuvre	Effectiveness	Levels of evidence*	Recommendation*
Routine teaching of BSE to women aged 40-49 yr	Evidence of no benefit in terms of survival from breast cancer	RCTs (I), ^{27,31,32} nonrandomized trial (II-1), ³⁴ cohort study (II-3), ³⁵ case-control studies (II-3) ³⁶⁻³⁸	Because there is fair evidence of no benefit, and good evidence of harm, there is fair evidence to recommend that routine teaching of BSE be excluded from the periodic health examination of women aged 40-49 (grade D†)
	Evidence of increased no. of physician visits for breast problems and increased rate of benign biopsy results	RCTs (I), ^{27,31,32} nonrandomized trial (II-1) ³⁴	
Routine teaching of BSE to women aged 50-69 yr	Evidence of no benefit in terms of survival from breast cancer	RCTs (I), ^{27,31,32} nonrandomized trial (II-1), ³⁴ cohort study (II-3), ³⁵ case-control studies (II-3) ³⁶⁻³⁸	Because there is fair evidence of no benefit, and good evidence of harm, there is fair evidence to recommend that routine teaching of BSE be excluded from the periodic health examination of women aged 50-69 (grade D†)
	Evidence of increased no. of physician visits for breast problems and increased rate of benign biopsy results	RCTs (I), ^{27,31,32} nonrandomized trial (II-1) ³⁴	

*See Appendix 1 for definitions of the levels of evidence and grades of recommendations.

†Although the evidence indicates no benefit from routine instruction, some women will ask to be taught BSE. The potential benefits and harms should be discussed with the woman, and if BSE is taught, care must be taken to ensure that she performs BSE in a proficient manner.

Note: The lack of sufficient evidence to evaluate the effectiveness of the manoeuvre in women younger than 40 years and those 70 years and older precludes making recommendations for teaching BSE to women in these age groups. The following issues may be important to consider: *Women younger than 40 years:* There is little evidence for effectiveness specific to this group. Because the incidence of breast cancer is low in this age group, the risk of net harm from BSE and BSE instruction is even more likely. *Women 70 years and older:* Although the incidence of breast cancer is high in this group, there is insufficient evidence to make a recommendation concerning BSE for women 70 years and older.

Women younger than 40 and those 70 years and older

The lack of sufficient evidence to evaluate the effectiveness of the manoeuvre in women younger than 40 and those 70 years and older precludes making recommendations for teaching BSE to women in these age groups. The following issues may be important to consider.

Women under 40 years: There is little evidence for the effectiveness of routinely teaching BSE that is specific to this age group. Because the incidence of breast cancer is low in this age group, the risk of net harm from BSE and BSE instruction is even more likely.

Women 70 years and older: Although the incidence of breast cancer is high in this age group, there is insufficient evidence to make a recommendation concerning BSE for women in this age group.

Important note

Although the evidence indicates no benefit from routine instruction, some women will ask to be taught BSE. The potential benefits and harms should be discussed with the woman, and if BSE is taught, care must be taken to ensure that she performs the examinations in a proficient manner.

By other organizations

Organizations that currently recommend the routine teaching of BSE include the American Academy of Family Physicians,⁶³ the American College of Obstetricians and Gynecologists,⁶⁴ the American Cancer Society⁷⁵ and the Canadian Cancer Society.⁷⁶

In 1993 the Evaluation Committee of the European Society of Mastology concluded that there was no benefit from routine BSE and no support for public health programs for BSE education.⁷⁷

The US Preventive Services Task Force found insufficient evidence to recommend for or against teaching BSE and insufficient evidence to recommend changing current BSE practices.⁷⁸ This recommendation is currently under review.

Research agenda

The RCTs of BSE effectiveness should be continued until adequate power is achieved. A well-designed trial of BSE effectiveness in a population receiving screening techniques of proven effectiveness (mammography and clinical breast examination for women 50–69 years) would be more applicable to the Canadian population but is unlikely to occur. Research into BSE technique may be useful. Ways of minimizing the detection of benign lesions should also be examined. There is a need to determine the extent to which older women (70 years and older) are likely to benefit from any screening programs for breast cancer.

This paper is based on the full technical report available from the office of the Canadian Task Force on Preventive Health Care (see the contact information at the end of the bibliography).

Competing interests: None declared.

Contributors: Dr. Baxter, as primary author, was responsible for all aspects related to initiating the topic, reviewing the evidence, drafting the original manuscript for review by the Canadian Task Force on Preventive Health Care and making subsequent revisions following input from the task force and external reviewers. The task force had 2 main roles: first, each task force member thoroughly reviewed the draft systematic review at several meetings, provided feedback to the author on the evidence review and deliberated with the author to arrive at practice recommendations; second, the task force office managed the committee analytical process, arranged external expert review and supported the author in preparing literature searches, drafting tables or other supplementary materials, and revising the various drafts of the manuscript.

Acknowledgements: The Canadian Task Force on Preventive Health Care thanks Dr. Suzanne W. Fletcher and Larissa Nekhlyudov, Department of Ambulatory Care and Prevention, Harvard Medical School, Boston, Mass., Dr. Russell Harris, University of North Carolina at Chapel Hill, Chapel Hill, NC, and Dr. Kathleen I. Pritchard, Toronto–Sunnybrook Regional Cancer Centre, Toronto, Ont., for reviewing an earlier draft of this manuscript. The views expressed in this report are those of the author and the task force and do not necessarily reflect the positions of reviewers.

References

- Morrison B. Screening for breast cancer. In: Canadian Task Force on Preventive Health Care. *The Canadian guide to clinical preventive health care*. Ottawa: Canada Communication Group; 1994. p. 787–95.
- Ringash J, with the Canadian Task Force on Preventive Health Care. Preventive health care, 2001 update: screening mammography among women aged 40–49 years at average risk of breast cancer. *CMAJ* 2001;164(4):469–76. Available: www.cma.ca/cmaj/vol-164/issue-4/0469.htm
- Auvainen A, Elovainio L, Hakama M. Breast self-examination and survival from breast cancer: a prospective follow up study. *Breast Cancer Res Treat* 1996;38(2):161–8.
- Miller AB, Baines CJ, To T, Wall C. Canadian National Breast Screening Study: 2. Breast cancer detection and death rates among women aged 50–59 years. *CMAJ* 1992;147:1477–88.
- McPherson CP, Swenson KK, Jolitz G, Murray CL. Survival of women ages 40–49 years with breast carcinoma according to method of detection. *Cancer* 1997;79:1923–32.
- Huguley CM, Brown RL, Greenberg RS, Clark WS. Breast self-examination and survival from breast cancer. *Cancer* 1988;62:1389–96.
- Canadian Cancer Statistics 2000*. Toronto: National Cancer Institute of Canada; 2000.
- Law MR, Morris JK, Wald NJ. The importance of age in screening for cancer. *J Med Screen* 1999;6:16–20.
- Kelsey JL, Bernstein L. Epidemiology and prevention of breast cancer. *Annu Rev Public Health* 1996;17:47–67.
- Forbes JF. The incidence of breast cancer: the global burden, public health considerations. *Semin Oncol* 1997;24:S1–20–S1–35.
- Vafiadis P. Breast self examination: Should general practice bother? *Aust Fam Physician* 1997;26(Suppl 1):S41–6.
- Henderson IC. Risk factors for breast cancer development. *Cancer* 1993;71(6 Suppl):2127–40.
- Mant D, Vessey MP. Epidemiology of breast cancer. In: Kirby I, Copeland EM, editors. *The breast: comprehensive management of benign and malignant diseases*. Philadelphia: WB Saunders; 1991;116:363–5.
- Woolf SH, Battista RN, Anderson GM, Logan AG, Wang E. Assessing the clinical effectiveness of preventive maneuvers: analytic principles and systematic methods in reviewing evidence and developing clinical practice recommendations. A report by the Canadian Task Force on the Periodic Health Examination. *J Clin Epidemiol* 1990;43(9):891–905.
- Fletcher SW, O'Malley MS, Earp JL, Morgan TM, Lin S, Degnan D. How best to teach women breast self-examination. A randomized controlled trial. *Ann Intern Med* 1990;112:772–9.
- Dorsay RH, Cuneo WD, Somkin CP, Tekawa IS. Breast self-examination: improving competence and frequency in a classroom setting. *Am J Public Health* 1988;78:520–2.
- Howe HL. Proficiency in performing breast self-examination. *Patient Couns Health Educ* 1980;4:151–3.
- Hall DC, Adams CK, Stein GH, Stephenson HS, Goldstein MK, Penny-packer HS. Improved detection of human breast lesions following experimental training. *Cancer* 1980;46:408–14.
- Bennett SE, Lawrence RS, Angiolillo DF, Bennett SD, Budman S, Schneider

- GM, et al. Effectiveness of methods used to teach breast self-examination. *Am J Prev Med* 1990;6:208-17.
20. Haughey BP, Marshall JR, Nemoto T, Krolbart K, Mettlin C, Swanson M. Breast self-examination: reported practices, proficiency, and stage of disease at diagnosis. *Oncol Nurs Forum* 1988;15:315-9.
 21. Rutledge DN. Sensitivity and specificity of lump detection in breast models. *Am J Prev Med* 1992;8:314-8.
 22. Janz NK, Becker MH, Anderson LA, Marcoux BC. Interventions to enhance breast self-examination practice: a review. *Public Health Rev* 1989-90;17:89-163.
 23. Woolf SH. United States Preventive Services Task Force recommendations on breast cancer screening. *Cancer* 1992;69(7 Suppl):1913-8.
 24. Baker LH. Breast Cancer Detection Demonstration Project: five-year summary report. *CA Cancer J Clin* 1982;32(4):194-225.
 25. Fletcher SW, O'Malley MS, Pilgrim CA, Gonzalez JJ. How do women compare with internal medicine residents in breast lump detection? A study with silicone models. *J Gen Intern Med* 1989;4:277-83.
 26. Rutledge DN. Effects of age on lump detection accuracy. *Nurs Res* 1992;41:306-8.
 27. Thomas DB, Gao DL, Self SG, Allison CJ, Tao Y, Mahloch J, et al. Randomized trial of breast self-examination in Shanghai: methodology and preliminary results. *J Natl Cancer Inst* 1997;89:355-65.
 28. Semiglazov VF, Moiseenko VM. Breast self-examination for the early detection of breast cancer: a USSR/WHO controlled trial in Leningrad. *Bull World Health Organ* 1987;65:391-6.
 29. Semiglazov VF, Moiseyenko VM, Bavli JL, Migmanova NS, Seleznyov NK, Popova RT, et al. The role of breast self-examination in early breast cancer detection (results of the 5-year USSR/WHO randomized study in Leningrad). *Eur J Epidemiol* 1992;8:498-502.
 30. Semiglazov VF, Sagaidak VN, Moiseyenko VM, Mikhailov EA. Study of the role of breast self-examination in the reduction of mortality from breast cancer. The Russian Federation/World Health Organization Study. *Eur J Cancer* 1993;29A:2039-46.
 31. Semiglazov VF, Moiseenko VM, Protsenko SA, Bavli IL, Orlov AA, Ivanova OA, et al. [Preliminary results of the Russia (St.Petersburg)/WHO program for the evaluation of the effectiveness of breast self-examination]. *Vopr Onkol* 1996;42:49-55.
 32. Semiglazov VF, Moiseenko VM, Manikhas AG, Protsenko SA, Kharikova RS, Popova RT, et al. [Interim results of a prospective randomized study of self-examination for early detection of breast cancer (Russia/St.Petersburg/WHO)]. *Vopr Onkol* 1999;45:265-71.
 33. UK Trial of Early Detection of Breast Cancer Group. First results on mortality reduction in the UK trial of early detection of breast cancer. *Lancet* 1988;2:411-6.
 34. UK Trial of Early Detection of Breast Cancer Group. 16-year mortality from breast cancer in the UK Trial of Early Detection of Breast Cancer. *Lancet* 1999;353:1909-14.
 35. Holmberg L, Ekblom A, Calle E, Mokdad A, Byers T. Breast cancer mortality in relation to self-reported use of breast self-examination. A cohort study of 450,000 women. *Breast Cancer Res Treat* 1997;43:137-40.
 36. Harvey BJ, Miller AB, Baines CJ, Corey PN. Effect of breast self-examination techniques on the risk of death from breast cancer. *CMAJ* 1997;157(9):1205-12. Available: www.cma.ca/cmaj/vol-157/issue-9/1205.htm
 37. Muscat JE, Huncharek MS. Breast self-examination and extent of disease: a population-based study. *Cancer Detect Prev* 1991;15:155-9.
 38. Newcomb PA, Weiss NS, Storer BE, Scholes D, Young BE, Voigt LF. Breast self-examination in relation to the occurrence of advanced breast cancer. *J Natl Cancer Inst* 1991;83:260-5.
 39. Gastrin G, Miller AB, To T, Aronson KJ, Wall C, Hakama M, et al. Incidence and mortality from breast cancer in the Mama Program for Breast Screening in Finland, 1973-1986. *Cancer* 1994;73:2168-74.
 40. Koroltschouk V, Stanley D, Stjernsward J. The control of breast cancer: a World Health Organization perspective. *Cancer* 1990;65:2803-10.
 41. Locker AP, Caseldine J, Mitchell A, Blamey RW, Roebuck EJ, Elston CW. Results from a seven-year programme of breast self-examination in 89,010 women. *Br J Cancer* 1989;60:401-5.
 42. Baines CJ, To T. Changes in breast self-examination behavior achieved by 89,835 participants in the Canadian National Breast Screening Study. *Cancer* 1990;66:570-6.
 43. Mant D, Vessey MP, Neil A, McPherson K, Jones L. Breast self examination and breast cancer stage at diagnosis. *Br J Cancer* 1987;55:207-11.
 44. Huguley CM, Brown RL. The value of breast self-examination. *Cancer* 1981;47:989-95.
 45. Foster RS, Lang SP, Costanza MC, Worden JK, Haines CR, Yates JW. Breast self-examination practices and breast cancer stage. *N Engl J Med* 1978;299:265-70.
 46. Tamburini M, Massara G, Bertario L, Re A, Di Pietro S. Usefulness of breast self-examination for an early detection of breast cancer. Results of a study on 500 breast cancer patients and 652 controls. *Tumori* 1981;67:219-24.
 47. Smith EM, Francis AM, Polissar L. The effect of breast self-exam practices and physician examinations on extent of disease at diagnosis. *Prev Med* 1980;9:409-17.
 48. Smith EM, Burns TL. The effects of breast self-examination in a population-based cancer registry. A report of differences in extent of disease. *Cancer* 1985;55:432-7.
 49. Hislop TG, Coldman AJ, Skippen DH. Breast self-examination: importance of technique in early diagnosis. *CMAJ* 1984;131:1349-52.
 50. Senie RT, Rosen PP, Lesser ML, Kinne DW. Breast self-examination and medical examination related to breast cancer stage. *Am J Public Health* 1981;71:583-90.
 51. Feldman JG, Carter AC, Nicastrì AD, Hosat ST. Breast self-examination, relationship to stage of breast cancer at diagnosis. *Cancer* 1981;47:2740-5.
 52. Gould-Martin K, Paganini-Hill A, Casagrande C, Mack T, Ross RK. Behavioral and biological determinants of surgical stage of breast cancer. *Prev Med* 1982;11:429-40.
 53. Owen WL, Hoge AF, Asal NR, Anderson PL, Cucchiara AJ. Self-examination of the breast: use and effectiveness. *South Med J* 1985;78:1170-3.
 54. Greenwald P, Nasca PC, Lawrence CE, Horton J, McGarrath RP, Gabriele T, et al. Estimated effect of breast self-examination and routine physician examinations on breast-cancer mortality. *N Engl J Med* 1978;299:271-3.
 55. Shugg D, Hill D, Cooper D, Shepherd A. Practice of breast self-examination and the treatment of primary breast cancer. *Aust N Z J Surg* 1990;60:455-62.
 56. GIVIO (Interdisciplinary Group for Cancer Care Evaluation). Practice of breast self examination: disease extent at diagnosis and patterns of surgical care. A report from an Italian study. *J Epidemiol Community Health* 1991;45:112-6.
 57. Foster RS, Costanza MC. Breast self-examination practices and breast cancer survival. *Cancer* 1984;53:999-1005.
 58. Le Geyte M, Mant D, Vessey MP, Jones L, Yudin P. Breast self examination and survival from breast cancer. *Br J Cancer* 1992;66:917-8.
 59. Ogawa H, Tominaga S, Yoshida M, Kubo K, Takeuchi S. Breast self-examination practice and clinical stage of breast cancer. *Jpn J Cancer Res* 1987;78:447-52.
 60. Kuroishi T, Tominaga S, Ota J, Horino T, Taguchi T, Ishida T, et al. The effect of breast self-examination on early detection and survival. *Jpn J Cancer Res* 1992;83:344-50.
 61. Chie WC, Chang KJ. Factors related to tumor size of breast cancer at treatment in Taiwan. *Prev Med* 1994;23:91-7.
 62. Grady KE, Kegeles SS, Lund AK, Wolk CH, Farber NJ. Who volunteers for a breast self-examination program? Evaluating the bases for self-selection. *Health Educ Q* 1983;10:79-94.
 63. American Academy of Family Physicians. *Age charts for periodic health examination*. Kansas City (MO): The Academy; 1994.
 64. American College of Obstetricians and Gynecologists. *The obstetrician-gynecologist and primary-preventive health care*. Washington: The College; 1993.
 65. Brett J, Austoker J, Ong G. Do women who undergo further investigation for breast screening suffer adverse psychological consequences? A multi-centre follow-up study comparing different breast screening result groups five months after their last breast screening appointment. *J Public Health Med* 1998;20(4):396-403.
 66. Northouse LL, Jeffs M, Cracchiolo-Caraway A, Lampman L, Dorris G. Emotional distress reported by women and husbands prior to a breast biopsy. *Nurs Res* 1995;44(4):196-201.
 67. Benedict S, Williams RD, Baron PL. Recalled anxiety: from discovery to diagnosis of a benign breast mass. *Oncol Nurs Forum* 1994;21(10):1723-7.
 68. Ellman R, Moss SM, Coleman D, Chamberlain J. Breast self-examination programmes in the trial of early detection of breast cancer: ten year findings. *Br J Cancer* 1993;68:208-12.
 69. Calnan MW, Chamberlain J, Moss S. Compliance with a class teaching breast self examination. *J Epidemiol Community Health* 1983;37:264-70.
 70. MacFarlane ME, Sony SD. Women, breast lump discovery, and associated stress. *Health Care Women Int* 1992;13:23-32.
 71. Lee ECG, Maguire GP. Proceedings: emotional distress in patients attending a breast clinic. *Br J Surg* 1975;62:162.
 72. Frank JW, Mai V. Breast self-examination in young women: More harm than good? *Lancet* 1985;2:654-7.
 73. O'Malley M. Cost-effectiveness of two nurse-led programs to teach breast self-examination. *Am J Prev Med* 1993;9:139-45.
 74. Angiolillo DF, Bennett SE, Lawrence RS, Fenton T, Bennett SD, Schneider GM, et al. Health services utilization following a breast self-examination program [abstract]. *Clin Res* 1987;35:729.
 75. American Cancer Society. Summary of American Cancer Society recommendations for early detection of cancer in asymptomatic people. *CA Cancer J Clin* 1993;43:45.
 76. Canadian Cancer Society. *Seven steps to health*. Toronto: The Society; 1992.
 77. Wald N, Chamberlain J, Hackshaw A. Consensus conference on breast cancer screening. Paris, February 4-5, 1993. Report of the Evaluation Committee. *Oncology* 1994;51:380-9.
 78. United States Preventive Services Task Force. *Guide to clinical preventive services*. 2nd ed. Alexandria (VA): International Medical Publishing; 1996.

Reprint requests to: Canadian Task Force on Preventive Health Care, Parkwood Hospital, 801 Commissioners Rd. E, London ON N6C 5J1; ctf@ctfphc.org

Members of the Canadian Task Force on Preventive Health Care

Chairman: Dr. John W. Feightner, Professor, Department of Family Medicine, University of Western Ontario, London, Ont.; **Members:** Drs. R. Wayne Elford, Professor and Chair of Research, Department of Family Medicine, University of Calgary, Calgary, Alta.; Denice Feig, Assistant Professor, Department of Endocrinology, University of Toronto, Toronto, Ont.; Michel Labrecque, Professeur, Unité de médecine familiale, Université Laval, Rimouski, Qué.; Robin McLeod, Professor, Department of Surgery, Mount Sinai Hospital and University of Toronto, Toronto, Ont.; Harriet MacMillan, Departments of Psychiatry and Behavioural Neurosciences and of Pediatrics, Canadian Centre for Studies of Children at Risk, McMaster University, Hamilton, Ont.; Jean-Marie Moutquin, Professeur titulaire et directeur, Département d'obstétrique-gynécologie, Université de Sherbrooke, Sherbrooke, Que.; Valerie Palda, Assistant Professor, Department of General Internal Medicine, University of Toronto, Toronto, Ont.; Christopher Patterson, Professor and Head, Division of Geriatric Medicine, Department of Medicine, McMaster University, Hamilton, Ont.; and Elaine E.L. Wang, Associate Professor, Departments of Pediatrics and of Public Health Sciences, Faculty of Medicine, University of Toronto, Toronto, Ont. **Resource people:** Nadine Wathen, Coordinator, and Ruth Walton, Research Associate, Canadian Task Force on Preventive Health Care, Department of Family Medicine, University of Western Ontario, London, Ont.

Appendix 1: Canadian Task Force on Preventive Health Care levels of evidence and grades of recommendations**Levels of evidence**

I	Evidence from at least one well-designed randomized controlled trial
II-1	Evidence from well-designed controlled trials without randomization
II-2	Evidence from well-designed cohort or case-control analytic studies, preferably from more than one centre or research group
II-3	Evidence from comparisons between times or places with or without the intervention; dramatic results from uncontrolled studies could be included here
III	Opinions of respected authorities, based on clinical experience; descriptive studies or reports of expert committees

Grades of recommendations

A	Good evidence to support the recommendation that the condition or manoeuvre be specifically considered in a periodic health examination (PHE)
B	Fair evidence to support the recommendation that the condition or manoeuvre be specifically considered in a PHE
C	Insufficient evidence regarding inclusion of the condition or manoeuvre in, or its exclusion from, a PHE, but recommendations may be made on other grounds
D	Fair evidence to support the recommendation that the condition or manoeuvre be specifically excluded from a PHE
E	Good evidence to support the recommendation that the condition or manoeuvre be specifically excluded from a PHE

HOLIDAY REVIEW 2001

CALL FOR PAPERS

Does the only writing you get to do these days involve patients' charts or grant applications? Here's a chance to give your writing muscles a different kind of workout.

We're looking for spoofs of medical research, reflective essays on life and tales of medical adventure (or misadventure) for our 2001 Holiday Review. For inspiration, click on Back Issues at www.cma.ca/cmaj and go to the December issues for 1998, 1999 and 2000. Last year, for example, we published a report on the psychiatric problems facing Winnie T. Pooh and colleagues.

This year, we plan to sprinkle a variety of tidbits throughout the issue, and we need your help. Send us:

- a letter to the editor that could find a home nowhere but the Holiday Review
- a postcard from the place where you live, with an anecdote about your practice
- an original cartoon inspired by your medical career
- a photograph of a day in the life of your office, hospital or clinic (you'll need to get signed consent from any people in the photo)
- an obscure quotation on a holiday theme
- the title of the book you would bring with you if you were admitted to hospital, and the reason why you made this selection
- instructions on how someone in your medical specialty should approach the task of preparing, cooking and carving the holiday bird. The prize: the glory of publishing a winning entry in the first (and probably only) *CMAJ* Talk Turkey contest.



To discuss an idea for the Holiday Review issue, contact the Editor, Dr. John Hoey (tel 800 663-7336 x2118; hoeyj@cma.ca) or the News Editor, Pat Sullivan (800 663-7336 x2126; sullip@cma.ca). Articles should be no more than 1200 words, and illustrations are encouraged. Submissions received by Oct. 1, 2001, are more likely to be published.