

COHOMOLOGY THEORY IN 2-CATEGORIES

HIROYUKI NAKAOKA

ABSTRACT. Recently, symmetric categorical groups are used for the study of the Brauer groups of symmetric monoidal categories. As a part of these efforts, some algebraic structures of the 2-category of symmetric categorical groups SCG are being investigated. In this paper, we consider a 2-categorical analogue of an abelian category, in such a way that it contains SCG as an example. As a main theorem, we construct a long cohomology 2-exact sequence from any extension of complexes in such a 2-category. Our axiomatic and self-dual definition will enable us to simplify the proofs, by analogy with abelian categories.

1. Introduction

In 1970s, B. Pareigis started his study on the Brauer groups of symmetric monoidal categories in [6]. Around 2000, the notion of symmetric categorical groups are introduced to this study by E. M. Vitale in [9] (see also [8]). By definition, a symmetric categorical group is a categorification of an abelian group, and in this sense the 2-category of symmetric categorical groups SCG can be regarded as a 2-dimensional analogue of the category Ab of abelian groups. As such, SCG and its variants (e.g. 2-category \mathbb{G} -SMod of symmetric categorical groups with \mathbb{G} -action where \mathbb{G} is a fixed categorical group) admit a 2-dimensional analogue of the homological algebra in Ab.

For example, E. M. Vitale constructed for any monoidal functor $F : \mathbf{C} \rightarrow \mathbf{D}$ between symmetric monoidal categories \mathbf{C} and \mathbf{D} , a 2-exact sequence of Picard and Brauer categorical groups

$$\mathcal{P}(\mathbf{C}) \rightarrow \mathcal{P}(\mathbf{D}) \rightarrow \overline{\mathcal{F}} \rightarrow \mathcal{B}(\mathbf{C}) \rightarrow \mathcal{B}(\mathbf{D}).$$

By taking π_0 and π_1 , we can induce the well-known Picard-Brauer and Unit-Picard exact sequences of abelian groups respectively. In [7], A. del Río, J. Martínez-Moreno and E. M. Vitale defined a more subtle notion of the relative 2-exactness, and succeeded in constructing a cohomology long 2-exact sequence from any short relatively 2-exact sequence of complexes in SCG. In this paper, we consider a 2-categorical analogue of an abelian category, in such a way that it contains SCG as an example, so as to treat SCG and their variants in a more abstract, unified way.

The author is supported by JSPS. The author wishes to thank Professor Toshiyuki Katsura for his encouragement.

Received by the editors 2007-11-30 and, in revised form, 2008-11-16.

Transmitted by Ross Street. Published on 2008-11-25.

2000 Mathematics Subject Classification: 18D05.

Key words and phrases: symmetric categorical group, 2-category, cohomology, exact sequence.

© Hiroyuki Nakaoka, 2008. Permission to copy for private use granted.

In section 2, we review general definitions in a 2-category and properties of SCG, with simple comments. In section 3, we define the notion of a relatively exact 2-category as a generalization of SCG, also as a 2-dimensional analogue of an abelian category. We try to make the homological algebra in SCG ([7]) work well in this general 2-category. It will be worthy to note that our definition of a relatively exact 2-category is self-dual.

	category	2-category
general theory	abelian category	relatively exact 2-category
example	Ab	SCG

In section 4, we show the existence of *proper factorization systems* in any relatively exact 2-category, which will make several diagram lemmas more easy to handle. In any abelian category, any morphism f can be written in the form $f = e \circ m$ (uniquely up to an isomorphism), where e is epimorphic and m is monomorphic. As a 2-dimensional analogue, we show that any 1-cell f in a relatively exact 2-category \mathbf{S} admits the following two ways of factorization:

- (1) $i \circ m \implies f$ where i is fully cofaithful and m is faithful.
- (2) $e \circ j \implies f$ where e is cofaithful and j is fully faithful.

(In the case of SCG, see [3].) In section 5, complexes in \mathbf{S} and the relative 2-exactness are defined, generalizing those in SCG ([7]). Since we start from the self-dual definition, we can make good use of duality in the proofs. In section 6, as a main theorem, we construct a long cohomology 2-exact sequence from any short relatively 2-exact sequence (i.e. an extension) of complexes. Our proof is purely diagrammatic, and is an analogy of that for an abelian category. In section 5 and 6, several 2-dimensional diagram lemmas are shown. Most of them have 1-dimensional analogues in an abelian category, so we only have to be careful about the compatibility of 2-cells.

Since SCG is an example of a relatively exact 2-category, we expect some other 2-categories constructed from SCG will be a relatively exact 2-category. For example, $\mathbb{G}\text{-SMod}$, $\text{SCG} \times \text{SCG}$ and the 2-category of bifunctors from SCG are candidates. We will examine such examples in forthcoming papers.

2. Preliminaries

DEFINITIONS IN A 2-CATEGORY.

2.1. NOTATION. Throughout this paper, \mathbf{S} denotes a 2-category (in the strict sense). We use the following notation.

$\mathbf{S}^0, \mathbf{S}^1, \mathbf{S}^2$: class of 0-cells, 1-cells, and 2-cells in \mathbf{S} , respectively.

$\mathbf{S}^1(A, B)$: 1-cells from A to B , where $A, B \in \mathbf{S}^0$.

$\mathbf{S}^2(f, g)$: 2-cells from f to g , where $f, g \in \mathbf{S}^1(A, B)$ for certain $A, B \in \mathbf{S}^0$.

$\mathbf{S}(A, B)$: Hom-category between A and B (i.e. $\text{Ob}(\mathbf{S}(A, B)) = \mathbf{S}^1(A, B)$, $\mathbf{S}(A, B)(f, g) = \mathbf{S}^2(f, g)$).

In diagrams, \longrightarrow represents a 1-cell, \implies represents a 2-cell, \circ represents a horizontal composition, and \cdot represents a vertical composition. We use capital letters A, B, \dots for 0-cells, small letters f, g, \dots for 1-cells, and Greek symbols α, β, \dots for 2-cells.

For example, one of the conditions in the definition of a 2-category can be written as follows (see for example [4]):

2.2. REMARK. For any diagram in \mathbf{S}

$$\begin{array}{ccccc}
 & & f_1 & & g_1 \\
 & & \curvearrowright & & \curvearrowright \\
 A & & & B & & C, \\
 & & \alpha \Downarrow & & \beta \Downarrow \\
 & & \curvearrowleft & & \curvearrowleft \\
 & & f_2 & & g_2
 \end{array}$$

we have

$$(f_1 \circ \beta) \cdot (\alpha \circ g_2) = (\alpha \circ g_1) \cdot (f_2 \circ \beta). \tag{1}$$

(Note: composition is always written diagrammatically.)

This equality is frequently used in later arguments.

Products, pullbacks, difference kernels and their duals are defined by the universality.

2.3. DEFINITION. For any A_1 and $A_2 \in \mathbf{S}^0$, their product $(A_1 \times A_2, p_1, p_2)$ is defined as follows:

(a) $A_1 \times A_2 \in \mathbf{S}^0$, $p_i \in \mathbf{S}^1(A_1 \times A_2, A_i)$ ($i = 1, 2$).

(b1) (existence of a factorization)

For any $X \in \mathbf{S}^0$ and $q_i \in \mathbf{S}^1(X, A_i)$ ($i = 1, 2$), there exist $q \in \mathbf{S}^1(X, A_1 \times A_2)$ and $\xi_i \in \mathbf{S}^2(q \circ p_i, q_i)$ ($i = 1, 2$).

$$\begin{array}{ccccc}
 & & X & & \\
 & q_1 \swarrow & \downarrow q & \searrow q_2 & \\
 & \xi_1 \swarrow & & \searrow \xi_2 & \\
 A_1 & \xleftarrow{p_1} & A_1 \times A_2 & \xrightarrow{p_2} & A_2
 \end{array}$$

(b2) (uniqueness of the factorization)

For any factorizations (q, ξ_1, ξ_2) and (q', ξ'_1, ξ'_2) which satisfy (b1), there exists a unique 2-cell $\eta \in \mathbf{S}^2(q, q')$ such that $(\eta \circ p_i) \cdot \xi'_i = \xi_i$ ($i = 1, 2$).

$$\begin{array}{ccc}
 q \circ p_i & \xrightarrow{\eta \circ p_i} & q' \circ p_i \\
 \searrow \xi_i & \circlearrowleft & \searrow \xi'_i \\
 & q_i &
 \end{array}$$

The coproduct of A_1 and A_2 is defined dually.


2.4. DEFINITION. For any $A_1, A_2, B \in \mathbf{S}^0$ and $f_i \in \mathbf{S}^1(A_i, B)$ ($i = 1, 2$), the pullback $(A_1 \times_B A_2, f'_1, f'_2, \xi)$ of f_1 and f_2 is defined as follows:

(a) $A_1 \times_B A_2 \in \mathbf{S}^0$, $f'_i \in \mathbf{S}^1(A_1 \times_B A_2, A_i)$, $f_i \in \mathbf{S}^1(A_i, B)$, $\xi \in \mathbf{S}^2(f'_1 \circ f_2, f'_2 \circ f_1)$.

$$\begin{array}{ccccc}
 & & A_2 & & \\
 & f'_1 \swarrow & & \searrow f_2 & \\
 A_1 \times_B A_2 & & & & B \\
 & f'_2 \searrow & \Downarrow \xi & \swarrow f_1 & \\
 & & A_1 & &
 \end{array}$$

(b1) (existence of a factorization)

For any $X \in \mathbf{S}^0$, $g_1 \in \mathbf{S}^1(X, A_2)$, $g_2 \in \mathbf{S}^1(X, A_1)$ and $\eta \in \mathbf{S}^2(g_1 \circ f_2, g_2 \circ f_1)$, there exist $g \in \mathbf{S}^1(X, A_1 \times A_2)$, $\xi_i \in \mathbf{S}^2(g \circ f'_i, g_i)$ ($i = 1, 2$) such that $(\xi_1 \circ f_2) \cdot \eta = (g \circ \xi) \cdot (\xi_2 \circ f_1)$.



(b2) (uniqueness of the factorization)

For any factorizations (g, ξ_1, ξ_2) and (g', ξ'_1, ξ'_2) which satisfy (b1), there exists a unique 2-cell $\zeta \in \mathbf{S}^2(g, g')$ such that $(\zeta \circ f'_i) \cdot \xi'_i = \xi_i$ ($i = 1, 2$).


The pushout of $f_i \in \mathbf{S}^1(A, B_i)$ ($i = 1, 2$) is defined dually.

2.5. DEFINITION. For any $A, B \in \mathbf{S}^0$ and $f, g \in \mathbf{S}^1(A, B)$, the difference kernel

$$(\text{DK}(f, g), d_{(f,g)}, \varphi_{(f,g)})$$

of f and g is defined as follows:

(a) $\text{DK}(f, g) \in \mathbf{S}^0$, $d_{(f,g)} \in \mathbf{S}^1(\text{DK}(f, g), A)$, $\varphi_{(f,g)} \in \mathbf{S}^2(d_{(f,g)} \circ f, d_{(f,g)} \circ g)$.



(b1) (existence of a factorization)

For any $X \in \mathbf{S}^0$, $d \in \mathbf{S}^1(X, A)$, $\varphi \in \mathbf{S}^2(d \circ f, d \circ g)$, there exist $\underline{d} \in \mathbf{S}^1(X, \text{DK}(f, g))$, $\underline{\varphi} \in \mathbf{S}^2(\underline{d} \circ d_{(f,g)}, d)$ such that $(\underline{d} \circ \varphi_{(f,g)}) \cdot (\underline{\varphi} \circ g) = (\underline{\varphi} \circ f) \cdot \varphi$.

(b2) (uniqueness of the factorization)

For any factorizations $(\underline{d}, \underline{\varphi})$ and $(\underline{d}', \underline{\varphi}')$ which satisfy (b1), there exists a unique 2-cell $\eta \in \mathbf{S}^2(\underline{d}, \underline{d}')$ such that $(\eta \circ d_{(f,g)}) \cdot \underline{\varphi}' = \underline{\varphi}$.

The difference cokernel of f and g is defined dually.

The following definition is from [2].

2.6. DEFINITION. Let $f \in \mathbf{S}^1(A, B)$.

(1) f is said to be faithful if $f^\flat := - \circ f : \mathbf{S}^1(C, A) \rightarrow \mathbf{S}^1(C, B)$ is faithful for any $C \in \mathbf{S}^0$.

(2) f is said to be fully faithful if f^\flat is fully faithful for any $C \in \mathbf{S}^0$.

(3) f is said to be cofaithful if $f^\sharp := f \circ - : \mathbf{S}^1(B, C) \rightarrow \mathbf{S}^1(A, C)$ is faithful for any $C \in \mathbf{S}^0$.

(4) f is said to be fully cofaithful if f^\sharp is fully faithful for any $C \in \mathbf{S}^0$.

PROPERTIES OF SCG. By definition, a symmetric categorical group is a symmetric monoidal category $(\mathbb{G}, \otimes, 0)$, in which each arrow is an isomorphism and each object has an inverse up to an equivalence with respect to the tensor \otimes . More precisely;

2.7. DEFINITION. A symmetric categorical group $(\mathbb{G}, \otimes, 0)$ consists of

- (a1) a category \mathbb{G}
- (a2) a tensor functor $\otimes : \mathbb{G} \times \mathbb{G} \rightarrow \mathbb{G}$
- (a3) a unit object $0 \in \text{Ob}(\mathbb{G})$
- (a4) natural isomorphisms

$$\begin{aligned} \alpha_{A,B,C} &: A \otimes (B \otimes C) \rightarrow (A \otimes B) \otimes C, \\ \lambda_A &: 0 \otimes A \rightarrow A, \quad \rho_A : A \otimes 0 \rightarrow A, \quad \gamma_{A,B} : A \otimes B \rightarrow B \otimes A \end{aligned}$$

which satisfy certain compatibility conditions (cf. [5]), and the following two conditions are satisfied:

- (b1) For any $A, B \in \text{Ob}(\mathbb{G})$ and $f \in \mathbb{G}(A, B)$, there exists $g \in \mathbb{G}(B, A)$ such that $f \circ g = \text{id}_A$, $g \circ f = \text{id}_B$.
- (b2) For any $A \in \text{Ob}(\mathbb{G})$, there exist $A^* \in \text{Ob}(\mathbb{G})$ and $\eta_A \in \mathbb{G}(0, A \otimes A^*)$.

In particular, there is a ‘zero categorical group’ 0 , which consists of only one object 0 and one morphism id_0 .

2.8. DEFINITION. For symmetric categorical groups \mathbb{G} and \mathbb{H} , a monoidal functor F from \mathbb{G} to \mathbb{H} consists of

- (a1) a functor $F : \mathbb{G} \rightarrow \mathbb{H}$
- (a2) natural isomorphisms

$$F_{A,B} : F(A \otimes B) \rightarrow F(A) \otimes F(B) \text{ and } F_I : F(0) \rightarrow 0$$

which satisfy certain compatibilities with α , λ , ρ , γ . (cf. [5])

2.9. REMARK. For any monoidal functors $F : \mathbb{G} \rightarrow \mathbb{H}$ and $G : \mathbb{H} \rightarrow \mathbb{K}$, their composition $F \circ G : \mathbb{G} \rightarrow \mathbb{K}$ is defined by

$$(F \circ G)_{A,B} := G(F_{A,B}) \circ G_{F(A), F(B)} \quad (2)$$

$$(F \circ G)_I := G(F_I) \circ G_I. \quad (3)$$

In particular, there is a ‘zero monoidal functor’ $0_{\mathbb{G}, \mathbb{H}} : \mathbb{G} \rightarrow \mathbb{H}$ for each \mathbb{G} and \mathbb{H} , which sends every object in \mathbb{G} to $0_{\mathbb{H}}$, every arrow in \mathbb{G} to $\text{id}_{0_{\mathbb{H}}}$, and $(0_{\mathbb{G}, \mathbb{H}})_{A,B} = \lambda_0^{-1} = \rho_0^{-1}$, $(0_{\mathbb{G}, \mathbb{H}})_I = \text{id}_0$. It is easy to see that $0_{\mathbb{G}, \mathbb{H}} \circ 0_{\mathbb{H}, \mathbb{K}} = 0_{\mathbb{G}, \mathbb{K}}$ ($\forall \mathbb{G}, \mathbb{H}, \mathbb{K}$).

2.10. REMARK. Our notion of a monoidal functor is equal to that of a ‘ γ -monoidal functor’ in [7].

2.11. DEFINITION. For monoidal functors $F, G : \mathbb{G} \rightarrow \mathbb{H}$, a natural transformation φ from F to G is said to be a monoidal transformation if it satisfies

$$\begin{aligned}\varphi_{A \otimes B} \circ G_{A,B} &= F_{A,B} \circ (\varphi_A \otimes \varphi_B) \\ F_I &= \varphi_0 \circ G_I.\end{aligned}\tag{4}$$

The following remark is from [9].

2.12. REMARK. By condition (b2), it is shown that there exists a 2-cell $\varepsilon_A \in \mathbb{G}(A^* \otimes A, 0)$ for each object A , such that the following compositions are identities:

$$\begin{array}{ccccccccccc} A & \xrightarrow{\lambda_A^{-1}} & 0 \otimes A & \xrightarrow{\eta_{A \otimes 1}} & (A \otimes A^*) \otimes A & \xrightarrow{\alpha^{-1}} & A \otimes (A^* \otimes A) & \xrightarrow{1 \otimes \varepsilon_A} & A \otimes 0 & \xrightarrow{\rho_A} & A \\ A^* & \xrightarrow{\rho_{A^*}^{-1}} & A^* \otimes 0 & \xrightarrow{1 \otimes \eta_A} & A^* \otimes (A \otimes A^*) & \xrightarrow{\alpha} & (A^* \otimes A) \otimes A^* & \xrightarrow{\varepsilon_{A \otimes 1}} & 0 \otimes A^* & \xrightarrow{\lambda_{A^*}} & A^* \end{array}$$

For each monoidal functor $F : \mathbb{G} \rightarrow \mathbb{H}$, there exists a natural morphism $\iota_{F,A} : F(A^*) \rightarrow F(A)^*$.

2.13. DEFINITION. SCG is defined to be the 2-category whose 0-cells are symmetric categorical groups, 1-cells are monoidal functors, and 2-cells are monoidal transformations.

The following two propositions are satisfied in SCG (see for example [1]).

2.14. PROPOSITION. For any symmetric categorical groups \mathbb{G} and \mathbb{H} , if we define a monoidal functor $F \otimes_{\mathbb{G}, \mathbb{H}} G : \mathbb{G} \rightarrow \mathbb{H}$ by

$$\begin{aligned} F \otimes_{\mathbb{G}, \mathbb{H}} G(A) &:= F(A) \otimes_{\mathbb{H}} G(A) \\ (F \otimes_{\mathbb{G}, \mathbb{H}} G)_{A,B} &:= (F(A \otimes B) \otimes G(A \otimes B)) \\ &\xrightarrow{F_{A,B} \otimes G_{A,B}} F(A) \otimes F(B) \otimes G(A) \otimes G(B) \\ &\xrightarrow{\simeq} F(A) \otimes G(A) \otimes F(B) \otimes G(B) \\ (F \otimes_{\mathbb{G}, \mathbb{H}} G)_I &:= (F(I) \otimes G(I) \xrightarrow{F_I \otimes G_I} I \otimes I \xrightarrow{\simeq} I), \end{aligned}$$

then $(\text{SCG}(\mathbb{G}, \mathbb{H}), \otimes_{\mathbb{G}, \mathbb{H}}, 0_{\mathbb{G}, \mathbb{H}})$ becomes again a symmetric categorical group with appropriately defined $\alpha, \lambda, \rho, \gamma$, and

$$\text{Hom} = \text{SCG}(-, -) : \text{SCG} \times \text{SCG} \rightarrow \text{SCG}$$

becomes a 2-functor (cf. section 6 in [1]).

In SCG, by definition of the zero categorical group we have $\mathbf{S}^1(\mathbb{G}, 0) = \{0_{\mathbb{G}, 0}\}$, while $\mathbf{S}^1(0, \mathbb{G})$ may have more than one objects. In this point SCG might be said to have ‘non self-dual’ structure, but $\mathbf{S}^1(\mathbb{G}, 0)$ and $\mathbf{S}^1(0, \mathbb{G})$ have the following ‘self-dual’ property.

2.15. REMARK. (1) For any symmetric categorical group \mathbb{G} and any monoidal functor $F : \mathbb{G} \rightarrow 0$, there exists a unique 2-cell $\varphi : F \Longrightarrow 0_{\mathbb{G}, 0}$.

(2) For any symmetric categorical group \mathbb{G} and any monoidal functor $F : 0 \rightarrow \mathbb{G}$, there exists a unique 2-cell $\varphi : F \Longrightarrow 0_{0, \mathbb{G}}$.

PROOF. (1) follows from the fact that the zero categorical group has only one morphism id_0 . (2) follows from condition (4) in Definition 2.11. \blacksquare

The usual compatibility arguments show the following Lemma.

2.16. LEMMA. *Let $F : \mathbb{G} \rightarrow \mathbb{H}$ be a monoidal functor. For any $A, B \in \text{Ob}(\mathbb{G})$,*

$$\begin{array}{ccc} \Phi_{A,B} : \mathbb{G}(A, B) & \longrightarrow & \mathbb{G}(A \otimes B^*, 0) \\ \Downarrow & & \Downarrow \\ f & \longmapsto & (f \otimes 1_{B^*}) \circ \eta_B^{-1} \end{array}$$

and

$$\begin{array}{ccc} \Psi_{A,B} : \mathbb{G}(A \otimes B^*, 0) & \longrightarrow & \mathbb{G}(A, B) \\ \Downarrow & & \Downarrow \\ g & \longmapsto & \rho_A^{-1} \circ (1_A \otimes \varepsilon_B^{-1}) \circ \alpha_{A, B^*, B} \circ (g \otimes 1_B) \circ \lambda_B \end{array}$$

are mutually inverse, and the following diagram is commutative;

$$\begin{array}{ccc} \mathbb{G}(A, B) & \xrightarrow{\Phi_{A,B}} & \mathbb{G}(A \otimes B^*, 0) \\ \swarrow F & & \searrow F \\ \mathbb{H}(F(A), F(B)) & \circ & \mathbb{H}(F(A \otimes B^*), F(0)) \\ \searrow \Phi_{F(A), F(B)} & & \swarrow \Theta_{A,B}^F \\ & \mathbb{H}(F(A) \otimes F(B)^*, 0) & \end{array}$$

where $\Theta_{A,B}^F$ is defined by

$$\begin{array}{ccc} \Theta_{A,B}^F : \mathbb{H}(F(A \otimes B^*), F(0)) & \longrightarrow & \mathbb{H}(F(A) \otimes F(B)^*, 0) \\ \Downarrow & & \Downarrow \\ h & \longmapsto & (1_{F(A)} \otimes (\iota_B^F)^{-1}) \circ (F_{A, B^*})^{-1} \circ h \circ F_I. \end{array}$$

3. Definition of a relatively exact 2-category

LOCALLY SCG 2-CATEGORY. We define a locally SCG 2-category not only as a 2-category whose Hom-categories are SCG, but with some more conditions, in order to let it be a 2-dimensional analogue of that of an additive category.

3.1. DEFINITION. *A locally small 2-category \mathbf{S} is said to be locally SCG if the following conditions are satisfied:*

(A1) *For every $A, B \in \mathbf{S}^0$, there is a given functor $\otimes_{A,B} : \mathbf{S}(A, B) \times \mathbf{S}(A, B) \rightarrow \mathbf{S}(A, B)$, and a given object $0_{A,B} \in \text{Ob}(\mathbf{S}(A, B)) = \mathbf{S}^1(A, B)$ such that $(\mathbf{S}(A, B), \otimes_{A,B}, 0_{A,B})$ becomes a symmetric categorical group, and the following naturality conditions are satisfied:*

$$0_{A,B} \circ 0_{B,C} = 0_{A,C} \quad (\forall A, B, C \in \mathbf{S}^0)$$

(A2) *Hom = $\mathbf{S}(-, -) : \mathbf{S} \times \mathbf{S} \rightarrow \text{SCG}$ is a 2-functor which satisfies for any $A, B, C \in \mathbf{S}^0$,*

$$(0_{A,B})_I^\sharp = \text{id}_{0_{A,C}} \in \mathbf{S}^2(0_{A,C}, 0_{A,C}) \quad (5)$$

$$(0_{A,B})_I^\flat = \text{id}_{0_{C,B}} \in \mathbf{S}^2(0_{C,B}, 0_{C,B}). \quad (6)$$

(A3) There is a 0-cell $0 \in \mathbf{S}^0$ called a zero object, which satisfy the following conditions:

(a3-1) $\mathbf{S}(0, 0)$ is the zero categorical group.

(a3-2) For any $A \in \mathbf{S}^0$ and $f \in \mathbf{S}^1(0, A)$, there exists a unique 2-cell $\theta_f \in \mathbf{S}^2(f, 0_{0,A})$.

(a3-3) For any $A \in \mathbf{S}^0$ and $f \in \mathbf{S}^1(A, 0)$, there exists a unique 2-cell $\tau_f \in \mathbf{S}^2(f, 0_{A,0})$.

(A4) For any $A, B \in \mathbf{S}^0$, their product and coproduct exist.

Let us explain about these conditions.

3.2. REMARK. By condition (A1) of Definition 3.1, every 2-cell in a locally SCG 2-category becomes invertible, as in the case of SCG (cf. [9]). This helps us to avoid being fussy about the directions of 2-cells in many propositions and lemmas, and we use the word ‘dual’ simply to reverse 1-cells.

3.3. REMARK. By condition (A2) in Definition 3.1,

$$\begin{aligned} f^\sharp &:= f \circ - : \mathbf{S}(B, C) \rightarrow \mathbf{S}(A, C) \\ f^\flat &:= - \circ f : \mathbf{S}(C, A) \rightarrow \mathbf{S}(C, B) \end{aligned}$$

are monoidal functors ($\forall C \in \mathbf{S}^0$) for any $f \in \mathbf{S}^1(A, B)$, and the following naturality conditions are satisfied:

(a2-1) For any $f \in \mathbf{S}^1(A, B)$, $g \in \mathbf{S}^1(B, C)$ and $D \in \mathbf{S}^0$, we have $(f \circ g)^\sharp = g^\sharp \circ f^\sharp$ as monoidal functors.

$$\begin{array}{ccc} A \xrightarrow{f} B \xrightarrow{g} C \longrightarrow D & & \begin{array}{ccc} \mathbf{S}(C, D) & \xrightarrow{g^\sharp} & \mathbf{S}(B, D) \\ & \searrow (f \circ g)^\sharp & \circ & \swarrow f^\sharp \\ & & \mathbf{S}(A, D) \end{array} \end{array}$$

(a2-2) The dual of (a2-1) for $-^\flat$.

(a2-3) For any $f \in \mathbf{S}^1(A, B)$, $g \in \mathbf{S}^1(C, D)$, we have $f^\sharp \circ g^\flat = g^\flat \circ f^\sharp$ as monoidal functors.

$$\begin{array}{ccc} A \xrightarrow{f} B \longrightarrow C \xrightarrow{g} D & & \begin{array}{ccc} \mathbf{S}(B, C) & \xrightarrow{f^\sharp} & \mathbf{S}(A, C) \\ g^\flat \downarrow & \circ & \downarrow g^\flat \\ \mathbf{S}(B, D) & \xrightarrow{f^\sharp} & \mathbf{S}(A, D) \end{array} \end{array}$$

Since already $(f \circ g)^\sharp = g^\sharp \circ f^\sharp$ as functors, (a2-1) means $(f \circ g)_I^\sharp = (g^\sharp \circ f^\sharp)_I$, and by (3) in Remark 2.9, this is equivalent to

$$(f \circ g)_I^\sharp = f^\sharp(g_I^\sharp) \cdot f_I^\sharp = (f \circ g_I^\sharp) \cdot f_I^\sharp.$$

Similarly, we obtain

$$(f \circ g)_I^\flat = (f_I^\flat \circ g) \cdot g_I^\flat, \quad (7)$$

$$(f_I^\sharp \circ g) \cdot g_I^\flat = (f \circ g_I^\flat) \cdot f_I^\sharp. \quad (8)$$

3.4. **REMARK.** By condition (A2), for any $f, g \in \mathbf{S}^1(A, B)$ and any $\alpha \in \mathbf{S}^2(f, g)$, $\alpha \circ - : f^\sharp \Rightarrow g^\sharp$ becomes a monoidal transformation. So, the diagrams

$$\begin{array}{ccc} f \circ (k \otimes h) & \xrightarrow{\alpha \circ (k \otimes h)} & g \circ (k \otimes h) \\ f_{k,h}^\sharp \Downarrow & \circlearrowleft & \Downarrow g_{k,h}^\sharp \\ (f \circ k) \otimes (f \circ h) & \xrightarrow{(\alpha \circ k) \otimes (\alpha \circ h)} & (g \circ k) \otimes (g \circ h) \end{array} \quad \text{and} \quad \begin{array}{ccc} f \circ 0_{B,C} & \xrightarrow{\alpha \circ 0_{B,C}} & g \circ 0_{B,C} \\ f_I^\sharp \searrow & \circlearrowleft & \swarrow g_I^\sharp \\ & 0_{A,C} & \end{array}$$

are commutative for any $C \in \mathbf{S}^0$ and $k, h \in \mathbf{S}^1(B, C)$. Similar statement also holds for $- \circ \alpha : f^\flat \Rightarrow g^\flat$.

3.5. **COROLLARY.** *In a locally SCG 2-category \mathbf{S} , the following are satisfied:*

(1) *For any diagram in \mathbf{S}*

$$\begin{array}{ccc} C & \xrightarrow{h} & A & \xrightarrow{f} & B \\ & \searrow \varepsilon & \Downarrow \alpha & \searrow & \\ & 0_{C,A} & & g & \end{array}$$

we have

$$h \circ \alpha = (\varepsilon \circ f) \cdot f_I^\flat \cdot g_I^{\flat-1} \cdot (\varepsilon^{-1} \circ g). \quad (9)$$

(2) *For any diagram in \mathbf{S}*

$$\begin{array}{ccc} A & \xrightarrow{f} & B & \xrightarrow{h} & C \\ & \searrow \alpha & \Downarrow \varepsilon & \searrow & \\ & g & & 0_{B,C} & \end{array}$$

we have

$$\alpha \circ h = (f \circ \varepsilon) \cdot f_I^\sharp \cdot g_I^{\sharp-1} \cdot (g \circ \varepsilon^{-1}). \quad (10)$$

(3) *For any diagram in \mathbf{S}*

$$\begin{array}{ccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ & \searrow \alpha & \Downarrow \beta & \searrow & \\ & 0_{A,B} & & 0_{B,C} & \end{array}$$

we have

$$(f \circ \beta) \cdot f_I^\sharp = (\alpha \circ g) \cdot g_I^\flat. \quad (11)$$

PROOF. (1) $(h \circ \alpha) = (\varepsilon \circ f) \cdot (0_{C,A} \circ \alpha) \cdot (\varepsilon^{-1} \circ g) = (\varepsilon \circ f) \cdot f_I^\flat \cdot g_I^{\flat-1} \cdot (\varepsilon^{-1} \circ g)$. (2) is the dual of (1). And (3) follows from (5), (6), (9), (10). \blacksquare

3.6. **REMARK.** We don't require a locally SCG 2-category to satisfy $\mathbf{S}^1(A, 0) = \{0_{A,0}\}$, for the sake of duality (see the comments before Remark 2.15).

RELATIVELY EXACT 2-CATEGORY.

3.7. DEFINITION. Let \mathbf{S} be a locally SCG 2-category. \mathbf{S} is said to be relatively exact if the following conditions are satisfied:

- (B1) For any 1-cell $f \in \mathbf{S}^1(A, B)$, its kernel and cokernel exist.
- (B2) For any 1-cell $f \in \mathbf{S}^1(A, B)$, f is faithful if and only if $f = \ker(\text{cok}(f))$.
- (B3) For any 1-cell $f \in \mathbf{S}^1(A, B)$, f is cofaithful if and only if $f = \text{cok}(\ker(f))$.


It is shown in [9] that SCG satisfies these conditions.

Let us explain about these conditions.

3.8. DEFINITION. Let \mathbf{S} be a locally SCG 2-category. For any $f \in \mathbf{S}^1(A, B)$, its kernel $(\text{Ker}(f), \ker(f), \varepsilon_f)$ is defined by universality as follows (we abbreviate $\ker(f)$ to $k(f)$) :

- (a) $\text{Ker}(f) \in \mathbf{S}^0$, $k(f) \in \mathbf{S}^1(\text{Ker}(f), A)$, $\varepsilon_f \in \mathbf{S}^2(k(f) \circ f, 0)$.
- (b1) (existence of a factorization)

For any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, A)$ and $\varepsilon \in \mathbf{S}^2(k \circ f, 0)$, there exist $\underline{k} \in \mathbf{S}^1(K, \text{Ker}(f))$ and $\underline{\varepsilon} \in \mathbf{S}^2(\underline{k} \circ k(f), k)$ such that $(\underline{\varepsilon} \circ f) \cdot \varepsilon = (\underline{k} \circ \varepsilon_f) \cdot (\underline{k})_f^\sharp$.



- (b2) (uniqueness of the factorization)

For any factorizations $(\underline{k}, \underline{\varepsilon})$ and $(\underline{k}', \underline{\varepsilon}')$ which satisfy (b1), there exists a unique 2-cell $\xi \in \mathbf{S}^2(\underline{k}, \underline{k}')$ such that $(\xi \circ k(f)) \cdot \underline{\varepsilon}' = \underline{\varepsilon}$.

3.9. REMARK. (1) By its universality, the kernel of f is unique up to an equivalence. We write this equivalence class again $\text{Ker}(f) = [\text{Ker}(f), k(f), \varepsilon_f]$.


(2) It is also easy to see that if f and f' are equivalent, then

$$[\text{Ker}(f), k(f), \varepsilon_f] = [\text{Ker}(f'), k(f'), \varepsilon_{f'}].$$

For any f , its cokernel $\text{Cok}(f) = [\text{Cok}(f), c(f), \pi_f]$ is defined dually, and the dual statements also hold for the cokernel.

3.10. REMARK. Let \mathbf{S} be a locally SCG 2-category, and let f be in $\mathbf{S}^1(A, B)$.

For any pair (k, ε) with $k \in \mathbf{S}^1(0, A)$, $\varepsilon \in \mathbf{S}^2(k \circ f, 0)$



and for any pair (k', ε') with $k' \in \mathbf{S}^1(0, A)$, $\varepsilon' \in \mathbf{S}^2(k' \circ f, 0)$, there exists a unique 2-cell $\xi \in \mathbf{S}^2(k, k')$ such that $(\xi \circ f) \cdot \varepsilon' = \varepsilon$.

PROOF. By condition (a3-2) of Definition 3.1, $\varepsilon \in \mathbf{S}^2(k \circ f, 0)$ must be equal to the unique 2-cell $(\theta_k \circ f) \cdot f_I^\flat$. Similarly we have $\varepsilon' = (\theta_{k'} \circ f) \cdot f_I^\flat$, and, ξ should be the unique 2-cell $\theta_k \cdot \theta_{k'}^{-1} \in \mathbf{S}^2(k, k')$, which satisfies $(\xi \circ f) \cdot \varepsilon' = \varepsilon$. ■

From this, it makes no ambiguity if we abbreviate $\text{Ker}(f) = [0, 0_{0,A}, f_I^\natural]$ to $\text{Ker}(f) = 0$, because $[0, k, \varepsilon] = [0, k', \varepsilon']$ for any (k, ε) and (k', ε') . Dually, we abbreviate $\text{Cok}(f) = [0, 0_{A,0}, f_I^\sharp]$ to $\text{Cok}(f) = 0$.

By using condition (A3) of Definition 3.1, we can show the following easily:

- 3.11. EXAMPLE. (1) For any $A \in \mathbf{S}^0$, $\text{Ker}(0_{A,0} : A \rightarrow 0) = [A, \text{id}_A, \text{id}_0]$.
- (2) For any $A \in \mathbf{S}^0$, $\text{Cok}(0_{0,A} : 0 \rightarrow A) = [A, \text{id}_A, \text{id}_0]$.

3.12. CAUTION. (1) $\text{Ker}(0_{0,A} : 0 \rightarrow A)$ need not be equivalent to 0. Indeed, in the case of SCG, for any symmetric categorical group \mathbb{G} , $\text{Ker}(0_{0,\mathbb{G}} : 0 \rightarrow \mathbb{G})$ is equivalent to an important invariant $\pi_1(\mathbb{G})[0]$.

(2) $\text{Cok}(0_{A,0} : A \rightarrow 0)$ need not be equivalent to 0 either. In the case of SCG, $\text{Cok}(0_{\mathbb{G},0} : \mathbb{G} \rightarrow 0)$ is equivalent to $\pi_0(\mathbb{G})[1]$.

3.13. REMARK. The precise meaning of condition (B2) in Definition 3.7 is that, for any 1-cell $f \in \mathbf{S}^1(A, B)$ and its cokernel $[\text{Cok}(f), \text{cok}(f), \pi_f]$, f is faithful if and only if $\text{Ker}(\text{cok}(f)) = [A, f, \pi_f]$. Similarly for condition (B3).

RELATIVE (CO-)KERNEL AND FIRST PROPERTIES OF A RELATIVELY EXACT 2-CATEGORY. Throughout this subsection, \mathbf{S} is a relatively exact 2-category.

3.14. DEFINITION. For any diagram in \mathbf{S}

$$\begin{array}{ccc}
 & 0 & \\
 & \uparrow \varphi & \\
 A & \xrightarrow{f} B & \xrightarrow{g} C,
 \end{array}
 \tag{12}$$


its relative kernel $(\text{Ker}(f, \varphi), \text{ker}(f, \varphi), \varepsilon_{(f,\varphi)})$ is defined as follows (we abbreviate $\text{ker}(f, \varphi)$ to $k(f, \varphi)$):

- (a) $\text{Ker}(f, \varphi) \in \mathbf{S}^0$, $k(f, \varphi) \in \mathbf{S}^1(\text{Ker}(f, \varphi), A)$, $\varepsilon_{(f,\varphi)} \in \mathbf{S}^2(k(f, \varphi) \circ f, 0)$.
- (b0) (compatibility of the 2-cells)

$$\varepsilon_{(f,\varphi)} \text{ is compatible with } \varphi \text{ i.e. } (k(f, \varphi) \circ \varphi) \cdot k(f, \varphi)_I^\sharp = (\varepsilon_{(f,\varphi)} \circ g) \cdot g_I^\flat.$$

- (b1) (existence of a factorization)

For any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, A)$ and $\varepsilon \in \mathbf{S}^2(k \circ f, 0)$ which are compatible with φ , i.e. $(k \circ \varphi) \cdot k_I^\sharp = (\varepsilon \circ g) \cdot g_I^\flat$, there exist $\underline{k} \in \mathbf{S}^1(K, \text{Ker}(f, \varphi))$ and $\underline{\varepsilon} \in \mathbf{S}^2(\underline{k} \circ k(f, \varphi), k)$ such that $(\underline{\varepsilon} \circ f) \cdot \varepsilon = (\underline{k} \circ \varepsilon_{(f,\varphi)}) \cdot (\underline{k})_I^\sharp$.



- (b2) (uniqueness of the factorization)

For any factorizations $(\underline{k}, \underline{\varepsilon})$ and $(\underline{k}', \underline{\varepsilon}')$ which satisfy (b1), there exists a unique 2-cell $\xi \in \mathbf{S}^2(\underline{k}, \underline{k}')$ such that $(\xi \circ k(f, \varphi)) \cdot \underline{\varepsilon}' = \underline{\varepsilon}$.

3.15. **REMARK.** By its universality, the relative kernel of (f, φ) is unique up to an equivalence. We write this equivalence class $[\text{Ker}(f, \varphi), k(f, \varphi), \varepsilon_{(f, \varphi)}]$.

3.16. **DEFINITION.** Let \mathbf{S} be a relatively exact 2-category. For any diagram (12) in \mathbf{S} , its relative cokernel $(\text{Cok}(g, \varphi), \text{cok}(g, \varphi), \pi_{(g, \varphi)})$ is defined dually by universality. We abbreviate $\text{cok}(g, \varphi)$ to $c(g, \varphi)$, and write the equivalence class of the relative cokernel $[\text{Cok}(g, \varphi), c(g, \varphi), \pi_{(g, \varphi)}]$.

3.17. **CAUTION.** In the rest of this paper, \mathbf{S} denotes a relatively exact 2-category, unless otherwise specified. In the following propositions and lemmas, we often omit the statement and the proof of their duals. Each term should be replaced by its dual. For example, kernel and cokernel, faithfulness and cofaithfulness are mutually dual.

3.18. **REMARK.** By using condition (A3) of Definition 3.1, we can show the following easily. (These are also corollaries of Proposition 3.20.)

(1) $\text{Ker}(f, f_I^\sharp) = \text{Ker}(f)$ (and thus the ordinary kernel can be regarded as a relative kernel).

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \curvearrowright & \uparrow \uparrow f_I^\sharp & \curvearrowleft & \\
 A & \xrightarrow{f} & B & \xrightarrow{0} & 0
 \end{array}$$

(2) $\ker(f, \varphi)$ is faithful.

3.19. **LEMMA.** Let $f \in \mathbf{S}^1(A, B)$ and take its kernel $[\text{Ker}(f), k(f), \varepsilon_f]$. If $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, \text{Ker}(f))$ and $\sigma \in \mathbf{S}^2(k \circ k(f), 0)$

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \uparrow \uparrow \varepsilon_f & & \\
 K & \xrightarrow{k} & \text{Ker}(f) & \xrightarrow{k(f)} & A & \xrightarrow{f} & B \\
 & \searrow \sigma & \downarrow & \nearrow & & & \\
 & & 0 & & & &
 \end{array}$$

is compatible with ε_f , i.e. if σ satisfies

$$(\sigma \circ f) \cdot f_I^\flat = (k \circ \varepsilon_f) \cdot k_I^\sharp, \tag{13}$$

then there exists a unique 2-cell $\zeta \in \mathbf{S}^2(k, 0)$ such that $\sigma = (\zeta \circ k(f)) \cdot k(f)_I^\flat$.

PROOF. By (13), $\sigma : k \circ k(f) \implies 0$ is a factorization compatible with ε_f and f_I^\flat . On the other hand, by Remark 3.4, $k(f)_I^\flat : 0 \circ k(f) \implies 0$ is also a factorization compatible with ε_f, f_I^\flat . So, by the universality of the kernel, there exists a unique 2-cell $\zeta \in \mathbf{S}^2(k, 0)$ such that $\sigma = (\zeta \circ k(f)) \cdot k(f)_I^\flat$. ■

It is easy to see that the same statement also holds for relative (co-)kernels. In any relatively exact 2-category, the relative (co-)kernel always exist. More precisely, the following proposition holds.

3.20. PROPOSITION. Consider diagram (12) in \mathbf{S} . By the universality of $\text{Ker}(g) = [\text{Ker}(g), \ell, \varepsilon]$, f factors through ℓ uniquely up to an equivalence as $\underline{\varphi} : \underline{f} \circ \ell \Rightarrow f$, where $\underline{f} \in \mathbf{S}^1(A, \text{Ker}(g))$ and $\underline{\varphi} \in \mathbf{S}^2(\underline{f} \circ \ell, f)$:

$$(\underline{f} \circ \varepsilon) \cdot (\underline{f})_I^\sharp = (\underline{\varphi} \circ g) \cdot \varphi$$

Then we have $\text{Ker}(f, \varphi) = [\text{Ker}(f), k(f), \eta]$, where $\eta := (k(f) \circ \varphi^{-1}) \cdot (\varepsilon_f \circ \ell) \cdot \ell_I^\flat \in \mathbf{S}^2(k(f) \circ f, 0)$. We abbreviate this to $\text{Ker}(f, \varphi) = \text{Ker}(f)$.

PROOF. For any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, A)$ and $\sigma \in \mathbf{S}^2(k \circ f, 0)$ which are compatible with φ , i.e. $(\sigma \circ g) \cdot g_I^\flat = (k \circ \varphi) \cdot k_I^\sharp$, if we put

$$\rho := (k \circ \underline{\varphi}) \cdot \sigma \in \mathbf{S}^2(k \circ \underline{f} \circ \ell, 0),$$

then ρ is compatible with ε . By Lemma 3.19, there exists a 2-cell $\zeta : k \circ \underline{f} \Rightarrow 0$ such that $\rho = (\zeta \circ \ell) \cdot \ell_I^\flat$. So, by the universality of $\text{Ker}(f)$, there exist $\underline{k} \in \mathbf{S}^1(K, \text{Ker}(f))$ and $\underline{\sigma} \in \mathbf{S}^2(\underline{k} \circ k(f), k)$ such that $(\underline{\sigma} \circ \underline{f}) \cdot \zeta = (\underline{k} \circ \varepsilon_f) \cdot (\underline{k})_I^\sharp$. Then, $\underline{\sigma}$ is compatible with σ and η ,

and the existence of a factorization is shown. To show the uniqueness of the factorization, let $(\underline{k}', \underline{\sigma}')$ be another factorization which is compatible with σ , η , i.e. $(\underline{\sigma}' \circ \underline{f}) \cdot \sigma = (\underline{k}' \circ \eta) \cdot (\underline{k}')_I^\sharp$. Then, by using $\eta = (k(f) \circ \varepsilon_f^{-1}) \cdot (\varepsilon_f \circ \ell) \cdot \ell_I^\flat$ and $\zeta \circ \ell = \rho \cdot \ell_I^{\flat-1} = (k \circ \varepsilon) \cdot \sigma \cdot \ell_I^{\flat-1}$, we can show $((\underline{\sigma}' \circ \underline{f}) \cdot \zeta) \circ \ell = ((\underline{k}' \circ \varepsilon_f) \cdot (\underline{k}')_I^\sharp) \circ \ell$. Since ℓ is faithful, we obtain $((\underline{\sigma}' \circ \underline{f}) \cdot \zeta) = (\underline{k}' \circ \varepsilon_f) \cdot (\underline{k}')_I^\sharp$. Thus, $\underline{\sigma}'$ is compatible with ζ and ε_f . By the universality of $\text{Ker}(f)$, there exists a 2-cell $\xi \in \mathbf{S}^2(\underline{k}, \underline{k}')$ such that $(\xi \circ k(f)) \cdot \underline{\sigma}' = \underline{\sigma}$. Uniqueness of such $\xi \in \mathbf{S}^2(\underline{k}, \underline{k}')$ follows from the faithfulness of $k(f)$. ■

3.21. PROPOSITION. Let $f \in \mathbf{S}^1(A, B)$, $g \in \mathbf{S}^1(B, C)$ and suppose g is fully faithful. Then, $\text{Ker}(f \circ g) = [\text{Ker}(f), k(f), (\varepsilon_f \circ g) \cdot g_I^\flat]$. We abbreviate this to $\text{Ker}(f \circ g) = \text{Ker}(f)$.

PROOF. Since g is fully faithful, for any $K \in \mathbf{S}^0, k \in \mathbf{S}^1(K, A)$ and $\sigma \in \mathbf{S}^2(k \circ f \circ g, 0)$, there exists $\rho \in \mathbf{S}^2(k \circ f, 0)$ such that $\sigma = (\rho \circ g) \cdot g_I^b$. And by the universality of $\text{Ker}(f)$, there are $\underline{k} \in \mathbf{S}^1(K, \text{Ker}(f))$ and $\underline{\sigma} \in \mathbf{S}^2(\underline{k} \circ k(f), k)$ such that $(\underline{\sigma} \circ f) \cdot \rho = (\underline{k} \circ \varepsilon_f) \cdot (\underline{k})_I^\sharp$. Then, it can be easily seen that $\underline{\sigma}$ is compatible with σ and $(\varepsilon_f \circ g) \cdot g_I^b$:

$$(\underline{\sigma} \circ f \circ g) \cdot \sigma = (\underline{k} \circ ((\varepsilon_f \circ g) \cdot g_I^b)) \cdot (\underline{k})_I^\sharp$$

Thus we obtain a desired factorization. To show the uniqueness of the factorization, let $(\underline{k}', \underline{\sigma}')$ be another factorization of k which satisfies

$$(\underline{\sigma}' \circ f \circ g) \cdot \sigma = (\underline{k}' \circ ((\varepsilon_f \circ g) \cdot g_I^b)) \cdot (\underline{k}')_I^\sharp$$

Then, we can show $\underline{\sigma}'$ is compatible with ρ and ε_f . By the universality of $\text{Ker}(f)$, there exists a 2-cell $\xi \in \mathbf{S}^2(\underline{k}, \underline{k}')$ such that $(\xi \circ k(f)) \cdot \underline{\sigma}' = \underline{\sigma}$. Uniqueness of such ξ follows from the faithfulness of $k(f)$. ■

By definition, $f \in \mathbf{S}^1(A, B)$ is faithful (resp. fully faithful) if and only if $- \circ f : \mathbf{S}^2(g, h) \rightarrow \mathbf{S}^2(g \circ f, h \circ f)$ is injective (resp. bijective) for any $K \in \mathbf{S}^0$ and any $g, h \in \mathbf{S}^1(K, A)$. Concerning this, we have the following lemma.

3.22. LEMMA. Let $f \in \mathbf{S}^1(A, B)$.

(1) f is faithful if and only if for any $K \in \mathbf{S}^0$ and $k \in \mathbf{S}^1(K, A)$,

$$- \circ f : \mathbf{S}^2(k, 0) \rightarrow \mathbf{S}^2(k \circ f, 0 \circ f) \text{ is injective.}$$

(2) f is fully faithful if and only if for any $K \in \mathbf{S}^0$ and $k \in \mathbf{S}^1(K, A)$,

$$- \circ f : \mathbf{S}^2(k, 0) \rightarrow \mathbf{S}^2(k \circ f, 0 \circ f) \text{ is bijective.}$$

PROOF. By Lemma 2.16, we have the following commutative diagram for any $g, h \in \mathbf{S}^1(K, A)$:

$$\begin{array}{ccc}
 \mathbf{S}^2(g, h) & \xrightarrow[\text{bij.}]{\Phi_{g,h}} & \mathbf{S}^2(g \otimes h^*, 0) \\
 \swarrow - \circ f & & \searrow - \circ f \\
 \mathbf{S}^2(g \circ f, h \circ f) & & \mathbf{S}^2((g \otimes h^*) \circ f, 0 \circ f) \\
 \searrow \Phi_{g \circ f, h \circ f} & \circlearrowleft & \swarrow \Theta_{g,h}^{f^b} \\
 & & \mathbf{S}^2((g \circ f) \otimes (h \circ f)^*, 0) \\
 \text{bij.} \swarrow & & \searrow \text{bij.}
 \end{array}$$

So we have

$$\begin{aligned} & - \circ f : \mathbf{S}^2(g, h) \rightarrow \mathbf{S}^2(g \circ f, h \circ f) \text{ is injective (resp. bijective)} \\ \Leftrightarrow & - \circ f : \mathbf{S}^2(g \otimes h^*, 0) \rightarrow \mathbf{S}^2((g \otimes h^*) \circ f, 0 \circ f) \text{ is injective (resp. bijective).} \end{aligned}$$

■

3.23. COROLLARY. *For any $f \in \mathbf{S}^1(A, B)$, f is faithful if and only if the following condition is satisfied:*

$$\alpha \circ f = \text{id}_{0 \circ f} \Rightarrow \alpha = \text{id}_0 \quad (\forall K \in \mathbf{S}^0, \forall \alpha \in \mathbf{S}^2(0_{K,A}, 0_{K,A})) \quad (14)$$

PROOF. If f is faithful, (14) is trivially satisfied, since we have $\text{id}_{0 \circ f} = \text{id}_0 \circ f$. To show the other implication, by Lemma 3.22, it suffices to show that $- \circ f : \mathbf{S}^2(k, 0) \rightarrow \mathbf{S}^2(k \circ f, 0 \circ f)$ is injective. For any $\alpha_1, \alpha_2 \in \mathbf{S}^2(k, 0)$ which satisfy $\alpha_1 \circ f = \alpha_2 \circ f$, we have $(\alpha_1^{-1} \cdot \alpha_2) \circ f = (\alpha_1 \circ f)^{-1} \cdot (\alpha_2 \circ f) = \text{id}_{0 \circ f}$. From the assumption we obtain $\alpha_1^{-1} \cdot \alpha_2 = \text{id}_0$, i.e. $\alpha_1 = \alpha_2$. ■

The next corollary immediately follows from Lemma 3.22.

3.24. COROLLARY. *For any $f \in \mathbf{S}^1(A, B)$, f is fully faithful if and only if for any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, A)$, and any $\sigma \in \mathbf{S}^2(k \circ f, 0)$, there exists unique $\tau \in \mathbf{S}^2(k, 0)$ such that $\sigma = (\tau \circ f) \cdot f_I^b$.*

3.25. COROLLARY. *For any $f \in \mathbf{S}^1(A, B)$, the following are equivalent:*

- (1) f is fully faithful.
- (2) $\text{Ker}(f) = 0$.

PROOF. (1) \Rightarrow (2)

Since f is fully faithful, for any $k \in \mathbf{S}^1(K, A)$ and $\varepsilon \in \mathbf{S}^2(k \circ f, 0)$, there exists a 2-cell $\underline{\varepsilon} \in \mathbf{S}^2(0_{K,A}, k)$ such that $(\underline{\varepsilon} \circ f) = (0 \circ f_I^b) \cdot 0_I^\sharp \cdot \varepsilon^{-1} = (0 \circ f_I^b) \cdot \varepsilon^{-1}$, and the existence of a factorization is shown. To show the uniqueness of the factorization, it suffices to show that for any other factorization $(\underline{k}', \underline{\varepsilon}')$ with $(\underline{\varepsilon}' \circ f) \cdot \varepsilon = (\underline{k}' \circ f_I^b) \cdot (\underline{k}')_I^\sharp$, the unique 2-cell $\tau \in \mathbf{S}^2(\underline{k}', 0)$ (see condition (a3-2) in Definition 3.1) satisfies $(\tau \circ 0) \cdot \underline{\varepsilon} = \underline{\varepsilon}'$. Since f is faithful, this is equivalent to $(\tau \circ 0 \circ f) \cdot (\underline{\varepsilon} \circ f) \cdot \varepsilon = (\underline{\varepsilon}' \circ f) \cdot \varepsilon$, and this follows easily from $\tau \circ 0 = (\tau \circ 0) \cdot 0_I^\sharp = (\underline{k}')_I^\sharp$ and $(\tau \circ 0 \circ f) \cdot (0 \circ f_I^b) = (\underline{k}' \circ f_I^b) \cdot (\tau \circ 0)$. (see Corollary 3.5.)

(2) \Rightarrow (1) Since $\text{Ker}(f) = [0, 0, f_I^b]$, for any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, A)$ and any $\sigma \in \mathbf{S}^2(k \circ f, 0)$, there exist $\underline{k} \in \mathbf{S}^1(K, 0)$ and $\underline{\sigma} \in \mathbf{S}^2(\underline{k} \circ 0, k)$ such that $(\underline{\sigma} \circ f) \cdot \sigma = (\underline{k} \circ f_I^b) \cdot (\underline{k})_I^\sharp$. Thus $\tau := \underline{\sigma}^{-1} \cdot \underline{k}_I^\sharp$ satisfies $\sigma = (\tau \circ f) \cdot f_I^b$. If there exists another $\tau' \in \mathbf{S}^2(k, 0)$ satisfying $\sigma = (\tau' \circ f) \cdot f_I^b$, then by the universality of the kernel, there exists $v \in \mathbf{S}^2(\underline{k}, 0)$ such that $(v \circ 0) \cdot \tau'^{-1} = \underline{\tau}$. Since $v \circ 0 = \underline{k}_I^\sharp$ by (11), we obtain $\tau = \tau'$. Thus τ is uniquely determined. ■


3.26. PROPOSITION. For any $f \in \mathbf{S}^1(A, B)$, the following are equivalent.

- (1) f is an equivalence.
- (2) f is cofaithful and fully faithful.
- (3) f is faithful and fully cofaithful.

PROOF. Since (1) \Leftrightarrow (3) is the dual of (1) \Leftrightarrow (2), we show only (1) \Leftrightarrow (2).


(1) \Rightarrow (2) : trivial.

(2) \Rightarrow (1) : Since f is cofaithful, we have $f = \text{cok}(\ker(f))$, $\text{Cok}(k(f)) = [B, f, \varepsilon_f]$. On the other hand, since f is fully faithful, we have $\text{Ker}(f) = [0, 0, f_I^b]$, and so we have $\text{Cok}(k(f)) = [A, \text{id}_A, \text{id}_0]$. And by the uniqueness (up to an equivalence) of the cokernel, there is an equivalence from A to B , which is equivalent to f . Thus, f becomes an equivalence.



■


3.27. LEMMA. Let $f : A \rightarrow B$ be a faithful 1-cell in \mathbf{S} . Then, for any $K \in \mathbf{S}^0$ and $k \in \mathbf{S}^1(K, 0)$, we have $\mathbf{S}^2(k \circ 0_{0, \text{Ker}(f)}, 0_{K, \text{Ker}(f)}) = \{k_I^\sharp\}$.



PROOF. For any $\sigma \in \mathbf{S}^2(k \circ 0_{0, \text{Ker}(f)}, 0_{K, \text{Ker}(f)})$, we can show $((\sigma \circ k(f)) \cdot k(f)_I^b) \circ f = ((k \circ k(f)_I^b) \cdot k_I^\sharp) \circ f$. By the faithfulness of f , we have $(\sigma \circ k(f)) \cdot k(f)_I^b = (k \circ k(f)_I^b) \cdot k_I^\sharp$. Thus, we have $\sigma \circ k(f) = k_I^\sharp \circ k(f)$. By the faithfulness of $k(f)$, we obtain $\sigma = k_I^\sharp$. ■

3.28. COROLLARY. $f : A \rightarrow B$ is faithful if and only if $\text{Ker}(0_{0,A}, f_I^b) = 0$.

PROOF. Since there is a factorization diagram with $(0_{0, \text{Ker}(f)} \circ \varepsilon_f) \cdot (0_{0, \text{Ker}(f)})_I^\sharp = (k(f)_I^b \circ f) \cdot f_I^b$




(see (a3-2) in Definition 3.1) we have $\text{Ker}(0_{0,A}, f_I^b) = \text{Ker}(0_{0, \text{Ker}(f)})$ by Proposition 3.20. So, it suffices to show $\text{Ker}(0_{0, \text{Ker}(f)}) = 0$. For any $K \in \mathbf{S}^0$ and $k \in \mathbf{S}^1(K, 0)$, we have

$\mathbf{S}^2(k \circ 0_{0, \text{Ker}(f)}, 0_{K, \text{Ker}(f)}) = \{k_I^\sharp\}$ by the Lemma 3.27. So $0_{0, \text{Ker}(f)}$ becomes fully faithful, and thus $\text{Ker}(0_{0, \text{Ker}(f)}) = 0$.

Conversely, assume $\text{Ker}(0_{0,A}, f_I^b) = 0$. For any $K \in \mathbf{S}^0$ and $\alpha \in \mathbf{S}^2(0_{K,A}, 0_{K,A})$ satisfying $\alpha \circ f = \text{id}_{0 \circ f}$, we show $\alpha = \text{id}_0$ (Corollary 3.23).

By $\alpha \circ f = \text{id}_{0 \circ f}$, α is compatible with f_I^b :



So there exist $k \in \mathbf{S}^1(K, 0)$ and $\varepsilon \in \mathbf{S}^2(k \circ \text{id}_0, 0_{K,0})$ satisfying

$$(\varepsilon \circ 0_{0,A}) \cdot \alpha = (k \circ \text{id}_0) \cdot k_I^\sharp.$$

Since $\varepsilon \circ 0_{0,A} = k_I^\sharp$ by (5) and (10), we obtain $\alpha = \text{id}_0$. ■

In any relatively exact 2-category \mathbf{S} , the difference kernel of any pair of 1-cells $g, h : A \rightarrow B$ always exists. More precisely, we have the following proposition:

3.29. PROPOSITION. *For any $g, h \in \mathbf{S}^1(A, B)$, if we take the kernel $\text{Ker}(g \otimes h^*) = [\text{Ker}(g \otimes h^*), k, \varepsilon]$ of $g \otimes h^*$ and put $\tilde{\varepsilon} := \Psi_{k \circ g, k \circ h}(\Theta_{g,h}^{k^\sharp}(\varepsilon \cdot k_I^{\sharp-1})) \in \mathbf{S}^2(k \circ g, k \circ h)$, then $(\text{Ker}(g \otimes h^*), k, \tilde{\varepsilon})$ is the difference kernel of g and h .*


PROOF. For any $K \in \mathbf{S}^0$ and $\ell \in \mathbf{S}^1(K, A)$, there exists a natural isomorphism (Lemma 2.16)

$$\begin{array}{ccc} \mathbf{S}^2(\ell \circ (g \otimes h^*), 0) & \xrightarrow{\quad} & \mathbf{S}^2(\ell \circ g, \ell \circ h) \\ \Downarrow \sigma & & \Downarrow \tilde{\sigma} \\ \sigma & \xrightarrow{\quad} & \tilde{\sigma} := \Psi_{\ell \circ g, \ell \circ h}(\Theta_{g,h}^{\ell^\sharp}(\sigma \cdot \ell_I^\sharp)). \end{array}$$

So, to give a 2-cell $\sigma \in \mathbf{S}^2(\ell \circ (g \otimes h^*), 0)$ is equivalent to give a 2-cell $\tilde{\sigma} \in \mathbf{S}^2(\ell \circ g, \ell \circ h)$. And, by using Remark 3.4 and Corollary 3.5, the usual compatibility argument shows the proposition. ■

In any relatively exact 2-category \mathbf{S} , the pullback of any pair of morphisms $f_i : A_i \rightarrow B$ ($i = 1, 2$) always exists. More precisely, we have the following proposition:

3.30. PROPOSITION. *For any $f_i \in \mathbf{S}^1(A_i, B)$ ($i = 1, 2$), if we take the product of A_1 and A_2 ($A_1 \times A_2, p_1, p_2$), and take the difference kernel (D, d, φ) of $p_1 \circ f_1$ and $p_2 \circ f_2$*



then, $(D, d \circ p_1, d \circ p_2, \varphi)$ is the pullback of f_1 and f_2 .

PROOF OF CONDITION (b1) (IN DEFINITION 2.4). For any $X \in \mathbf{S}^0$, $g_i \in \mathbf{S}^1(X, A_i)$ ($i = 1, 2$) and $\eta \in \mathbf{S}^2(g_1 \circ f_1, g_2 \circ f_2)$, by the universality of $A_1 \times A_2$, there exist $g \in \mathbf{S}^1(X, A_1 \times A_2)$ and $\xi_i \in \mathbf{S}^2(d \circ p_i, g_i)$ ($i = 1, 2$). Applying the universality of the difference kernel to the 2-cell

$$\zeta := (\xi_1 \circ f_1) \cdot \eta \cdot (\xi_2^{-1} \circ f_2) \in \mathbf{S}^2(g \circ p_1 \circ f_1, g \circ p_2 \circ f_2), \quad (15)$$

we see there exist $\underline{g} \in \mathbf{S}^1(X, D)$ and $\underline{\zeta} \in \mathbf{S}^2(\underline{g} \circ d, g)$

$$\begin{array}{ccccc} & X & & & \\ & \swarrow \underline{g} & & \searrow g & \\ D & \xrightarrow{d} & A_1 \times A_2 & \xrightarrow[p_2 \circ f_2]{p_1 \circ f_1} & B \\ & & \nearrow \zeta & & \end{array} \quad (16)$$

such that

$$(\underline{g} \circ \varphi) \cdot (\underline{\zeta} \circ p_2 \circ f_2) = (\underline{\zeta} \circ p_1 \circ f_1) \cdot \zeta. \quad (17)$$

By (15) and (17), we have $(\underline{g} \circ \varphi) \cdot (((\underline{\zeta} \circ p_2) \cdot \xi_2) \circ f_2) = (((\underline{\zeta} \circ p_1) \cdot \xi_1) \circ f_1) \cdot \eta$, and thus condition (b1) is satisfied.

$$\begin{array}{ccccc} & & (\underline{\zeta} \circ p_1) \cdot \xi_1 & & \\ & & \cdots & & \\ & g_1 & \nearrow & A_1 & \searrow f_1 \\ X & \xrightarrow{g} & D & \xrightarrow{\varphi} & B \\ & \searrow g_2 & \nearrow & A_2 & \swarrow f_2 \\ & & (\underline{\zeta} \circ p_2) \cdot \xi_2 & & \end{array}$$

proof of condition (b2)

If we take $h \in \mathbf{S}^1(X, D)$ and $\eta_i \in \mathbf{S}^2(h \circ d \circ p_i, g_i)$ ($i = 1, 2$) which satisfy $(h \circ \varphi) \cdot (\eta_2 \circ f_2) = (\eta_1 \circ f_1) \cdot \eta$, then by the universality of $A_1 \times A_2$, there exists a unique 2-cell $\kappa \in \mathbf{S}^2(h \circ d, g)$ such that

$$(\kappa \circ p_i) \cdot \xi_i = \eta_i \quad (i = 1, 2). \quad (18)$$

Then, κ becomes compatible with φ and ζ , i.e. $(h \circ \varphi) \cdot (\kappa \circ p_2 \circ f_2) = (\kappa \circ p_1 \circ f_1) \cdot \zeta$. So, comparing this with factorization (16), by the universality of the difference kernel, we see there exists a unique 2-cell $\chi \in \mathbf{S}^2(h, \underline{g})$ which satisfies

$$(\chi \circ d) \cdot \underline{\zeta} = \kappa \quad (19)$$

Then we have $(\chi \circ d \circ p_i) \cdot (\underline{\zeta} \circ p_i) \cdot \xi_i = (\kappa \circ p_i) \cdot \xi_i = \eta_i$ ($i = 1, 2$). Thus χ is the desired 2-cell in condition (b2), and the uniqueness of such a χ follows from the uniqueness of κ and χ which satisfy (18) and (19). \blacksquare

By the universality of the pullback, we have the following remark:

3.31. REMARK. Let

$$\begin{array}{ccccc}
 & & A_2 & & \\
 & f'_1 \nearrow & & f_2 \searrow & \\
 A_1 \times_B A_2 & & A_2 & \Downarrow \xi & B \\
 & f'_2 \searrow & & f_1 \nearrow & \\
 & & A_1 & &
 \end{array} \tag{20}$$

be a pull-back diagram. Then, for any $K \in \mathbf{S}^0$, $g, h \in \mathbf{S}^1(K, A_1 \times_B A_2)$ and $\alpha, \beta \in \mathbf{S}^2(g, h)$, we have

$$\alpha \circ f'_i = \beta \circ f'_i \ (i = 1, 2) \implies \alpha = \beta.$$

PROOF. To the diagram

$$\begin{array}{ccccc}
 & & A_2 & & \\
 & g \circ f'_1 \nearrow & & f_2 \searrow & \\
 K & & A_2 & \Downarrow g \circ \xi & B \\
 & g \circ f'_2 \searrow & & f_1 \nearrow & \\
 & & A_1 & &
 \end{array} ,$$

the following diagram gives a factorization which satisfies condition (b1) in Definition 2.4.

$$\begin{array}{ccccc}
 & & g \circ f'_1 \nearrow & & A_2 & & f_2 \searrow & & \\
 & & \circ & & & & & & \\
 K & \xrightarrow{g} & A_1 \times_B A_2 & \xrightarrow{f'_1} & A_2 & \xrightarrow{f_2} & B \\
 & & \circ & & \Downarrow \xi & & \\
 & & & & A_1 & \xrightarrow{f_1} & \\
 & & g \circ f'_2 \searrow & & & &
 \end{array}$$

Since each of $\text{id}_g : g \implies g$ and $\alpha \circ \beta^{-1} : g \implies g$ gives a 2-cell which satisfies condition (b2), we have $\alpha \circ \beta^{-1} = \text{id}$ by the uniqueness. Thus $\alpha = \beta$. ■

3.32. PROPOSITION. (See also Proposition 5.12.) Let (20) be a pull-back diagram. We have

- (1) f_1 : faithful $\implies f'_1$: faithful.
- (2) f_1 : fully faithful $\implies f'_1$: fully faithful.
- (3) f_1 : cofaithful $\implies f'_1$: cofaithful.

PROOF. proof of (1) By Corollary 3.23, it suffices to show $\alpha \circ f'_1 = \text{id}_{0 \circ f'_1} \implies \alpha = \text{id}_0$ for any $K \in \mathbf{S}^0$ and $\alpha \in \mathbf{S}^2(0_{K, A_1 \times_B A_2}, 0_{K, A_1 \times_B A_2})$. Since $(0 \circ \xi) \cdot (\alpha \circ f'_2 \circ f_1) = (\alpha \circ f'_1 \circ f_2) \cdot (0 \circ \xi) = \text{id}_{0 \circ f'_1 \circ f_2} \cdot (0 \circ \xi) = 0 \circ \xi$, we have $\alpha \circ f'_2 \circ f_1 = \text{id}_{0 \circ f'_2 \circ f_1} = \text{id}_{0 \circ f'_2} \circ f_1$. Since f_1 is faithful, we obtain $\alpha \circ f'_2 = \text{id}_{0 \circ f'_2}$. Thus, we have $\alpha \circ f'_i = \text{id}_{0 \circ f'_i} = \text{id}_0 \circ f'_i \ (i = 1, 2)$. By Remark 3.31, this implies $\alpha = \text{id}_0$.

proof of (2) By (1), f'_1 is already faithful. By Corollary 3.23, it suffices to show that for any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, A_1 \times_B A_2)$ and any $\sigma \in \mathbf{S}^2(k \circ f'_1, 0)$, there exists a unique 2-cell $\kappa \in \mathbf{S}^2(k, 0)$ such that $\sigma = (\kappa \circ f'_1) \cdot (f'_1)_I^b$. Since f_1 is fully faithful, for any $K \in \mathbf{S}^0$,

$k \in \mathbf{S}^1(K, A_1 \times_B A_2)$ and any $\sigma \in \mathbf{S}^2(k \circ f'_1, 0)$, there exists $\tau \in \mathbf{S}^2(k \circ f'_2, 0)$ such that $(\tau \circ f_1) \cdot (f_1)_I^\flat = (k \circ \xi^{-1}) \cdot (\sigma \circ f_2) \cdot (f_2)_I^\flat$. Then, for the diagram

$$\begin{array}{ccccc}
 & & A_1 & & \\
 & \nearrow 0 & & \searrow f_1 & \\
 K & & \parallel & & B \\
 & \searrow 0 & (f_1)_I^\flat \cdot (f_2)_I^{\flat^{-1}} & \nearrow f_2 & \\
 & & A_2 & &
 \end{array}$$

each of the factorizations

$$\begin{array}{ccc}
 \begin{array}{ccccc}
 & & 0 & & \\
 & \nearrow & & \searrow & \\
 K & \xrightarrow{k} & A_1 \times_B A_2 & \xrightarrow{f_2} & B \\
 & \searrow & \Downarrow \xi & \nearrow & \\
 & & A_1 & &
 \end{array}
 &
 &
 \begin{array}{ccccc}
 & & 0 & & \\
 & \nearrow & & \searrow & \\
 K & \xrightarrow{k} & A_1 \times_B A_2 & \xrightarrow{f_2} & B \\
 & \searrow & \Downarrow \xi & \nearrow & \\
 & & A_1 & &
 \end{array}
 \end{array}$$

satisfies condition (b1) in Definition 2.4. So there exists a 2-cell $\kappa \in \mathbf{S}^2(k, 0)$ such that $\sigma = (\kappa \circ f'_1) \cdot (f'_1)_I^\flat$. Uniqueness of such κ follows from the faithfulness of f'_1 .

proof of (3) Let $(A_1 \times A_2, p_1, p_2)$ be the product of A_1 and A_2 . For $\text{id}_{A_1} \in \mathbf{S}^1(A_1, A_1)$ and $0 \in \mathbf{S}^1(A_1, A_2)$, by the universality of $A_1 \times A_2$, there exist $i_1 \in \mathbf{S}^1(A_1, A_1 \times A_2)$, $\xi_1 \in \mathbf{S}^2(i_1 \circ p_1, \text{id}_{A_1})$ and $\xi_2 \in \mathbf{S}^2(i_2 \circ p_2, 0)$. Similarly, there is a 1-cell $i_2 \in \mathbf{S}^1(A_2, A_1 \times A_2)$ such that there are equivalences $i_2 \circ p_2 \simeq \text{id}_{A_2}$, $i_2 \circ p_1 \simeq 0$. If we put $t := (p_1 \circ f_1) \otimes (p_2 \circ f_2)^*$, then by Proposition 3.29 and 3.30, we have $A_1 \times_B A_2 = \text{Ker}(t)$. So we may assume $\text{Ker}(t) = [A_1 \times_B A_2, d, \varepsilon_t]$ and $f'_1 = d \circ p_2$.

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \nearrow & \uparrow \varepsilon_t & \searrow & \\
 A_1 \times_B A_2 & \xrightarrow{d} & A_1 \times A_2 & \xrightarrow{t} & B
 \end{array}$$

Since $i_1 \circ t$ and f_1 are equivalent;


$$i_1 \circ t \simeq (i_1 \circ p_1 \circ f_1) \otimes (i_1 \circ p_2 \circ f_2^*) \simeq (\text{id}_{A_1} \circ f_1) \otimes (0 \circ f_2^*) \simeq f_1,$$

by the cofaithfulness of f_1 , it follows that t is cofaithful. Thus, we have $B = \text{Cok}(\text{ker}(t))$, i.e. $\text{Cok}(d) = [B, t, \varepsilon_t]$. By (the dual of) Corollary 3.23, it suffices to show $f'_1 \circ \alpha = \text{id}_{f'_1 \circ 0} \Rightarrow \alpha = \text{id}_0$ for any $C \in \mathbf{S}^0$ and any $\alpha \in \mathbf{S}^2(0_{A_2, C}, 0_{A_2, C})$. For the 2-cell $(d \circ p_2)_I^\sharp \in \mathbf{S}^2(d \circ p_2 \circ 0_{A_2, C}, 0)$ (see the following diagram), by the universality of $\text{Cok}(d)$, there exist $u \in \mathbf{S}^1(B, C)$ and $\gamma \in \mathbf{S}^2(t \circ u, p_2 \circ 0)$ such that $(d \circ \gamma) \cdot (d \circ p_2)_I^\sharp = (\varepsilon_t \circ u) \cdot u_I^\flat$. Thus, if we put $\gamma' := \gamma \cdot (p_2 \circ \alpha)$, we have

$$\begin{aligned}
 (d \circ \gamma') \cdot (d \circ p_2)_I^\sharp &= (d \circ \gamma) \cdot (d \circ p_2 \circ \alpha) \cdot (d \circ p_2)_I^\sharp \\
 &= (d \circ \gamma) \cdot (f'_1 \circ \alpha) \cdot (d \circ p_2)_I^\sharp = (\varepsilon_t \circ u) \cdot u_I^\flat.
 \end{aligned}$$


So, γ and $\gamma' \in \mathbf{S}^2(t \circ u, p_2 \circ 0)$ give two factorization of $p_2 \circ 0$ compatible with ε_t and $(d \circ p_2)_I^\sharp$. By the universality of $\text{Cok}(d) = [B, t, \varepsilon_t]$, there exists a unique 2-cell $\beta \in \mathbf{S}^2(u, u)$ such that

$$(t \circ \beta) \cdot \gamma = \gamma'. \tag{21}$$



Then we have $(i_1 \circ t \circ \beta) \cdot (i_1 \circ \gamma) = i_1 \circ \gamma' = (i_1 \circ \gamma) \cdot (i_1 \circ p_2 \circ \alpha) = (i_1 \circ \gamma) \cdot (\xi_2 \circ 0) \cdot (0 \circ \alpha) \cdot (\xi_2^{-1} \circ 0) = (i_1 \circ \gamma)$, and thus, $(i_1 \circ t) \circ \beta = \text{id}_{i_1 \circ t \circ u}$. Since $i_1 \circ t \simeq f_1$ is cofaithful, we obtain $\beta = \text{id}_u$. So, by (21), we have $\gamma = \gamma' = \gamma \cdot (p_2 \circ \alpha)$, and consequently $p_2 \circ \alpha = \text{id}_{p_2 \circ 0}$. Since p_2 is cofaithful (because $i_2 \circ p_2 \simeq \text{id}_{A_2}$ is cofaithful), we obtain $\alpha = \text{id}_0$. ■

3.33. PROPOSITION. *Consider diagram (12) in \mathbf{S} . If we take $\text{Ker}(f, \varphi) = [\text{Ker}(f, \varphi), \ell, \varepsilon]$, then by the universality of $\text{Ker}(f) = [\text{Ker}(f), k(f), \varepsilon_f]$, ℓ factors uniquely up to an equivalence as*



where $(\underline{\varepsilon} \circ f) \cdot \varepsilon = (\underline{\ell} \circ \varepsilon_f) \cdot (\underline{\ell})_I^\sharp$. Then, $\underline{\ell}$ becomes fully faithful.

PROOF. Since $\underline{\ell} \circ k(f)$ is equivalent to a faithful 1-cell ℓ , so $\underline{\ell}$ becomes faithful. For any $K \in \mathbf{S}^0$, $k \in \mathbf{S}^1(K, \text{Ker}(f, \varphi))$ and $\sigma \in \mathbf{S}^2(k \circ \underline{\ell}, 0)$, if we put $\sigma' := (k \circ \underline{\varepsilon}^{-1}) \cdot (\sigma \circ k(f)) \cdot k(f)_I^\sharp \in \mathbf{S}^2(k \circ \ell, 0)$, then σ' becomes compatible with ε . So, by Lemma 3.19, there exists $\tau \in \mathbf{S}^2(k, 0)$ such that $\sigma' = (\tau \circ \ell) \cdot \ell_I^\sharp$, i.e.

$$(k \circ \underline{\varepsilon}^{-1}) \cdot (\sigma \circ k(f)) \cdot (k(f))_I^\sharp = (\tau \circ \ell) \cdot \ell_I^\sharp. \tag{22}$$

Now, since $(k \circ \underline{\varepsilon}) \cdot (\tau \circ \ell) \cdot \ell_I^\sharp = (\tau \circ \underline{\ell} \circ k(f)) \cdot (\underline{\ell} \circ k(f))_I^\sharp$ by Corollary 3.5, (22) is equivalent to $(\sigma \circ k(f)) \cdot (k(f))_I^\sharp = (\tau \circ \underline{\ell} \circ k(f)) \cdot (\underline{\ell}_I^\sharp \circ k(f)) \cdot (k(f))_I^\sharp$.

Thus, we obtain $\sigma \circ k(f) = ((\tau \circ \underline{\ell}) \cdot \underline{\ell}_I^\sharp) \circ k(f)$. Since $k(f)$ is faithful, it follows that $\sigma = (\tau \circ \underline{\ell}) \cdot \underline{\ell}_I^\sharp$. Uniqueness of such τ follows from the faithfulness of $\underline{\ell}$. Thus $\underline{\ell}$ becomes fully faithful by Corollary 3.24. ■

4. Existence of proper factorization systems

4.1. DEFINITION. *For any $A, B \in \mathbf{S}^0$ and $f \in \mathbf{S}^1(A, B)$, we define its image as $\text{Ker}(\text{cok}(f))$.*

4.2. **REMARK.** By the universality of the kernel, there exist $i(f) \in \mathbf{S}^1(A, \text{Im}(f))$ and $\iota \in \mathbf{S}^2(i(f) \circ k(c(f)), f)$ such that $(\iota \circ c(f)) \cdot \pi_f = (i(f) \circ \varepsilon_{c(f)}) \cdot i(f)_I^\sharp$. Coimage of f is defined dually, and we obtain a factorization through $\text{Coim}(f)$.

4.3. **PROPOSITION.** [1st factorization] *For any $f \in \mathbf{S}^1(A, B)$, the factorization $\iota : i(f) \circ k(c(f)) \implies f$ through $\text{Im}(f)$*

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ & \searrow i(f) & \nearrow k(c(f)) \\ & \Uparrow \iota & \\ & \text{Im}(f) & \end{array}$$

satisfies the following properties:

(A) $i(f)$ is fully cofaithful and $k(c(f))$ is faithful.

(B) For any factorization $\eta : i \circ m \implies f$ where m is faithful, following (b1) and (b2) hold:

(b1) There exist $t \in \mathbf{S}^1(\text{Im}(f), C)$, $\zeta_m \in \mathbf{S}^2(t \circ m, k(c(f)))$, $\zeta_i \in \mathbf{S}^2(i(f) \circ t, i)$

$$\begin{array}{ccccc} & & C & & \\ & i \nearrow & \uparrow & \searrow m & \\ A & \zeta_i \Uparrow t & & \Downarrow \zeta_m & B \\ & \searrow i(f) & \uparrow & \nearrow k(c(f)) & \\ & & \text{Im}(f) & & \end{array}$$

such that $(i(f) \circ \zeta_m) \cdot \iota = (\zeta_i \circ m) \cdot \eta$.

(b2) If both (t, ζ_m, ζ_i) and (t', ζ'_m, ζ'_i) satisfy (b1), then there is a unique 2-cell $\kappa \in \mathbf{S}^2(t, t')$ such that $(i(f) \circ \kappa) \cdot \zeta'_i = \zeta_i$ and $(\kappa \circ m) \cdot \zeta'_m = \zeta_m$.

Dually, we obtain the following proposition for the coimage factorization.

4.4. **PROPOSITION.** [2nd factorization] *For any $f \in \mathbf{S}^1(A, B)$, the factorization $\mu : c(k(f)) \circ j(f) \implies f$ through $\text{Coim}(f)$*


$$\begin{array}{ccc} & \text{Coim}(f) & \\ c(k(f)) \nearrow & \Downarrow \mu & \searrow j(f) \\ A & \xrightarrow{f} & B \end{array}$$

satisfies the following properties:

(A) $j(f)$ is fully faithful and $c(k(f))$ is cofaithful.

(B) For any factorization $\nu : e \circ j \implies f$ where e is cofaithful, following (b1) and (b2) (the dual of the conditions in Proposition 4.3) hold:

(b1) *There exists $s \in \mathbf{S}^1(C, \text{Coim}(f))$, $\zeta_e \in \mathbf{S}^2(e \circ s, c(k(f)))$, and $\zeta_j \in \mathbf{S}^2(s \circ j(f), j)$*




such that $(e \circ \zeta_j) \cdot \nu = (\zeta_e \circ j(f)) \cdot \mu$.

(b2) *If both (s, ζ_e, ζ_j) and (s', ζ'_e, ζ'_j) satisfy (b1), then there is a unique 2-cell $\lambda \in \mathbf{S}^2(t, t')$ such that $(\lambda \circ j(f)) \cdot \zeta'_j = \zeta_j$ and $(e \circ \lambda) \cdot \zeta'_e = \zeta_e$.*


In the rest of this section, we show Proposition 4.3.

4.5. LEMMA. *For any $f \in \mathbf{S}^1(A, B)$, $i(f)$ is cofaithful.*

PROOF. It suffices to show that for any $C \in \mathbf{S}^0$ and $\alpha \in \mathbf{S}^2(0_{\text{Im}(f), C}, 0_{\text{Im}(f), C})$



we have $i(f) \circ \alpha = \text{id}_{i(f) \circ 0} \implies \alpha = \text{id}_0$. Take the pushout of $k(c(f))$ and $0_{\text{Im}(f), C}$



and put

$$\xi_1 := \xi \cdot (\xi_1 \circ f_2) \cdot \eta = (g \circ \xi) \cdot (\xi_2 \circ f_1)(i_C)_I^b = (k(c(f)) \circ i_B \xrightarrow{\xi} 0 \circ i_C \xrightarrow{(i_C)_I^b} 0)$$


$$\xi_2 := \xi \cdot (\alpha \circ i_C) \cdot (i_C)_I^b = (k(c(f)) \circ i_B \xrightarrow{\xi} 0 \circ i_C \xrightarrow{\alpha \circ i_C} 0 \circ i_C \xrightarrow{(i_C)_I^b} 0).$$

Then, since i_C is faithful by (the dual of) Lemma 3.32, we have

$$\alpha = \text{id}_0 \iff \alpha \circ i_C = \text{id}_{0 \circ i_C} \iff \xi \cdot (\alpha \circ i_C) \cdot (i_C)_I^b = \xi \cdot \text{id}_{0 \circ i_C} \cdot (i_C)_I^b \iff \xi_1 = \xi_2.$$

So, it suffices to show $\xi_1 = \xi_2$. For each $i = 1, 2$, since $\text{Cok}(k(c(f))) = [\text{Cok}(f), c(f), \varepsilon_{c(f)}]$, there exist $e_i \in \mathbf{S}^1(\text{Cok}(f), C \amalg_{\text{Im}(f)} B)$ and $\varepsilon_i \in \mathbf{S}^2(c(f) \circ e_i, i_B)$ such that


$$(k(c(f)) \circ \varepsilon_i) \cdot \xi_i = (\varepsilon_{c(f)} \circ e_i) \cdot (e_i)_I^b. \tag{23}$$



Since by assumption $i(f) \circ \alpha = \text{id}_{i(f) \circ 0}$, we have

$$\begin{aligned} i(f) \circ \xi_2 &= (i(f) \circ \xi) \cdot (i(f) \circ \alpha \circ i_C) \cdot (i(f) \circ (i_C)_I^b) \\ &= (i(f) \circ \xi) \cdot (\text{id}_{i(f) \circ 0 \circ i_C}) \cdot (i(f) \circ (i_C)_I^b) = i(f) \circ \xi_1. \end{aligned}$$

So, if we put $\varpi := (\iota^{-1} \circ i_B) \cdot (i(f) \circ \xi_i) \cdot (i(f))_I^\sharp \in \mathbf{S}^2(f \circ i_B, 0)$, this doesn't depend on $i = 1, 2$. We can show easily $(f \circ \varepsilon_i) \cdot \varpi = (\pi_f \circ e_i) \cdot (e_i)_I^b$ ($i = 1, 2$). Thus (e_1, ε_1) and (e_2, ε_2) are two factorizations of i_B compatible with ϖ and π_f .



By the universality of $\text{Cok}(f)$, there exists a 2-cell $\beta \in \mathbf{S}^2(e_1, e_2)$ such that $(c(f) \circ \beta) \cdot \varepsilon_2 = \varepsilon_1$, and thus we have $\varepsilon_1^{-1} = \varepsilon_2^{-1} \cdot (c(f) \circ \beta^{-1})$. So, by (23), we have


$$\begin{aligned} \xi_1 &= (k(c(f)) \circ \varepsilon_1^{-1}) \cdot (\varepsilon_{c(f)} \circ e_1) \cdot (e_1)_I^b \\ &= (k(c(f)) \circ \varepsilon_2^{-1}) \cdot (k(c(f)) \circ c(f) \circ \beta^{-1}) \cdot (\varepsilon_{c(f)} \circ e_1) \cdot (e_1)_I^b \\ &\underset{9}{=} (k(c(f)) \circ \varepsilon_2^{-1}) \cdot (\varepsilon_{c(f)} \circ e_2) \cdot (e_2)_I^b = \xi_2. \end{aligned}$$

■

4.6. LEMMA. Let $f \in \mathbf{S}^1(A, B)$. Let $\iota : i(f) \circ k(c(f)) \Rightarrow f$ be the factorization of f through $\text{Im}(f)$ as before. If we are given a factorization $\eta : i \circ m \Rightarrow f$ of f where $i \in \mathbf{S}^1(A, C)$, $m \in \mathbf{S}^1(C, B)$ and m is faithful, then there exist $t \in \mathbf{S}^1(\text{Im}(f), C)$, $\zeta_i \in \mathbf{S}^2(i(f) \circ t, i)$ and $\zeta_m \in \mathbf{S}^2(t \circ m, k(c(f)))$ such that $(\zeta_i \circ m) \cdot \eta = (i(f) \circ \zeta_m) \cdot \iota$.

PROOF. By the universality of $\text{Cok}(f)$, for $\pi := (\eta^{-1} \circ c(m)) \cdot (i \circ \pi_m) \cdot i_I^\sharp \in \mathbf{S}^2(f \circ c(m), 0)$, there exist $\bar{m} \in \mathbf{S}^1(\text{Cok}(f), \text{Cok}(m))$ and $\bar{\eta} \in \mathbf{S}^2(c(f) \circ \bar{m}, c(m))$ such that

$$(f \circ \bar{\eta}) \cdot \pi = (\pi_f \circ \bar{m}) \cdot (\bar{m})_I^b. \tag{24}$$



Since m is faithful by assumption, it follows $\text{Ker}(c(m)) = [C, m, \pi_m]$. By the universality of $\text{Ker}(c(m))$, for the 2-cell

$$\zeta := (k(c(f)) \circ \bar{\eta}^{-1}) \cdot (\varepsilon_{c(f)} \circ \bar{m}) \cdot (\bar{m})_I^b \in \mathbf{S}^2(k(c(f)) \circ c(m), 0), \quad (25)$$


there exist $t \in \mathbf{S}^1(\text{Im}(f), C)$ and $\zeta_m \in \mathbf{S}^2(t \circ m, k(c(f)))$ such that $(\zeta_m \circ c(m)) \cdot \zeta = (t \circ \pi_m) \cdot t_I^\sharp$.

If we put $\bar{\zeta} := (i(f) \circ \zeta_m) \cdot \iota$, then the following claim holds:

4.7. CLAIM. *Each of the two factorizations of f through $\text{Ker}(c(m))$*

$$\eta : i \circ m \implies f \quad \text{and} \quad \bar{\zeta} : i(f) \circ t \circ m \implies f$$

is compatible with π_m and π .



If the above claim is proven, then by the universality of $\text{Ker}(c(m)) = [C, m, \pi_m]$, there exists a unique 2-cell $\zeta_i \in \mathbf{S}^2(i(f) \circ t, i)$ such that $(\zeta_i \circ m) \cdot \eta = \bar{\zeta}$. Thus we obtain (t, ζ_m, ζ_i) which satisfies $(\zeta_i \circ m) \cdot \eta = \bar{\zeta} = (i(f) \circ \zeta_m) \cdot \iota$, and the lemma is proven. So, we show Claim 4.7.

(a) compatibility of η with π_m, π

This follows immediately from the definition of π .

(b) compatibility of $\bar{\zeta}$ with π_m, π

We have

$$\begin{aligned} i(f) \circ \zeta &\stackrel{25}{=} (\iota \circ c(m)) \cdot (f \circ \bar{\eta}^{-1}) \cdot (\iota^{-1} \circ c(f) \circ \bar{m}) \\ &\quad \cdot (i(f) \circ \varepsilon_{c(f)} \circ \bar{m}) \cdot (i(f) \circ (\bar{m})_I^b) \\ &\stackrel{24}{=} (\iota \circ c(m)) \cdot \pi \cdot i(f)_I^{\sharp-1}. \end{aligned}$$

From this, we obtain $(i(f) \circ t \circ \pi_m) \cdot (i(f) \circ t_I^\sharp) = (\bar{\zeta} \circ c(m)) \cdot \pi \cdot i(f)_I^{\sharp-1}$. So we have

$$(\bar{\zeta} \circ c(m)) \cdot \pi = (i(f) \circ t \circ \pi_m) \cdot (i(f) \circ t_I^\sharp) \cdot i(f)_I^{\sharp-1} = (i(f) \circ t \circ \pi_m) \cdot (i(f) \circ t_I^\sharp).$$

■

4.8. LEMMA. Let $A, B, C \in \mathbf{S}^0$, $f, m, i \in \mathbf{S}^1$, $\eta \in \mathbf{S}^2$ be as in Lemma 4.6. If a triplet (t', ζ'_m, ζ'_i) (where $t' \in \mathbf{S}^1(\text{Im}(f), C)$, $\zeta'_m \in \mathbf{S}^2(t' \circ m, k(c(f)))$, $\zeta'_i \in \mathbf{S}^2(i(f) \circ t', i)$) satisfies

$$(i(f) \circ \zeta'_m) \cdot \iota = (\zeta'_i \circ m) \cdot \eta, \tag{26}$$

then ζ'_m becomes compatible with ζ and π_m (in the notation of the proof of Lemma 4.6), i.e. we have $(\zeta'_m \circ c(m)) \cdot \zeta = (t' \circ \pi_m) \cdot (t')^\sharp_I$.

4.9. REMARK. Since m is faithful, ζ'_m which satisfies (26) is uniquely determined by t' and ζ'_i if it exists.

PROOF OF LEMMA 4.8. Since we have

$$\begin{aligned} & i(f) \circ ((\zeta'_m \circ c(m)) \cdot \zeta) \\ & \stackrel{26,1}{=} (\zeta'_i \circ m \circ c(m)) \cdot (\eta \circ c(m)) \cdot (f \circ \bar{\eta}^{-1}) \cdot (\iota^{-1} \circ c(f) \circ \bar{m}) \\ & \quad \cdot (i(f) \circ \varepsilon_{c(f)} \circ \bar{m}) \cdot (i(f) \circ (\bar{m})^\flat_I) \\ & \stackrel{24}{=} ((i(f) \circ t' \circ \pi_m) \cdot (i(f) \circ (t')^\sharp_I)), \end{aligned}$$

we obtain $(\zeta'_m \circ c(m)) \cdot \zeta = (t' \circ \pi_m) \cdot (t')^\sharp_I$ by the cofaithfulness of $i(f)$. ■

4.10. COROLLARY. Let A, B, C, f, m, i, η as in Proposition 4.3. If both (t, ζ_m, ζ_i) and (t', ζ'_m, ζ'_i) satisfy (b1), then there exists a unique 2-cell $\kappa \in \mathbf{S}^2(t, t')$ such that $(i(f) \circ \kappa) \cdot \zeta'_i = \zeta_i$ and $(\kappa \circ m) \cdot \zeta'_m = \zeta_m$.

PROOF. By Lemma 4.8, there exists a 2-cell $\kappa \in \mathbf{S}^2(t, t')$ such that $(\kappa \circ m) \cdot \zeta'_m = \zeta_m$ by the universality of $\text{Ker}(c(m)) = [C, m, \pi_m]$. This κ also satisfies $\zeta_i = (i(f) \circ \kappa) \cdot \zeta'_i$, and unique by the cofaithfulness of $i(f)$. ■

Considering the case of $C = \text{Im}(f)$, we obtain the following corollary.

4.11. COROLLARY. For any $t \in \mathbf{S}^1(\text{Im}(f), \text{Im}(f))$, $\zeta_m \in \mathbf{S}^2(t \circ k(c(f)), k(c(f)))$ and $\zeta_i \in \mathbf{S}^2(i(f) \circ t, i(f))$ satisfying $(\zeta_i \circ k(c(f))) \cdot \iota = (i(f) \circ \zeta_m) \cdot \iota$, there exists a unique 2-cell $\kappa \in \mathbf{S}^2(t, \text{id}_{\text{Im}(f)})$ such that $i(f) \circ \kappa = \zeta_i$ and $\kappa \circ k(c(f)) = \zeta_m$.

Now, we can prove Proposition 4.3.

PROOF OF PROPOSITION 4.3. Since all the other is already shown, it suffices to show the following:

4.12. CLAIM. For any $C \in \mathbf{S}^0$ and any $g, h \in \mathbf{S}^1(\text{Im}(f), C)$,

$$i(f) \circ - : \mathbf{S}^2(g, h) \longrightarrow \mathbf{S}^2(i(f) \circ g, i(f) \circ h)$$

is surjective.

So, we show Claim 4.12. If we take the difference kernel of g and h ;

$$d_{(g,h)} : \text{DK}(g, h) \longrightarrow \text{Im}(f), \quad \varphi_{(g,h)} : d_{(g,h)} \circ g \implies d_{(g,h)} \circ h,$$

then by the universality of the difference kernel, for any $\beta \in \mathbf{S}^2(i(f) \circ g, i(f) \circ h)$ there exist $i \in \mathbf{S}^1(A, \text{DK}(g, h))$ and $\lambda \in \mathbf{S}^2(i \circ d_{(g,h)}, i(f))$

$$\begin{array}{ccccc}
 & & A & & \\
 & \swarrow & & \searrow & \\
 & i & & i(f) & \\
 \text{DK}(g, h) & \xrightarrow{\lambda} & \text{Im}(f) & \xrightarrow[g]{h} & C
 \end{array}$$

such that $(i \circ \varphi_{(g,h)}) \cdot (\lambda \circ h) = (\lambda \circ g) \cdot \beta$.

If we put $m := d_{(g,h)} \circ k(c(f))$, then m becomes faithful since $d_{(g,h)}$ and $k(c(f))$ are faithful. Applying Lemma 4.6 to the factorization $\eta := (\lambda \circ k(c(f))) \cdot \iota : i \circ m \implies f$, we obtain $t \in \mathbf{S}^1(\text{Im}(f), \text{DK}(g, h))$, $\zeta_m \in \mathbf{S}^2(t \circ m, k(c(f)))$ and $\zeta_i \in \mathbf{S}^2(i(f) \circ t, i)$ such that $(\zeta_i \circ m) \cdot \eta = (i(f) \circ \zeta_m) \cdot \iota$. Thus we have

$$(\zeta_i \circ d_{(g,h)} \circ k(c(f))) \cdot (\lambda \circ k(c(f))) \cdot \iota = (i(f) \circ \zeta_m) \cdot \iota.$$

So, if we put $\bar{\zeta}_i := (\zeta_i \circ d_{(g,h)}) \cdot \lambda \in \mathbf{S}^2(i(f) \circ t \circ d_{(g,h)}, i(f))$, then we have

$$(\bar{\zeta}_i \circ k(c(f))) \cdot \iota = (i(f) \circ \zeta_m) \cdot \iota.$$

By Corollary 4.11, there exists a 2-cell $\kappa \in \mathbf{S}^2(t \circ d_{(g,h)}, \text{id}_{\text{Im}(f)})$ such that $\kappa \circ k(c(f)) = \zeta_m$ and $i(f) \circ \kappa = \bar{\zeta}_i$. If we put $\alpha := (\kappa^{-1} \circ g) \cdot (t \circ \varphi_{(g,h)}) \cdot (\kappa \circ h) \in \mathbf{S}^2(g, h)$, we can show that α satisfies $i(f) \circ \alpha = \beta$. Thus $i(f) \circ - : \mathbf{S}^2(g, h) \longrightarrow \mathbf{S}^2(i(f) \circ g, i(f) \circ h)$ is surjective. ■

4.13. REMARK. In condition (B) of Proposition 4.3, if moreover i is fully cofaithful, then t becomes fully cofaithful since i and $i(f)$ are fully cofaithful. On the other hand, t is faithful since $k(c(f))$ is faithful. So, in this case t becomes an equivalence by Proposition 3.26.

Together with Corollary 4.11, we can show easily the following corollary:

4.14. COROLLARY. For any $f \in \mathbf{S}^1(A, B)$, the following (b1) and (b2) hold:
 (b1) If in the factorizations

$$\begin{array}{ccc}
 & C & \\
 i \nearrow & \Downarrow \eta & \searrow m \\
 A & \xrightarrow{f} & B
 \end{array}
 \quad
 \begin{array}{ccc}
 & C' & \\
 i' \nearrow & \Downarrow \eta' & \searrow m' \\
 A & \xrightarrow{f} & B,
 \end{array}$$

m, m' are faithful and i, i' are fully cofaithful, then there exist $t \in \mathbf{S}^1(C, C')$, $\zeta_m \in \mathbf{S}^2(t \circ m', m)$, and $\zeta_i \in \mathbf{S}^2(i \circ t, i')$ such that $(i \circ \zeta_m) \cdot \eta = (\zeta_i \circ m') \cdot \eta'$.

(b2) If both (t, ζ_m, ζ_i) and (t', ζ'_m, ζ'_i) satisfy (b1), then there is a unique 2-cell $\kappa \in \mathbf{S}^2(t, t')$ such that $(i \circ \kappa) \cdot \zeta'_i = \zeta_i$ and $(\kappa \circ m') \cdot \zeta'_m = \zeta_m$.

4.15. **REMARK.** Proposition 4.3 and Proposition 4.4 implies respectively the existence of (2,1)-proper factorization system and (1,2)-proper factorization system in any relatively exact 2-category, in the sense of [2].

In the notation of this section, condition (B2) and (B3) in Definition 3.7 can be written as follows:

- 4.16. **COROLLARY.** *For any $f \in \mathbf{S}^1(A, B)$, we have;*
 (1) *f is faithful iff $i(f) : A \rightarrow \text{Im}(f)$ is an equivalence.*
 (2) *f is cofaithful iff $j(f) : \text{Coim}(f) \rightarrow B$ is an equivalence.*

PROOF. Since (1) is the dual of (2), we show only (2).

In the coimage factorization diagram

$$\begin{array}{ccc}
 & \text{Coim}(f) & \\
 c(k(f)) \nearrow & \Downarrow \mu_f & \searrow j(f) \\
 A & \xrightarrow{f} & B,
 \end{array}$$

since $c(k(f))$ is cofaithful and $j(f)$ is fully faithful, we have f is cofaithful $\iff j(f)$ is cofaithful $\xleftrightarrow{\text{Prop. 3.26}}$ $j(f)$ is an equivalence. ■

5. Definition of relative 2-exactness

DIAGRAM LEMMAS (1).

5.1. **DEFINITION.** *A complex $A_\bullet = (A_n, d_n^A, \delta_n^A)$ is a diagram*

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \uparrow \delta_{n-1}^A & & \uparrow \delta_n^A & & \\
 \cdots & A_{n-2} & \xrightarrow{d_{n-2}^A} & A_{n-1} & \xrightarrow{d_{n-1}^A} & A_n & \xrightarrow{d_n^A} & A_{n+1} & \xrightarrow{d_{n+1}^A} & A_{n+2} & \cdots \\
 & & \downarrow d_{n-2}^A & & \downarrow d_{n-1}^A & & \downarrow d_n^A & & \downarrow d_{n+1}^A & & \\
 & & 0 & & 0 & & 0 & & & &
 \end{array}$$

where $A_n \in \mathbf{S}^0$, $d_n^A \in \mathbf{S}^1(A_n, A_{n+1})$, $\delta_n^A \in \mathbf{S}^2(d_{n-1}^A \circ d_n^A, 0)$, and satisfies the following compatibility condition for each $n \in \mathbb{Z}$:

$$(d_{n-1}^A \circ \delta_{n+1}^A) \cdot (d_{n-1}^A)_I^\sharp = (\delta_n^A \circ d_{n+1}^A) \circ (d_{n+1}^A)_I^\flat$$

5.2. **REMARK.** We consider a bounded complex as a particular case of a complex, by adding zeroes.

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \uparrow (d_0^A)_I^\flat & & \uparrow \delta_2^A & & \\
 \cdots & 0 & \xrightarrow{0} & 0 & \xrightarrow{d_0^A} & A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \cdots \\
 & & \downarrow \text{id} & & \downarrow \delta_1^A & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & 0 & & & &
 \end{array}$$

5.3. DEFINITION. For any complexes $A_\bullet = (A_n, d_n^A, \delta_n^A)$ and $B_\bullet = (B_n, d_n^B, \delta_n^B)$, a complex morphism $f_\bullet = (f_n, \lambda_n) : A_\bullet \rightarrow B_\bullet$ consists of $f_n \in \mathbf{S}^1(A_n, B_n)$ and $\lambda_n \in \mathbf{S}^2(d_n^A \circ f_{n+1}, f_n \circ d_n^B)$ for each n , satisfying

$$(\delta_n^A \circ f_{n+1}) \cdot (f_{n+1})_I^\flat = (d_{n-1}^A \circ \lambda_n) \cdot (\lambda_{n-1} \circ d_n^B) \cdot (f_{n-1} \circ \delta_n^B) \cdot (f_{n-1})_I^\sharp.$$

$$\begin{array}{ccccccccccc} \cdots & A_{n-2} & \xrightarrow{d_{n-2}^A} & A_{n-1} & \xrightarrow{d_{n-1}^A} & A_n & \xrightarrow{d_n^A} & A_{n+1} & \xrightarrow{d_{n+1}^A} & A_{n+2} & \cdots \\ & \downarrow f_{n-2} & & \downarrow f_{n-1} & & \downarrow f_n & & \downarrow f_{n+1} & & \downarrow f_{n+2} & \\ & \lambda_{n-2} \Downarrow & & \lambda_{n-1} \Downarrow & & \lambda_n \Downarrow & & \lambda_{n+1} \Downarrow & & \lambda_{n+2} \Downarrow & \\ \cdots & B_{n-2} & \xrightarrow{d_{n-2}^B} & B_{n-1} & \xrightarrow{d_{n-1}^B} & B_n & \xrightarrow{d_n^B} & B_{n+1} & \xrightarrow{d_{n+1}^B} & B_{n+2} & \cdots \end{array}$$

5.4. PROPOSITION. Consider the following diagram in \mathbf{S} .

$$\begin{array}{ccc} A_1 & \xrightarrow{f_1} & B_1 \\ a \downarrow & \lambda \Downarrow & \downarrow b \\ A_2 & \xrightarrow{f_2} & B_2 \end{array} \tag{27}$$

If we take the cokernels of f_1 and f_2 , then there exist $\bar{b} \in \mathbf{S}^1(\text{Cok}(f_1), \text{Cok}(f_2))$ and $\bar{\lambda} \in \mathbf{S}^2(c(f_1) \circ \bar{b}, b \circ c(f_2))$ such that

$$(\pi_{f_1} \circ \bar{b}) \cdot (\bar{b})_I^\flat = (f_1 \circ \bar{\lambda}) \cdot (\lambda \circ c(f_2)) \cdot (a \circ \pi_{f_2}) \cdot a_I^\sharp.$$

$$\begin{array}{ccccc} & & 0 & & \\ & & \uparrow \pi_{f_1} c(f_1) \bar{b} & & \\ A_1 & \xrightarrow{f_1} & B_1 & \xrightarrow{c(f_1)} & \text{Cok}(f_1) \\ a \downarrow & \lambda \Downarrow & \downarrow b & \Downarrow \bar{\lambda} & \downarrow \bar{b} \\ A_2 & \xrightarrow{f_2} & B_2 & \xrightarrow{c(f_2)} & \text{Cok}(f_2) \\ & & \downarrow \pi_{f_2} & & \\ & & 0 & & \end{array}$$

If $(\bar{b}', \bar{\lambda}')$ also satisfies this condition, there exists a unique 2-cell $\xi \in \mathbf{S}^2(\bar{b}, \bar{b}')$ such that $(c(f_1) \circ \xi) \cdot \bar{\lambda}' = \bar{\lambda}$.

PROOF. This follows immediately if we apply the universality of $\text{Cok}(f_1)$ to $(\lambda \circ c(f_2)) \cdot (a \circ \pi_{f_2}) \cdot a_I^\sharp \in \mathbf{S}^2(f_1 \circ b \circ c(f_2), 0)$. ■

5.5. PROPOSITION. Consider the following diagrams in \mathbf{S} ,

$$\begin{array}{ccc} \begin{array}{ccc} A_1 & \xrightarrow{f_1} & B_1 \\ a_1 \downarrow & \lambda_1 \Downarrow & \downarrow b_1 \\ a \Leftarrow A_2 & \xrightarrow{f_2} & B_2 \Rightarrow b \\ a_2 \downarrow & \lambda_2 \Downarrow & \downarrow b_2 \\ A_3 & \xrightarrow{f_3} & B_3 \end{array} & & \begin{array}{ccc} A_1 & \xrightarrow{f_1} & B_1 \\ a \downarrow & \lambda \Downarrow & \downarrow b \\ A_3 & \xrightarrow{f_3} & B_3 \end{array} \end{array}$$

which satisfy $(f_1 \circ \beta) \cdot \lambda = (\lambda_1 \circ b_2) \cdot (a_1 \circ \lambda_2) \cdot (\alpha \circ f_3)$. Applying Proposition 5.4, we obtain diagrams

$$\begin{array}{ccccc} B_1 \xrightarrow{c(f_1)} \text{Cok}(f_1) & B_1 \xrightarrow{c(f_1)} \text{Cok}(f_1) & B_2 \xrightarrow{c(f_2)} \text{Cok}(f_2) & & \\ b \downarrow \quad \Downarrow \bar{\lambda} \quad \downarrow \bar{b} & b_1 \downarrow \quad \Downarrow \bar{\lambda}_1 \quad \downarrow \bar{b}_1 & b_2 \downarrow \quad \Downarrow \bar{\lambda}_2 \quad \downarrow \bar{b}_2 & & \\ B_3 \xrightarrow{c(f_3)} \text{Cok}(f_3) & B_2 \xrightarrow{c(f_2)} \text{Cok}(f_2) & B_3 \xrightarrow{c(f_3)} \text{Cok}(f_3) & & \end{array}$$


with

$$\begin{aligned} (\pi_{f_1} \circ \bar{b}) \cdot (\bar{b})_I^\flat &= (f_1 \circ \bar{\lambda}) \cdot (\lambda \circ c(f_3)) \cdot (a \circ \pi_{f_3}) \cdot a_I^\sharp & (28) \\ (\pi_{f_1} \circ \bar{b}_1) \cdot (\bar{b}_1)_I^\flat &= (f_1 \circ \bar{\lambda}_1) \cdot (\lambda_1 \circ c(f_2)) \cdot (a_1 \circ \pi_{f_2}) \cdot (a_1)_I^\sharp \\ (\pi_{f_2} \circ \bar{b}_2) \cdot (\bar{b}_2)_I^\flat &= (f_2 \circ \bar{\lambda}_2) \cdot (\lambda_2 \circ c(f_3)) \cdot (a_2 \circ \pi_{f_3}) \cdot (a_2)_I^\sharp. \end{aligned}$$

Then, there exists a unique 2-cell $\bar{\beta} \in \mathbf{S}^2(\bar{b}_1 \circ \bar{b}_2, \bar{b})$ such that

$$(c(f_1) \circ \bar{\beta}) \cdot \bar{\lambda} = (\bar{\lambda}_1 \circ \bar{b}_2) \cdot (b_1 \circ \bar{\lambda}_2) \cdot (\beta \circ c(f_3)).$$

PROOF. By (28), $\bar{\lambda}$ is compatible with π_{f_1} and $(\lambda \circ c(f_3)) \cdot (a \circ \pi_{f_3}) \cdot a_I^\sharp$.



On the other hand, $\bar{\lambda}' := (\bar{\lambda}_1 \circ \bar{b}_2) \cdot (b_1 \circ \bar{\lambda}_2) \cdot (\beta \circ c(f_3))$ is also compatible with π_{f_1} and $(\lambda \circ c(f_3)) \cdot (a \circ \pi_{f_3}) \cdot a_I^\sharp$. So, by the universality of the $\text{Cok}(f_1)$, there exists a unique 2-cell $\bar{\beta} \in \mathbf{S}^2(\bar{b}_1 \circ \bar{b}_2, \bar{b})$ such that $(c(f_1) \circ \bar{\beta}) \cdot \bar{\lambda} = \bar{\lambda}'$. ■

5.6. COROLLARY. Let $(f_n, \lambda_n) : (A_n, d_n^A, \delta_n^A) \longrightarrow (B_n, d_n^B, \delta_n^B)$ be a complex morphism. Then, by taking the cokernels, we obtain a complex morphism $(c(f_n), \bar{\lambda}_n) : (B_n, d_n^B, \delta_n^B) \longrightarrow (\text{Cok}(f_n), \bar{d}_n^B, \bar{\delta}_n^B)$ which satisfies

$$(d_n^A \circ \pi_{f_{n+1}}) \cdot (d_n^A)_I^\sharp = (\lambda_n \circ c(f_{n+1})) \cdot (f_n \circ \bar{\lambda}_n) \cdot (\pi_{f_n} \circ \bar{d}_n^B) \cdot (\bar{d}_n^B)_I^\flat \tag{29}$$

for each n .

PROOF. By Proposition 5.4, we obtain \bar{d}_n^B and $\bar{\lambda}_n$ which satisfy (29). And by Proposition 5.5, for each n , there exists a unique 2-cell $\bar{\delta}_n^B \in \mathbf{S}^2(\bar{d}_{n-1}^B \circ \bar{d}_n^B, 0)$ such that $((\bar{\delta}_n^B \circ c(f_{n+1})) \cdot c(f_{n+1})_I^\flat = (d_{n-1}^B \circ \bar{\lambda}_n) \cdot (\bar{\lambda}_{n-1} \circ \bar{d}_n^B) \cdot (c(f_{n-1}) \circ \bar{\delta}_n^B) \cdot c(f_{n-1})_I^\sharp$. By the uniqueness of $\bar{\beta}$ in Proposition 5.5, it is easy to see that

$$(\bar{\delta}_n^B \circ \bar{d}_{n+1}^B) \cdot (\bar{d}_{n+1}^B)_I^\flat = (\bar{d}_{n-1}^B \circ \bar{\delta}_{n+1}^B) \cdot (\bar{d}_{n-1}^B)_I^\sharp.$$

These are saying that $(\text{Cok}(f_n), \bar{d}_n^B, \bar{\delta}_n^B)$ is a complex and $(c(f_n), \bar{\lambda}_n)$ is a complex morphism. ■

5.7. PROPOSITION. Consider the following diagram in \mathbf{S} .

$$\begin{array}{ccc} A_1 & \xrightarrow{f_1} & B_1 \\ a \downarrow & \lambda \Downarrow & \downarrow b \\ A_3 & \xrightarrow{f_3} & B_3 \end{array}$$

By taking the cokernels of f_1 and f_2 , we obtain

$$\begin{array}{ccccc} A_1 & \xrightarrow{f_1} & B_1 & \xrightarrow{c(f_1)} & \text{Cok}(f_1) \\ a \downarrow & \lambda \Downarrow & \downarrow b & \Downarrow \lambda' & \downarrow \bar{b} \\ A_2 & \xrightarrow{f_2} & B_2 & \xrightarrow{c(f_2)} & \text{Cok}(f_2) \end{array}$$

and from this diagram, by taking the cokernels of a, b, \bar{b} , we obtain

$$\begin{array}{ccccc} & & 0 & & \\ & & \uparrow \pi_{f_2} & & \\ A_2 & \xrightarrow{f_2} & B_2 & \xrightarrow{c(f_2)} & \text{Cok}(f_2) \\ c(a) \downarrow & \bar{\lambda} \Downarrow & \downarrow c(b) & \Downarrow \bar{\lambda}' & \downarrow c(\bar{b}) \\ \text{Cok}(a) & \xrightarrow{\bar{f}_2} & \text{Cok}(b) & \xrightarrow{c(f_2)} & \text{Cok}(\bar{b}) \\ & & \downarrow \bar{\pi}_{f_2} & & \\ & & 0 & & \end{array}$$

Then we have $\text{Cok}(\bar{f}_2) = [\text{Cok}(\bar{b}), \overline{c(f_2)}, \bar{\pi}_{f_2}]$. We abbreviate this to $\text{Cok}(\bar{f}_2) = \text{Cok}(\bar{b})$.

PROOF. Left to the reader. ■

5.8. PROPOSITION. In the following diagram, assume $f_\bullet : A_\bullet \rightarrow B_\bullet$ is a complex morphism.

$$\begin{array}{ccccc} & & 0 & & \\ & & \uparrow \delta_2^A & & \\ A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 \\ f_1 \downarrow & \lambda_1 \Downarrow & \downarrow f_2 & \lambda_2 \Downarrow & \downarrow f_3 \\ B_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{d_2^B} & B_3 \\ & & \downarrow \delta_2^B & & \\ & & 0 & & \end{array} \tag{30}$$

If we take the cokernels of d_1^A and d_1^B ,

$$\begin{array}{ccccc} A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{c(d_1^A)} & \text{Cok}(d_1^A) \\ f_1 \downarrow & \lambda_1 \Downarrow & \downarrow f_2 & \bar{\lambda}_1 \Downarrow & \downarrow \bar{f}_2 \\ B_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{c(d_1^B)} & \text{Cok}(d_1^B) \end{array}$$

then by the universality of cokernel, we obtain $\bar{d}_2^A \in \mathbf{S}^1(\text{Cok}(d_1^A), A_3)$ and $\bar{\delta}_2^A \in \mathbf{S}^2(c(d_1^A) \circ \bar{d}_2^A, d_2^A)$ such that $(d_1^A \circ \bar{\delta}_2^A) \cdot \delta_2^A = (\pi_{d_1^A} \circ \bar{d}_2^A) \cdot (\bar{d}_2^A)_I^b$. Similarly, we obtain $\bar{d}_2^B \in \mathbf{S}^1(\text{Cok}(d_1^B), B_3)$, $\bar{\delta}_2^B \in \mathbf{S}^2(c(d_1^B) \circ \bar{d}_2^B, d_2^B)$ with $(d_1^B \circ \bar{\delta}_2^B) \cdot \delta_2^B = (\pi_{d_1^B} \circ \bar{d}_2^B) \cdot (\bar{d}_2^B)_I^b$. Then, there exists a unique 2-cell $\bar{\lambda}_2 \in \mathbf{S}^2(\bar{d}_2^A \circ f_3, \bar{f}_2 \circ \bar{d}_2^B)$ such that $(c(d_1^A) \circ \bar{\lambda}_2) \cdot (\bar{\lambda}_1 \circ \bar{d}_2^B) \cdot (f_2 \circ \bar{\delta}_2^B) = (\bar{\delta}_2^A \circ f_3) \cdot \lambda_2$.

PROOF. If we put $\delta := (d_1^A \circ \lambda_2^{-1}) \cdot (\delta_2^A \circ f_3) \cdot (f_3)_I^b$, then both the factorizations

$$\begin{aligned} (\bar{\delta}_2^A \circ f_3) \cdot \lambda_2 & : c(d_1^A) \circ (\bar{d}_2^A \circ f_3) \implies f_2 \circ d_2^B \\ (\bar{\lambda}_1 \circ \bar{d}_2^B) \cdot (f_2 \circ \bar{\delta}_2^B) & : c(d_1^A) \circ (\bar{f}_2 \circ \bar{d}_2^B) \implies f_2 \circ d_2^B \end{aligned}$$

are compatible with $\pi_{d_1^A}$ and δ . So the proposition follows from the universality of $\text{Cok}(d_2^A)$. ■

5.9. PROPOSITION. In diagram (27), if we take the coimage factorization $\mu_a : c(k(a)) \circ j(a) \implies a$ and $\mu_b : c(k(b)) \circ j(b) \implies b$, then there exist $f \in \mathbf{S}^1(\text{Coim}(a), \text{Coim}(b))$, $\lambda_1 \in \mathbf{S}^2(f_1 \circ c(k(b)), c(k(a)) \circ f)$ and $\lambda_2 \in \mathbf{S}^2(f \circ j(b), j(a) \circ f_2)$ such that $(f_1 \circ \mu_b) \cdot \lambda = (\lambda_1 \circ j(b)) \cdot (c(k(a)) \circ \lambda_2) \cdot (\mu_a \circ f_2)$.

$$\begin{array}{ccc} A_1 & \xrightarrow{f_1} & B_1 \\ \downarrow c(k(a)) & \Downarrow \lambda_1 & \downarrow c(k(b)) \\ a & \xrightarrow{\mu_a} \text{Coim}(a) \xrightarrow{f} \text{Coim}(b) \xrightarrow{\mu_b} & b \\ \downarrow j(a) & \Downarrow \lambda_2 & \downarrow j(b) \\ A_2 & \xrightarrow{f_2} & B_2 \end{array} \tag{31}$$

Moreover, for any other $(f', \lambda'_1, \lambda'_2)$ with this property, there exists a unique 2-cell $\xi \in \mathbf{S}^2(f, f')$ such that $\lambda_1 \cdot (c(k(a)) \circ \xi) = \lambda'_1$ and $(\xi \circ j(b)) \cdot \lambda'_2 = \lambda_2$.

PROOF. Since the coimage factorization is unique up to an equivalence and is obtained by the factorization which fills in the following diagram, we may assume $\text{Ker}(a) = [\text{Ker}(a), k(a), \varepsilon_a]$, $\text{Cok}(k(a)) = [\text{Coim}(a), c(k(a)), \pi_{k(a)}]$, and $(k(a) \circ \mu_a) \cdot \varepsilon_a = (\pi_{k(a)} \circ j(a)) \cdot j(a)_I^b$.

$$\begin{array}{ccc} & \xrightarrow{0} & \text{Coim}(a) \\ & \swarrow \pi_{k(a)} & \nearrow c(k(a)) \\ \text{Ker}(a) & \xrightarrow{k(a)} & A_1 \\ & \searrow \varepsilon_a & \downarrow a \\ & \xrightarrow{0} & A_2 \end{array}$$

$\exists j(a)$

Similarly, we may assume

$$\begin{aligned} \text{Ker}(b) & = [\text{Ker}(b), k(b), \varepsilon_b], \\ \text{Cok}(k(b)) & = [\text{Coim}(b), c(k(b)), \pi_{k(b)}] \end{aligned}$$

and $(k(b) \circ \mu_b) \cdot \varepsilon_b = (\pi_{k(b)} \circ j(b)) \cdot j(b)_I^\flat$. By (the dual of) Proposition 5.4, there are $\underline{f}_1 \in \mathbf{S}^1(\text{Ker}(a), \text{Ker}(b))$ and $\underline{\lambda} \in \mathbf{S}^2(\underline{f}_1 \circ k(b), k(a) \circ \underline{f}_1)$ such that $(\underline{\lambda} \circ b) \cdot (k(a) \circ \underline{\lambda}) \cdot (\varepsilon_a \circ f_2) \cdot (f_2)_I^\flat = (\underline{f}_1 \circ \varepsilon_b) \cdot (\underline{f}_1)_I^\sharp$. Applying Proposition 5.8, we can show the existence of $(f, \lambda_1, \lambda_2)$. To show the uniqueness (up to an equivalence), let $(f', \lambda'_1, \lambda'_2)$ satisfy

$$(f_1 \circ \mu_b) \cdot \lambda = (\lambda'_1 \circ j(b)) \cdot (c(k(a)) \circ \lambda'_2) \cdot (\mu_a \circ f_2).$$

From this, we can obtain

$$(\underline{f}_1 \circ \pi_{k(b)}) \cdot (\underline{f}_1)_I^\sharp = (\underline{\lambda} \circ c(k(b))) \cdot (k(a) \circ \lambda'_1) \cdot (\pi_{k(a)} \circ f') \cdot f'_I{}^\flat.$$

And the uniqueness follows from the uniqueness of 2-cells in Proposition 5.4 and Proposition 5.8. ■

5.10. PROPOSITION. *Let $f_\bullet : A_\bullet \rightarrow B_\bullet$ be a complex morphism as in diagram (30). If we take the cokernels of f_1, f_2, f_3 and relative cokernels of the complex A_\bullet and B_\bullet as in the following diagram, then we have $\text{Cok}(\bar{f}_3) = \text{Cok}(\bar{d}_2^B, \bar{\delta}_2^B)$.*

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & & \uparrow \delta_2^A & & & & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{c(d_2^A, \delta_2^A)} & \text{Cok}(d_2^A, \delta_2^A) \\
 f_1 \downarrow & \lambda_1 \Downarrow & f_2 \downarrow & \lambda_2 \Downarrow & f_3 \downarrow & \exists \Downarrow & \bar{f}_3 \downarrow \\
 B_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{d_2^B} & B_3 & \xrightarrow{c(d_2^B, \delta_2^B)} & \text{Cok}(d_2^B, \delta_2^B) \\
 c(f_1) \downarrow & \bar{\lambda}_1 \Downarrow & c(f_2) \downarrow & \bar{\lambda}_2 \Downarrow & c(f_3) \downarrow & & \\
 \text{Cok}(f_1) & \xrightarrow{\bar{d}_1^B} & \text{Cok}(f_2) & \xrightarrow{\bar{d}_2^B} & \text{Cok}(f_3) & & \\
 & & \bar{\delta}_2^B \Downarrow & & & & \\
 & & 0 & & & &
 \end{array}$$

PROOF. Immediately follows from Proposition 5.7, Proposition 5.8 and (the dual of) Proposition 3.20. ■

5.11. PROPOSITION. *In diagram (27), if a is fully cofaithful, then the following diagram obtained in Proposition 5.4 is a pushout diagram.*

$$\begin{array}{ccc}
 B_1 & \xrightarrow{c(f_1)} & \text{Cok}(f_1) \\
 b \downarrow & \Downarrow \bar{\lambda} & \downarrow \bar{b} \\
 B_2 & \xrightarrow{c(f_2)} & \text{Cok}(f_2)
 \end{array}$$

PROOF. Left to the reader. ■

Concerning Proposition 3.32, we have the following proposition.

5.12. PROPOSITION. *Let*

$$\begin{array}{ccc} A_1 \times_B A_2 & \xrightarrow{f'_1} & A_2 \\ f'_2 \downarrow & \xi \Downarrow & \downarrow f_2 \\ A_1 & \xrightarrow{f_1} & B \end{array}$$

be a pullback diagram in \mathbf{S} . If f_1 is fully cofaithful, then f'_1 is fully cofaithful.

PROOF. Since f_1 is cofaithful, in the notation of the proof of Proposition 3.32, $\text{Cok}(i_1) = [A_2, p_2, \xi_2]$ and $\text{Cok}(d) = [B, t, \varepsilon_t]$. Applying Proposition 5.7 to the diagram

$$\begin{array}{ccc} 0 & \xrightarrow{0} & A_1 \\ 0 \downarrow & & \downarrow \\ A_1 \times_B A_2 & \xrightarrow{d} & A_1 \times A_2 \end{array}$$

we obtain

$$\text{Cok}(f_1) = 0 \iff \text{Cok}(f'_1) = 0.$$

$$\begin{array}{ccccccc} 0 & \xrightarrow{0} & A_1 & \xlongequal{\text{id}} & A_1 & & \\ 0 \downarrow & & \downarrow & \downarrow i_1 & \downarrow & & \downarrow f_1 \\ A_1 \times_B A_2 & \xrightarrow{d} & A_1 \times A_2 & \xrightarrow{t} & B & & \\ \text{id} \parallel & & \circlearrowleft & \downarrow p_2 & \downarrow & & \downarrow 0 \\ A_1 \times_B A_2 & \xrightarrow{f'_1} & A_2 & \xrightarrow{0} & 0 & & \end{array}$$

■

5.13. PROPOSITION. *In diagram (27), assume a is cofaithful. By Proposition 5.9, we obtain a coimage factorization diagram as (31). If we take the cokernel of this diagram as*

$$\begin{array}{ccc} B_1 & \xrightarrow{c(f_1)} & \text{Cok}(f_1) \\ \downarrow c(k(b)) & \Downarrow \bar{\lambda}_1 & \downarrow c(k(b)) \\ b & \xleftarrow{\mu_b} \text{Coim}(b) \xrightarrow{c(f)} \text{Cok}(f) \xrightarrow{\bar{\mu}_b} & \bar{b} \\ \downarrow j(b) & \Downarrow \bar{\lambda}_2 & \downarrow j(b) \\ B_2 & \xrightarrow{c(f_2)} & \text{Cok}(f_2) \end{array}$$

then the factorization

$$\begin{array}{ccc} & \text{Cok}(f) & \\ \nearrow c(k(b)) & \downarrow \bar{\mu}_b & \searrow j(b) \\ \text{Cok}(f_1) & \xrightarrow{\bar{b}} & \text{Cok}(f_2) \end{array}$$

becomes again a coimage factorization.

PROOF. It suffices to show that $\overline{c(k(b))}$ is cofaithful and $\overline{j(b)}$ is fully faithful. Since $c(k(b))$ and $c(f)$ are cofaithful, it follows that $\overline{c(k(b))}$ is cofaithful. Since $j(a)$ is an equivalence,

$$\begin{array}{ccc} \text{Coim}(b) & \xrightarrow{c(f)} & \text{Cok}(f) \\ j(b)\downarrow & \Downarrow \lambda_2 & \downarrow \overline{j(b)} \\ B_2 & \xrightarrow{c(f_2)} & \text{Cok}(f_2) \end{array}$$

is a pushout diagram by Proposition 5.11. By (the dual of) Proposition 5.12, $\overline{j(b)}$ becomes fully faithful. ■

DEFINITION OF THE RELATIVE 2-EXACTNESS.

5.14. LEMMA. Consider the following diagram in \mathbf{S} .

$$\begin{array}{ccccc} A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ & \searrow & \varphi \Downarrow & \nearrow & \\ & & 0 & & \end{array} \tag{32}$$

If we factor it as

$$\begin{array}{ccccc} & \text{Ker}(g) & & \text{Cok}(f) & \\ & \nearrow \underline{f} & \searrow k(g) & \nearrow c(f) & \searrow \overline{g} \\ A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ & \searrow & \varphi \Downarrow & \nearrow & \\ & & 0 & & \end{array} \tag{33}$$

with

$$\begin{aligned} (\underline{\varphi} \circ g) \cdot \varphi &= (\underline{f} \circ \varepsilon_g) \cdot (\underline{f})_I^\sharp \\ (f \circ \overline{\varphi}) \cdot \varphi &= (\pi_f \circ \overline{g}) \cdot (\overline{g})_I^b, \end{aligned}$$

then $\text{Cok}(\underline{f}) = 0$ if and only if $\text{Ker}(\overline{g}) = 0$.

PROOF. We show only $\text{Cok}(\underline{f}) = 0 \Rightarrow \text{Ker}(\overline{g}) = 0$, since the other implication can be shown dually. If $\text{Cok}(\underline{f}) = 0$, i.e. if \underline{f} is fully cofaithful, then we have

$$\text{Cok}(f) = \text{Cok}(\underline{f} \circ k(g)) = \text{Cok}(k(g)) = \text{Coim}(g).$$

Thus the following diagram is a coimage factorization, and \overline{g} becomes fully faithful.

$$\begin{array}{ccc} & \text{Cok}(f) & \\ c(f) \nearrow & & \searrow \overline{g} \\ B & \xrightarrow{g} & C \\ & \Downarrow \overline{\varphi} & \end{array}$$

■

5.15. DEFINITION. Diagram (32) is said to be 2-exact in B , if $\text{Cok}(f) = 0$ (or equivalently $\text{Ker}(\bar{g}) = 0$).

5.16. REMARK. In the notation of Lemma 5.14, the following are equivalent :

- (i) (32) is 2-exact in B .
- (ii) f is fully cofaithful.
- (iii) \bar{g} is fully faithful.
- (iv) $c(f) = \text{cok}(k(g))$ (i.e. $\text{Cok}(f) = \text{Coim}(g)$).
- (v) $k(g) = \text{ker}(c(f))$ (i.e. $\text{Ker}(g) = \text{Im}(f)$).

PROOF. By the duality, we only show (i) \Leftrightarrow (iii) \Leftrightarrow (v).

- (i) \Leftrightarrow (iii) follows from Corollary 3.25.
- (iii) \Rightarrow (v) follows from Proposition 3.21.
- (v) \Rightarrow (iii) follows from Proposition 4.3. ■


Let us fix the notation for relative (co-)kernels of a complex.

5.17. DEFINITION. For any complex $A_\bullet = (A_n, d_n, \delta_n)$ in \mathbf{S} , we put


- (1) $[Z^n(A_\bullet), z_n^A, \zeta_n^A] := \text{Ker}(d_n, \delta_{n+1})$.
- (2) $[Q^n(A_\bullet), q_n^A, \rho_n^A] := \text{Cok}(d_{n-1}, \delta_{n-1})$.

5.18. REMARK. By the universality of $\text{Ker}(d_n, \delta_{n+1})$ and Lemma 3.19, there exist $k_n \in \mathbf{S}^1(A_{n-1}, Z^n(A_\bullet))$, $\nu_{n,1} \in \mathbf{S}^2(k_n \circ z_n, d_{n-1})$ and $\nu_{n,2} \in \mathbf{S}^2(d_{n-2} \circ k_n, 0)$ such that

$$\begin{aligned} (\nu_{n,1} \circ d_n) \cdot \delta_n &= (k_n \circ \zeta_n) \cdot (k_n)_I^\sharp \\ (d_{n-2} \circ \nu_{n,1}) \cdot \delta_{n-1} &= (\nu_{n,2} \circ z_n) \cdot (z_n)_I^\flat \end{aligned}$$



On the other hand, by the universality of $\text{Ker}(d_n)$, we obtain a factorization diagram



which satisfy

$$\begin{aligned} (\underline{\delta}_n \circ d_n) \cdot \delta_n &= (\underline{d}_{n-1} \circ \varepsilon_{d_n}) \cdot (\underline{d}_{n-1})_I^\sharp \\ (d_{n-2} \circ \underline{\delta}_n) \cdot \delta_{n-1} &= (\underline{\delta}_{n-1} \circ k(d_n)) \cdot (k(d_n))_I^\flat \end{aligned}$$

By Proposition 3.20, there exists a factorization of z_n through $\text{Ker}(d_n)$

$$\begin{array}{ccccc}
 Z^n(A_\bullet) & \xrightarrow{\quad 0 \quad} & & & \\
 \downarrow \underline{z}_n & \nearrow z_n & \zeta_n \nearrow & & \\
 \text{Ker}(d_n) & \xrightarrow{k(d_n)} & A_n & \xrightarrow{d_n} & A_{n+1} \\
 & & \searrow \varepsilon_{d_n} & & \\
 & & & \xrightarrow{\quad 0 \quad} &
 \end{array}$$

which satisfies

$$(\underline{\zeta}_n \circ d_n) \cdot \zeta_n = (\underline{z}_n \circ \varepsilon_{d_n}) \cdot (\underline{z}_n)_I^\sharp$$

Moreover \underline{z}_n is fully faithful by Proposition 3.33.

By the universality of $\text{Ker}(d_n)$, we can show easily the following claim.

5.19. CLAIM. *There exists a unique 2-cell $\widehat{\zeta}_n \in S(k_n \circ z_n, \underline{d}_{n-1})$*

$$\begin{array}{ccccc}
 & & Z^n(A_\bullet) & & \\
 & \nearrow 0 & \downarrow z_n & \searrow 0 & \\
 A_{n-2} & \xrightarrow{d_{n-2}} & A_{n-1} & \xrightarrow{d_n} & A_{n+1} \\
 \downarrow \underline{\delta}_{n-1} & \nearrow \widehat{\zeta}_n & \downarrow \underline{z}_n & \nearrow \underline{\zeta}_n & \downarrow \varepsilon_{d_n} \\
 & & \text{Ker}(d_n) & & \\
 & \searrow 0 & \nearrow k(d_n) & \nearrow 0 &
 \end{array}$$

such that


$$(\widehat{\zeta}_n \circ k(d_n)) \cdot \underline{\delta}_n = (k_n \circ \underline{\zeta}_n) \cdot \nu_{n,1}$$

This $\widehat{\zeta}_n$ also satisfies

$$(d_{n-2} \circ \widehat{\zeta}_n) \cdot \underline{\delta}_{n-1} = (\nu_{n,2} \circ \underline{z}_n) \cdot (\underline{z}_n)_I^\flat$$

5.20. REMARK. Dually, by the universality of the cokernels, we obtain the following two factorization diagrams, where \bar{q}_n is fully cofaithful.

$$\begin{array}{ccccccc}
 & & Q^n(A_\bullet) & & & & \\
 & \nearrow 0 & \downarrow q_n & \searrow \ell_n & \nearrow 0 & & \\
 A_{n-2} & \xrightarrow{d_{n-2}} & A_{n-1} & \xrightarrow{d_n} & A_{n+1} & \xrightarrow{d_{n+1}} & A_{n+2} \\
 \downarrow \pi_{d_{n-1}} & \nearrow \rho_n & \downarrow \mu_{n,1} & \nearrow \mu_{n,2} & \downarrow \bar{\delta}_{n+1} & & \\
 & & \text{Cok}(d_{n-1}) & & & & \\
 & \searrow 0 & \nearrow c(d_{n-1}) & \nearrow \bar{d}_n & \nearrow 0 & &
 \end{array}$$



We define relative 2-cohomology in the following two ways, which will be shown to be equivalent later.


5.21. DEFINITION.

$$\begin{aligned}
 H_1^n(A_\bullet) &:= \text{Cok}(k_n, \nu_{n,2}) \\
 H_2^n(A_\bullet) &:= \text{Ker}(\ell_n, \mu_{n,2})
 \end{aligned}$$

5.22. LEMMA. *In the factorization diagram (33) in Lemma 5.14, if we take the cokernel of \underline{f} and the kernel of \bar{g} , then there exist $w \in \mathbf{S}^1(\text{Cok}(\underline{f}), \text{Ker}(\bar{g}))$ and $\omega \in \mathbf{S}^2(c(\underline{f}) \circ w \circ k(\bar{g}), k(\underline{g}) \circ c(\underline{f}))$ such that*

$$(\underline{f} \circ \omega) \cdot (\varphi \circ c(\underline{f})) \cdot \pi_f = (\pi_{\underline{f}} \circ w \circ k(\bar{g})) \cdot (w \circ k(\bar{g}))_I^\flat \tag{34}$$

$$(\omega \circ \bar{g}) \cdot (k(\underline{g}) \circ \bar{\varphi}) \cdot \varepsilon_g = (c(\underline{f}) \circ w \circ \varepsilon_{\bar{g}}) \cdot (c(\underline{f}) \circ w)_I^\sharp. \tag{35}$$



Moreover, for any other factorization (w', ω') with these properties, there exists a unique 2-cell $\kappa \in \mathbf{S}^2(w, w')$ such that $(c(\underline{f}) \circ \kappa \circ k(\bar{g})) \cdot \omega' = \omega$.

PROOF. Applying Proposition 5.4 to

$$\begin{array}{ccc}
 A & \xrightarrow{\text{id}_A} & A \\
 \underline{f} \downarrow \bar{\varphi}^{-1} \Downarrow & & \downarrow f \\
 \text{Ker}(g) & \xrightarrow[k(g)]{} & B,
 \end{array} \tag{36}$$

we obtain $w_1 \in \mathbf{S}^1(\text{Cok}(\underline{f}), \text{Cok}(f))$ and $\omega_1 \in \mathbf{S}^2(c(\underline{f}) \circ w_1, k(g) \circ c(f))$ which satisfy

$$(\underline{f} \circ w_1) \cdot (\underline{\varphi} \circ c(f)) \cdot \pi_f = (\pi_f \circ w_1) \cdot (w_1)_I^b. \tag{37}$$

Then $(\omega_1 \circ \bar{g}) \cdot (k(g) \circ \bar{\varphi}) \cdot \varepsilon_g \in \mathbf{S}^2(c(\underline{f}) \circ w_1 \circ \bar{g}, 0)$ becomes compatible with π_f .

$$\begin{array}{ccccc} & & 0 & & \\ & \curvearrowright & \uparrow \pi_f & \curvearrowleft & \\ A & \xrightarrow{\underline{f}} & \text{Ker}(g) & \xrightarrow{c(f)} & \text{Cok}(f)^{w_1 \circ \bar{g}} & \rightarrow C \\ & & \downarrow & & \downarrow & \\ & & 0 & & (\omega_1 \circ \bar{g}) \cdot (k(g) \circ \bar{\varphi}) \cdot \varepsilon_g & \end{array}$$

By Lemma 3.19, there exists a 2-cell $\delta \in S(w_1 \circ \bar{g}, 0)$ such that

$$(c(\underline{f}) \circ \delta) \cdot c(\underline{f})_I^\sharp = (\omega_1 \circ \bar{g}) \cdot (k(g) \circ \bar{\varphi}) \cdot \varepsilon_g.$$

So, if we take the cokernels of $k(g)$ and w_1 , then by Proposition 5.8, we obtain the following diagram:

$$\begin{array}{ccccccc} & & & & g & & \\ & & & & \uparrow \mu_g & & \\ \text{Ker}(g) & \xrightarrow{k(g)} & B & \xrightarrow{c(k(g))} & \text{Coim}(g) & \xrightarrow{j(g)} & C \\ c(f) \downarrow & \xRightarrow{\omega_1} & \downarrow c(f) & \xRightarrow{\exists \bar{\varphi}_1} & \downarrow \exists \bar{c} & \xRightarrow{\exists \bar{\varphi}_2} & \parallel \text{id}_C \\ \text{Cok}(\underline{f}) & \xrightarrow{w_1} & \text{Cok}(f) & \xrightarrow{c(w_1)} & \text{Cok}(w_1) & \xrightarrow{\exists g^\dagger} & C \\ & & & & \downarrow \exists & & \\ & & & & \bar{g} & & \end{array}$$

Applying Proposition 5.7 to (36), we obtain

$$[\text{Cok}(w_1), \bar{c}, (\bar{c})_I^b] = \text{Cok}(0 \xrightarrow{0} \text{Coim}(g)).$$

Thus \bar{c} is an equivalence. Since $j(g)$ is fully faithful, g^\dagger becomes fully faithful. Thus the following diagram is 2-exact in $\text{Cok}(f)$.

$$\begin{array}{ccc} \text{Cok}(\underline{f}) & \xrightarrow{w_1} & \text{Cok}(f) \xrightarrow{\bar{g}} C \\ & \searrow \delta \downarrow & \nearrow \\ & & 0 \end{array} \tag{38}$$

So if we factor (38) by $w \in \mathbf{S}^1(\text{Cok}(\underline{f}), \text{Ker}(\bar{g}))$ and $\omega_2 \in \mathbf{S}^2(w \circ k(\bar{g}), w_1)$ as in the diagram

$$\begin{array}{ccc} & \text{Ker}(\bar{g}) & \xrightarrow{0} \\ w \nearrow & \downarrow \omega_2 & \searrow k(\bar{g}) \\ \text{Cok}(\underline{f}) & \xrightarrow{w_1} & \text{Cok}(f) \xrightarrow{\bar{g}} C \\ & \searrow \delta \downarrow & \nearrow \varepsilon_{\bar{g}} \\ & & 0 \end{array} \tag{39}$$

which satisfies

$$(w \circ \varepsilon_{\bar{g}}) \cdot w_I^\sharp = (\omega_2 \circ \bar{g}) \cdot \delta,$$

then w becomes fully cofaithful by Lemma 5.14. If we put $\omega := (c(\underline{f}) \circ \omega_2) \cdot \omega_1$, then (w, ω) satisfies conditions (34) and (35).

If (w', ω') satisfies

$$\begin{aligned} (\underline{f} \circ \omega') \cdot (\underline{\varphi} \circ c(\underline{f})) \cdot \pi_f &= (\pi_{\underline{f}} \circ w' \circ k(\bar{g})) \cdot (w' \circ k(\bar{g}))_I^\flat & (40) \\ (\omega' \circ \bar{g}) \cdot (k(g) \circ \bar{\varphi}) \cdot \varepsilon_g &= (c(\underline{f}) \circ w' \circ \varepsilon_{\bar{g}}) \cdot (c(\underline{f}) \circ w')_I^\sharp, \end{aligned}$$

then, since both the factorization of $k(g) \circ c(f)$ through $\text{Cok}(\underline{f})$

$$\begin{aligned} \omega' &: c(\underline{f}) \circ w' \circ k(\bar{g}) \implies k(g) \circ c(f) \\ \omega_1 &: c(\underline{f}) \circ w_1 \implies k(g) \circ c(f) \end{aligned}$$

are compatible with $\pi_{\underline{f}}$ and $(\underline{\varphi} \circ c(\underline{f})) \cdot \pi_f$ by (37) and (40), there exists $\omega'_2 \in \mathbf{S}^2(w' \circ k(\bar{g}), w_1)$ such that

$$(c(\underline{f}) \circ \omega'_2) \cdot \omega_1 = \omega'.$$

Then we can see ω'_2 is compatible with $\varepsilon_{\bar{g}}$ and δ . So, comparing this with the factorization (39), by the universality of $\text{Ker}(\bar{g})$, we see there exists a unique 2-cell $\kappa \in \mathbf{S}^2(w, w')$ such that $(\kappa \circ k(\bar{g})) \cdot \omega'_2 = \omega_2$. Then κ satisfies $(c(\underline{f}) \circ \kappa \circ k(\bar{g})) \cdot \omega' = \omega$. Uniqueness of such κ follows from the fact that $c(\underline{f})$ is cofaithful and $k(\bar{g})$ is faithful. ■


5.23. PROPOSITION. *In Lemma 5.22, w is an equivalence.*

PROOF. We showed Lemma 5.22 by taking the cokernel first and the kernel second, but we obtain the same (w, ω) if we take the kernel first and the cokernel second, because of the symmetricity of the statement (and the uniqueness of (w, ω) up to an equivalence) of Lemma 5.22. As shown in the proof, since (38) is 2-exact in $\text{Cok}(\underline{f})$, w becomes fully cofaithful in the factorization (39). By the above remark, similarly w can be obtained also by the factorization

$$\begin{array}{ccccc} & & \text{Cok}(\underline{f}) & & \\ & & \nearrow & \Downarrow & \searrow w \\ A & \longrightarrow & \text{Ker}(g) & \longrightarrow & \text{Ker}(\bar{g}) \\ & \searrow & \Downarrow & \nearrow & \\ & & 0 & & \end{array}$$

where the bottom row is 2-exact in $\text{Ker}(g)$. So w becomes fully faithful. Thus, w is fully cofaithful and fully faithful, i.e. an equivalence. ■


5.24. COROLLARY. For any complex $A_\bullet = (A_n, d_n, \delta_n)$, if we factor it as



(in the notation of Definition 5.17, Remark 5.18 and Remark 5.20), then there exist $w \in \mathbf{S}^1(H_1^n(A_\bullet), H_2^n(A_\bullet))$ and $\omega \in \mathbf{S}^2(c(k_n, \nu_{n,2}) \circ w \circ k(\ell_n, \mu_{n,2}), z_n \circ q_n)$ such that


$$(k_n \circ \omega) \cdot (\nu_{n,1} \circ q_n) \cdot \rho_n = (\pi(k_n, \nu_{n,2}) \circ w \circ k(\ell_n, \mu_{n,2})) \cdot (w \circ k(\ell_n, \mu_{n,2}))_I^b$$

$$(\omega \circ \ell_n) \cdot (z_n \circ \mu_{n,1}) \cdot \zeta_n = (c(k_n, \nu_{n,2}) \circ w \circ \varepsilon(\ell_n, \mu_{n,2})) \cdot (c(k_n, \nu_{n,2}) \circ w)_I^\#.$$



For any other factorization (w', ω') with these conditions, there exists a unique 2-cell $\kappa \in \mathbf{S}^2(\omega, \omega')$ such that $(c(k_n, \nu_{n,2}) \circ \kappa \circ k(\ell_n, \mu_{n,2})) \cdot \omega' = \omega$. Moreover, this w becomes an equivalence.

PROOF. For the factorization diagrams



which satisfy

$$\begin{aligned}
 (d_{n-2} \circ \bar{\delta}_{n-1}) \cdot \delta_{n-1} &= (\pi_{d_{n-2}} \circ \bar{d}_{n-1}) \cdot (\bar{d}_{n-1})_I^\flat \\
 (\bar{\delta}_{n-1} \circ d_n) \cdot \delta_n &= (c(d_{n-2}) \circ \bar{\delta}_n) \cdot c(d_{n-2})_I^\sharp \\
 (\underline{\delta}_{n+1} \circ d_{n+1}) \cdot \delta_{n+1} &= (\underline{d}_n \circ \varepsilon_{d_{n+1}}) \cdot (\underline{d}_n)_I^\sharp \\
 (d_{n-1} \circ \underline{\delta}_{n+1}) \cdot \delta_n &= (\underline{\delta}_n \circ k(d_{n+1})) \cdot k(d_{n+1})_I^\flat,
 \end{aligned}$$

there exists a unique 2-cell $\delta_n^\dagger \in \mathbf{S}^2(\bar{d}_{n-1} \circ \underline{d}_n, 0)$ such that

$$\begin{aligned}
 (\delta_n^\dagger \circ k(d_{n+1})) \cdot k(d_{n+1})_I^\flat &= (\bar{d}_{n-1} \circ \underline{\delta}_{n+1}) \cdot \bar{\delta}_n \\
 (c(d_{n-2}) \circ \delta_n^\dagger) \cdot c(d_{n-2})_I^\sharp &= (\bar{\delta}_{n-1} \circ \underline{d}_n) \cdot \underline{\delta}_n.
 \end{aligned}$$

By Proposition 3.20, applying Lemma 5.22 and Proposition 5.23 to the following diagram, we can obtain Corollary 5.24.

$$\begin{array}{ccccc}
 & Z^n(A_\bullet) & & Q^n(A_\bullet) & \\
 & \nearrow & & \nearrow & \\
 \text{Cok}(d_{n-2}) & \xrightarrow{\bar{d}_{n-1}} & A_n & \xrightarrow{d_n} & \text{Ker}(d_{n+1}) \\
 & \searrow & \delta_n^\dagger \downarrow & \searrow & \\
 & & 0 & &
 \end{array}$$

■

Thus $H_1^n(A_\bullet)$ and $H_2^n(A_\bullet)$ are equivalent. We abbreviate this to $H^n(A_\bullet)$.

5.25. DEFINITION. A complex A_\bullet is said to be relatively 2-exact in A_n if $H^n(A_\bullet)$ is equivalent to zero.

5.26. REMARK. If the complex is bounded, we consider the relative 2-exactness after adding zeroes as in Remark 5.2. For example, a bounded complex

$$\begin{array}{ccccc}
 & & 0 & & \\
 & & \uparrow \varphi & & \\
 A & \xrightarrow{f} & B & \xrightarrow{g} & C \\
 & & & &
 \end{array}$$

is relatively 2-exact in B if and only if

$$\begin{array}{ccccccc}
 & & & 0 & & & \\
 & & & \uparrow \varphi & & & \\
 0 & \xrightarrow{0} & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \xrightarrow{0} & 0 \\
 & & f_I^\flat \downarrow & & & & g_I^\sharp \downarrow & & \\
 & & 0 & & & & 0 & &
 \end{array}$$

is relatively 2-exact in B , and this is equivalent to the 2-exactness in B by Remark 3.18.

6. Long cohomology sequence in a relatively exact 2-category


DIAGRAM LEMMAS (2).

6.1. LEMMA. Let A_\bullet be a complex in \mathbf{S} , in which $A_5 = 0$ and $d_4 = 0$:

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & \curvearrowright & \uparrow \delta_2^A & & \uparrow (d_3^A)_I^\sharp & \curvearrowright & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{d_3^A} & A_4 \xrightarrow{0} 0 \\
 & & \downarrow \delta_3^A & & & & \\
 & & 0 & & & &
 \end{array} \tag{41}$$

Then, (41) is relatively 2-exact in A_3 and A_4 if and only if $\text{Cok}(d_2, \delta_2) = A_4$, i.e. $[Q^3(A_\bullet), q_3, \rho_3] = [A_4, d_3, \delta_3]$.

PROOF. As in Remark 5.20, we have two factorization diagrams



where \bar{q}_3 is fully cofaithful. We have

$$\begin{aligned}
 (41) \text{ is relatively 2-exact in } A_4 & \Leftrightarrow \text{Cok}(d_3, \delta_3) = 0 \\
 & \Leftrightarrow \text{Cok}(\bar{d}_3) = 0 \Leftrightarrow \text{Cok}(\bar{q}_3 \circ l_3) = 0 \\
 & \Leftrightarrow \text{Cok}(l_3) = 0 \Leftrightarrow l_3 \text{ is fully cofaithful}
 \end{aligned}$$

and

$$\begin{aligned}
 (41) \text{ is relatively 2-exact in } A_3 & \Leftrightarrow \text{Ker}(l_3, (l_3)_I^\sharp) = 0 \\
 & \Leftrightarrow \text{Ker}(l_3) = 0 \Leftrightarrow l_3 \text{ is fully faithful.}
 \end{aligned}$$

Thus, (41) is relatively 2-exact in A_3 and A_4 if and only if l is fully cofaithful and fully faithful, i.e. l is an equivalence. ■

By Remark 3.18, we have the following corollary:

6.2. COROLLARY. Let (A_n, d_n, δ_n) be a bounded complex in \mathbf{S} , as follows:

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & \curvearrowright & \uparrow \delta_2^A & \curvearrowright & 0 & & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{0} & 0 \\
 & & & & \downarrow (d_2^A)_I^\# & & \\
 & & & & 0 & &
 \end{array} \tag{42}$$

Then, (42) is relatively 2-exact in A_2 and A_3 if and only if $\text{Cok}(d_1^A) = [A_3, d_2^A, \delta_2^A]$.

6.3. LEMMA. Let A_\bullet be a complex. As in Definition 5.17, Remark 5.18 and Remark 5.20, take a factorization diagram

$$\begin{array}{ccccccc}
 & & Z^{n+1}(A_\bullet) & & & & \\
 & \nearrow 0 & \uparrow k_{n+1} & \searrow z_{n+1} & \nearrow 0 & & \\
 A_{n-1} & \xrightarrow{d_{n-1}} & A_n & \xrightarrow{d_n} & A_{n+1} & \xrightarrow{d_{n+1}} & A_{n+2} \\
 & \searrow \rho_n & \downarrow \nu_{n+1,2} & \uparrow \nu_{n+1,1} & \downarrow \zeta_{n+1} & \nearrow & \\
 & & Q^n(A_\bullet) & & & & \\
 & \nearrow 0 & \uparrow q_n & \searrow \ell_n & \nearrow 0 & & \\
 & & & & & &
 \end{array}$$

which satisfies

$$\begin{aligned}
 (\nu_{n+1,1} \circ d_{n+1}) \cdot \delta_{n+1} &= (k_{n+1} \circ \zeta_{n+1}) \cdot (k_{n+1})_I^\# \\
 (d_{n-1} \circ \nu_{n+1,1}) \cdot \delta_n &= (\nu_{n+1,2} \circ z_{n+1}) \cdot (z_{n+1})_I^b \\
 (d_{n-1} \circ \mu_{n,1}) \cdot \delta_n &= (\rho_n \circ \ell_n) \cdot (\ell_n)_I^b \\
 (\mu_{n,1} \circ d_{n+1}) \cdot \delta_{n+1} &= (q_n \circ \mu_{n,2}) \cdot (q_n)_I^\#
 \end{aligned}$$

Then, there exist $x_n \in \mathbf{S}^1(Q^n(A_\bullet), Z^{n+1}(A_\bullet))$, $\xi_n \in \mathbf{S}^2(x_n \circ z_{n+1}, \ell_n)$ and $\eta_n \in \mathbf{S}^2(q_n \circ x_n, k_{n+1})$ such that

$$\begin{aligned}
 (\xi_n \circ d_{n+1}) \cdot \mu_{n,2} &= (x_n \circ \zeta_{n+1}) \cdot (x_n)_I^\# \\
 (q_n \circ \xi_n) \cdot \mu_{n,1} &= (\eta_n \circ z_{n+1}) \cdot \nu_{n+1,1} \\
 (d_{n-1} \circ \eta_n) \cdot \nu_{n+1,2} &= (\rho_n \circ x_n) \cdot (x_n)_I^b.
 \end{aligned} \tag{43}$$

Moreover, for any other (x'_n, ξ'_n, η'_n) with these properties, there exists a unique 2-cell $\kappa \in \mathbf{S}^2(x_n, x'_n)$ such that $(\kappa \circ z_{n+1}) \cdot \xi'_n = \xi_n$ and $(q_n \circ \kappa) \cdot \eta'_n = \eta_n$.

PROOF. By the cofaithfulness of q_n , we can show $\mu_{n,2}$ is compatible with δ_{n+2} . By the universality of the relative kernel $Z^{n+1}(A_\bullet)$, there exist $x_n \in \mathbf{S}^1(Q^n(A_\bullet), Z^{n+1}(A_\bullet))$ and $\xi_n \in \mathbf{S}^2(x_n \circ z_{n+1}, \ell_n)$ such that

$$(\xi_n \circ d_{n+1}) \cdot \mu_{n,2} = (x_n \circ \zeta_{n+1}) \cdot (x_n)_I^\#$$

Then, both the factorizations

$$\begin{aligned} \nu_{n+1,1} &: k_{n+1} \circ z_{n+1} \implies d_n \\ (q_n \circ \xi_n) \cdot \mu_{n,1} &: q_n \circ x_n \circ z_{n+1} \implies d_n \end{aligned}$$

are compatible with ζ_{n+1} and δ_{n+1} . Thus by the universality of relative kernel $Z^{n+1}(A_\bullet)$, there exists a unique 2-cell $\eta_n \in \mathbf{S}^2(q_n \circ x_n, k_{n+1})$ such that $(q_n \circ \xi_n) \cdot \mu_{n,1} = (\eta_n \circ z_{n+1}) \cdot \nu_{n+1,1}$. It can be easily seen that η_n also satisfies (43). Uniqueness (up to an equivalence) of (x_n, ξ_n, η_n) follows from the universality of the relative kernel $Z^{n+1}(A_\bullet)$ and the uniqueness of η_n . ■

6.4. LEMMA. Consider the following complex diagram in \mathbf{S} .

$$\begin{array}{ccc} & 0 & \\ & \curvearrowright & \\ A & \xrightarrow{f} & B \xrightarrow{g} C \\ & \uparrow \varphi & \\ & 0 & \end{array} \tag{44}$$

If (44) is 2-exact in B and g is cofaithful, then we have $\text{Cok}(f) = [C, g, \varphi]$.

PROOF. If we factor (44) as

$$\begin{array}{ccc} & 0 & \text{Cok}(f) \\ & \curvearrowright & \downarrow \bar{g} \\ A & \xrightarrow{f} & B \xrightarrow{g} C \\ & \uparrow \pi_f & \swarrow \varphi \\ & 0 & \end{array},$$

then, since (44) is 2-exact in B , \bar{g} becomes fully faithful. On the other hand, since g is cofaithful, \bar{g} is also cofaithful. Thus \bar{g} becomes an equivalence. ■

6.5. LEMMA. Consider the following complex morphism in \mathbf{S} .

$$\begin{array}{ccccccc} & & 0 & & & & \\ & & \curvearrowright & & & & \\ A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{0} & 0 \\ & \parallel \text{id} & \downarrow \lambda & \downarrow f_2 & \downarrow \kappa & \downarrow f_3 & \\ A_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{d_2^B} & B_3 & \xrightarrow{0} & 0 \\ & & \downarrow \delta_2^B & \curvearrowright & & & \\ & & 0 & & & & \end{array}$$

If the complexes are relatively 2-exact in A_2, A_3 and B_2, B_3 respectively, i.e. they satisfy $\text{Cok}(d_1^A) = [A_3, d_2^A, \delta_2^A]$ and $\text{Cok}(d_1^B) = [B_3, d_2^B, \delta_2^B]$ (see Corollary 6.2), then the following diagram obtained by taking the kernel of f_2 becomes 2-exact in A_3 .

$$\begin{array}{ccc} & 0 & \\ & \curvearrowright & \\ \text{Ker}(f_2) & \xrightarrow{k(f_2) \circ d_2^A} & A_3 \xrightarrow{f_3} B_3 \\ & & \downarrow (k(f_2) \circ \kappa) \cdot (\varepsilon_{f_2 \circ d_2^B}) \cdot (d_2^B)_I \end{array} \tag{45}$$

PROOF. By taking the kernels of id_{A_1} and f_2 in the diagram

$$\begin{array}{ccc} A_1 & \xrightarrow{d_1^A} & A_2 \\ \text{id} \parallel & \lambda \Downarrow & \downarrow f_2 \\ A_1 & \xrightarrow{d_1^B} & B_2 \end{array}$$

and taking the cokernels of $0_{0,A_1}$ and $k(f_2)$, we obtain the following diagram by Proposition 5.8, where $\theta = k(f_2)_I^b \cdot (d_1^A)_I^{b-1}$:

$$\begin{array}{ccccccc} 0 & \xrightarrow{0} & A_1 & \xlongequal{\text{id}} & A_1 & \xlongequal{\text{id}} & A_1 \\ 0 \downarrow & \xRightarrow{\theta} & \downarrow d_1^A & \xRightarrow{\lambda_1} & \bar{d}_1^A \downarrow & \xRightarrow{\lambda_2} & \downarrow d_1^B \\ \text{Ker}(f_2) & \xrightarrow{k(f_2)} & A_2 & \xrightarrow{c(k(f_2))} & \text{Coim}(f_2) & \xrightarrow{j(f_2)} & B_2 \\ & & & \searrow \mu_{f_2} \downarrow & & \nearrow f_2 & \\ & & & & & & \end{array} \tag{46}$$


By taking the cokernels of $0_{0,\text{Ker}(f_2)}$, d_1^A and d_1^B in (46), we obtain the left of the following diagrams, while by Proposition 5.13 we obtain the right as a coimage factorization if we take the cokernels of d_1^A , \bar{d}_1^A and d_1^B in (46):

$$\begin{array}{ccc} \begin{array}{ccc} 0 & \xrightarrow{\varepsilon_{f_2} \uparrow} & B_2 \\ \text{Ker}(f_2) \xrightarrow{k(f_2)} A_2 \xrightarrow{f_2} & & \\ \text{id} \parallel & \circlearrowleft & d_2^A \downarrow \xrightarrow{\kappa} \downarrow d_2^B \\ \text{Ker}(f_2) \xrightarrow{k(f_2) \circ d_2^A} A_3 \xrightarrow{f_3} & & B_3 \\ 0 & \xrightarrow{(k(f_2) \circ \kappa) \cdot (\varepsilon_{f_2} \circ d_2^B) \cdot (d_2^B)_I^b} & \end{array} & \begin{array}{ccc} f_2 & \xrightarrow{\mu_{f_2} \uparrow} & B_2 \\ A_2 \xrightarrow{c(k(f_2))} \text{Coim}(f_2) \xrightarrow{j(f_2)} & & \\ d_2^A \downarrow \xrightarrow{\kappa_1} \bar{d}_2^A \downarrow \xrightarrow{\kappa_2} \downarrow d_2^B & & \\ A_3 \xrightarrow{c(k(f_3))} \text{Coim}(f_3) \xrightarrow{j(f_3)} & & B_3 \\ f_3 & \xrightarrow{\downarrow \mu_{f_3}} & \end{array} \end{array}$$

On the other hand by Proposition 5.7, if we take the compatible 2-cell $v = (k(f_2) \circ \kappa_1) \cdot (\pi_{k(f_2)} \circ \bar{d}_2^A) \cdot (\bar{d}_2^A)_I^b \in \mathbf{S}^2(k(f_2) \circ d_2^A \circ c(k(f_3)), 0)$,

$$\begin{array}{ccc} 0 & \xrightarrow{\uparrow \pi_{k(f_2)} \circ \bar{d}_2^A} & \text{Coim}(f_2) \\ \text{Ker}(f_2) \xrightarrow{k(f_2)} A_2 \xrightarrow{c(k(f_2))} & & \\ \text{id} \parallel & \circlearrowleft & d_2^A \downarrow \xrightarrow{\kappa_1} \downarrow \bar{d}_2^A \\ \text{Ker}(f_2) \xrightarrow{k(f_2) \circ d_2^A} A_3 \xrightarrow{c(k(f_3))} & & \text{Coim}(f_3) \\ 0 & \xrightarrow{\downarrow v} & \end{array}$$

then we have $\text{Cok}(k(f_2) \circ d_2^A) = [\text{Coim}(f_3), c(k(f_3)), v]$. It can be easily shown that v is compatible with μ_{f_3} and $(k(f_2) \circ \kappa) \cdot (\varepsilon_{f_2} \circ d_2^B) \cdot (d_2^B)_I^\flat$.



Since $\text{Cok}(k(f_2) \circ d_2^A) = [\text{Coim}(f_3), c(k(f_3)), v]$ and $j(f_3)$ is fully faithful by Proposition 4.4, this means (45) is 2-exact in A_3 . ■

6.6. LEMMA. Consider the following complex morphism in \mathbf{S} .

$$\begin{array}{ccccc}
 & & 0 & & \\
 & & \curvearrowright & & \\
 & & \uparrow \delta_2^A & & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 \\
 f_1 \downarrow & \lambda_1 \downarrow & f_2 \downarrow & \lambda_2 \downarrow & \downarrow f_3 \\
 0 \xrightarrow{0} & B_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{d_2^B} & B_3 \\
 & & \delta_2^B \downarrow & & \\
 & & 0 & &
 \end{array} \tag{47}$$

If the complexes are relatively 2-exact in A_2 and B_1, B_2 respectively, then the following diagram obtained by taking the kernels

$$\begin{array}{ccccc}
 & & 0 & & \\
 & & \curvearrowright & & \\
 & & \uparrow \delta_2^A & & \\
 \text{Ker}(f_1) & \xrightarrow{d_1^A} & \text{Ker}(f_2) & \xrightarrow{d_2^A} & \text{Ker}(f_3) \\
 & & & &
 \end{array} \tag{48}$$

is 2-exact in $\text{Ker}(f_2)$.

PROOF. If we decompose (47) into

$$\begin{array}{ccc}
 A_1 \xrightarrow{d_1^{A^\dagger}} \text{Ker}(d_2^A) & & \text{Ker}(d_2^A) \xrightarrow{k(d_2^A)} A_2 \xrightarrow{d_2^A} A_3 \\
 f_1 \downarrow \quad \lambda_1^\dagger \downarrow \quad \downarrow f_2 & \text{and} & f_2 \downarrow \quad \lambda_1 \downarrow \quad f_2 \downarrow \quad \lambda_2 \downarrow \quad \downarrow f_3 \\
 0 \xrightarrow{0} B_1 \xrightarrow{d_1^{B^\dagger}} \text{Ker}(d_2^B) & & \text{Ker}(d_2^B) \xrightarrow{k(d_2^B)} B_2 \xrightarrow{d_2^B} B_3
 \end{array}$$

then by (the dual of) Proposition 5.7, we have $\text{Ker}(d_2^A) = \text{Ker}(f_2)$. Since $d_1^{B^\dagger}$ is an equivalence by (the dual of) Corollary 6.2, the diagram obtained by taking the kernels of f_1 and f_2

$$\begin{array}{ccc}
 \text{Ker}(f_1) \xrightarrow{d_1^{A^\dagger}} \text{Ker}(f_2) & & \\
 k(f_1) \downarrow \quad \lambda_1^\dagger \downarrow \quad \downarrow k(f_2) & & \\
 A_1 \xrightarrow{d_1^{A^\dagger}} \text{Ker}(d_2^A) & &
 \end{array}$$

becomes a pullback diagram by (the dual of) Proposition 5.11. Since $d_1^{A^\dagger}$ is fully cofaithful, $\underline{d}_1^{A^\dagger}$ becomes also fully cofaithful by Proposition 5.12. This means (48) is 2-exact in $\text{Ker}(f_2)$. ■

6.7. LEMMA. Consider the following complex morphism in \mathbf{S} .

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \curvearrowright & \uparrow \delta_2^A & \curvearrowright & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 \\
 \text{id} \parallel & \lambda_1 \Downarrow & \text{id} \parallel & \lambda_2 \Downarrow & \downarrow f_3 \\
 A_1 & \xrightarrow{d_1^B} & A_2 & \xrightarrow{d_2^B} & B_3 \\
 & \curvearrowleft & \delta_2^B \Downarrow & \curvearrowleft & \\
 & & 0 & &
 \end{array} \tag{49}$$

If f_3 is faithful and the bottom row is 2-exact in A_2 , then the top row is also 2-exact in A_2 .

PROOF. By taking the cokernels of d_1^A and d_1^B in (49), we obtain (by Proposition 5.8)

$$\begin{array}{ccc}
 \text{Cok}(d_1^A) & \xrightarrow{\bar{d}_2^A} & A_3 \\
 \text{id} \parallel & \bar{\lambda}_2 \Downarrow & \downarrow f_3 \\
 \text{Cok}(d_1^B) & \xrightarrow{\bar{d}_2^B} & B_3.
 \end{array}$$

Since \bar{d}_2^B is fully faithful, by taking the kernels in this diagram, we obtain the following diagram.

$$\begin{array}{ccccc}
 & & 0 & \xrightarrow{0} & \text{Ker}(f_3) \\
 & & 0 \downarrow & \exists \Downarrow & \downarrow k(f_3) \\
 \text{Ker}(\bar{d}_2^A) & \xrightarrow{k(\bar{d}_2^A)} & \text{Cok}(d_1^A) \cdot \bar{d}_2^A & \longrightarrow & A_3 \\
 0 \downarrow & \exists \Downarrow & \text{id} \parallel & \bar{\lambda}_2 \Downarrow & \downarrow f_3 \\
 0 & \xrightarrow{0} & \text{Cok}(d_1^B) & \xrightarrow{\bar{d}_2^B} & B_3
 \end{array}$$

In this diagram, we have


$$\text{Ker}(\bar{d}_2^A) = \text{Ker}(\text{Ker}(\bar{d}_2^A) \longrightarrow 0) \underset{\text{Prop. 5.7}}{=} \text{Ker}(0 \longrightarrow \text{Ker}(f_3)) \underset{\text{Cor. 3.28}}{=} 0.$$

This means that the top row in (49) is 2-exact in A_2 . ■

6.8. COROLLARY. Let

$$A_1 \xrightarrow{d_1^A} A_2 \xrightarrow{d_2^A} A_3 \quad \text{and} \quad B_1 \xrightarrow{d_1^B} A_2 \xrightarrow{d_2^B} B_3$$


be two complexes, and assume that there exist 1-cells f_1, f_3 and 2-cells $\lambda_1, \lambda_2, \sigma$ as in the following diagram




where f_1 is cofaithful and f_3 is faithful. Assume they satisfy

$$\begin{aligned}
 \text{(d1)} \quad & (\lambda_1 \circ d_2^B) \cdot \delta_2^B = (f_1 \circ \sigma) \cdot (f_1)_I^\sharp \\
 \text{(d2)} \quad & (d_1^A \circ \lambda_2) \cdot \sigma = (\delta_2^A \circ f_3) \cdot (f_3)_I^\flat.
 \end{aligned}$$

Then, if the diagram




is 2-exact in A_2 , then the diagram



is also 2-exact in A_2 .


PROOF. This follows if we apply Lemma 6.7 and its dual to the following diagrams:



■

By Corollary 6.8, it can be shown that the 2-exactness plus compatibility implies the relative 2-exactness (see [7] in the case of SCG):

6.9. COROLLARY. Let $A_\bullet = (A_n, d_n, \delta_n)$ be a complex in \mathbf{S} . If



is 2-exact in A_n , then A_\bullet is relatively 2-exact in A_n .

PROOF. This follows immediately if we apply Corollary 6.8 to the following diagram (see the proof of Corollary 5.24):

$$\begin{array}{ccccc}
 A_{n-1} & \xrightarrow{d_{n-1}} & & \xrightarrow{d_n} & \text{Ker}(d_{n+1}) \\
 \downarrow c(d_{n-2}) & \nearrow \bar{\delta}_{n-1} & & \searrow \delta_{n+1} & \downarrow k(d_{n+1}) \\
 & & A_n & & \\
 \text{Cok}(d_{n-2}) & \xrightarrow{\bar{d}_{n-1}} & & \xrightarrow{d_n} & A_{n+1} \\
 & \searrow \bar{\delta}_n & & \nearrow & \\
 & & 0 & &
 \end{array}$$

■

CONSTRUCTION OF THE LONG COHOMOLOGY SEQUENCE.

6.10. DEFINITION. A complex in \mathbf{S}

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \searrow & \uparrow \varphi & \nearrow & \\
 A & \xrightarrow{f} & B & \xrightarrow{g} & C \\
 & & & &
 \end{array} \tag{50}$$

is called an extension if it is relatively 2-exact in every 0-cell.

6.11. REMARK. By Corollary 6.2 (and its dual), (50) is an extension if and only if $\text{Ker}(g) = [A, f, \varphi]$ and $\text{Cok}(f) = [C, g, \varphi]$.

6.12. DEFINITION. Let $(f_\bullet, \lambda_\bullet) : A_\bullet \rightarrow B_\bullet$ and $(g_\bullet, \kappa_\bullet) : B_\bullet \rightarrow C_\bullet$ be complex morphisms and $\varphi_\bullet = \{\varphi_n : f_n \circ g_n \Rightarrow 0\}$ be 2-cells. Then,

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \searrow & \uparrow \varphi_\bullet & \nearrow & \\
 A_\bullet & \xrightarrow{f_\bullet} & B_\bullet & \xrightarrow{g_\bullet} & C_\bullet \\
 & & & &
 \end{array} \tag{51}$$

is said to be an extension of complexes if it satisfies the following properties:

(e1) For every n , the following complex is an extension:

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \searrow & \uparrow \varphi_n & \nearrow & \\
 A_n & \xrightarrow{f_n} & B_n & \xrightarrow{g_n} & C_n \\
 & & & &
 \end{array}$$

(e2) φ_\bullet satisfies

$$(\lambda_n \circ g_{n+1}) \cdot (f_n \circ \kappa_n) \cdot (\varphi_n \circ d_n^C) \cdot (d_n^C)_I^b = (d_n^A \circ \varphi_{n+1}) \cdot (d_n^A)_I^\sharp.$$

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \searrow & \uparrow \varphi_n & \nearrow & \\
 A_n & \xrightarrow{f_n} & B_n & \xrightarrow{g_n} & C_n \\
 \downarrow d_n^A & \xrightarrow{\lambda_n} & \downarrow d_n^B & \xrightarrow{\kappa_n} & \downarrow d_n^C \\
 A_{n+1} & \xrightarrow{f_{n+1}} & B_{n+1} & \xrightarrow{g_{n+1}} & C_{n+1} \\
 & \searrow \varphi_{n+1} & & \nearrow & \\
 & & 0 & &
 \end{array}$$

Our main theorem is the following:

6.13. THEOREM. For any extension of complexes in \mathbf{S}

$$A_{\bullet} \xrightarrow{f_{\bullet}} B_{\bullet} \xrightarrow{g_{\bullet}} C_{\bullet},$$

$\begin{array}{c} 0 \\ \curvearrowright \\ \uparrow \varphi_{\bullet} \\ \curvearrowleft \end{array}$

we can construct a long 2-exact sequence:

$$\cdots \rightarrow H^n(B_{\bullet}) \rightarrow H^n(C_{\bullet}) \rightrightarrows H^{n+1}(A_{\bullet}) \rightarrow H^{n+1}(B_{\bullet}) \rightarrow \cdots$$

$\begin{array}{c} 0 \\ \uparrow \\ \downarrow \\ 0 \end{array} \quad \begin{array}{c} 0 \\ \uparrow \\ \downarrow \\ 0 \end{array}$

6.14. CAUTION. This sequence is not necessarily a complex. (See Remark 6.19.)

We prove this theorem in the rest of this section.

6.15. LEMMA. In the notation of Lemma 6.3, we have

- (1) $\text{Ker}(x_n) = H^n(A_{\bullet})$,
- (2) $\text{Cok}(x_n) = H^{n+1}(A_{\bullet})$.

PROOF. We only show (1), since (2) can be shown in the same way. In the notation of Lemma 6.3 and Remark 5.18, we can show that the factorization

$$(x_n \circ \zeta_{n+1}) \cdot \xi_n : (x_n \circ z_{n+1}) \circ k(d_{n+1}) \implies \ell_n$$

is compatible with $\varepsilon_{d_{n+1}}$ and $\mu_{n,2}$.

$$\begin{array}{ccccc}
 Q^n(A_{\bullet}) & & & & 0 \\
 \downarrow x_n \circ z_{n+1} & \nearrow \ell_n & \mu_{n,2} \uparrow & \xrightarrow{d_{n+1}} & A_{n+1} \\
 & & & & \downarrow \varepsilon_{d_{n+1}} \\
 & & & & A_{n+2} \\
 & \nearrow k(d_{n+1}) & & & \\
 \text{Ker}(d_{n+1}) & & & & 0
 \end{array}$$

$(x_n \circ \zeta_{n+1}) \cdot \xi_n$

So, by Proposition 3.20, Proposition 3.21 and the fact that z_{n+1} is fully faithful, we have $H^n(A_{\bullet}) = \text{Ker}(\ell_n, \mu_{n,2}) = \text{Ker}(x_n \circ z_{n+1}) = \text{Ker}(x_n)$. ■

6.16. LEMMA. For any extension (51) of complexes in \mathbf{S} , we can construct a complex morphism

$$\begin{array}{ccccccc}
 & & & & 0 & & \\
 & & & & \uparrow Q^n(\varphi_{\bullet}) & & \\
 Q^n(A_{\bullet}) & \xrightarrow{Q^n(f_{\bullet})} & Q^n(B_{\bullet}) & \xrightarrow{Q^n(g_{\bullet})} & Q^n(C_{\bullet}) & \xrightarrow{0} & 0 \\
 x_n^A \downarrow & & \tilde{\lambda}_n \downarrow & & x_n^B \downarrow & & \downarrow \tilde{\kappa}_n \\
 0 \longrightarrow & Z^{n+1}(A_{\bullet}) & \xrightarrow{Z^{n+1}(f_{\bullet})} & Z^{n+1}(B_{\bullet}) & \xrightarrow{Z^{n+1}(g_{\bullet})} & Z^{n+1}(C_{\bullet}) & \\
 & & & & \downarrow Z^{n+1}(\varphi_{\bullet}) & & \\
 & & & & 0 & &
 \end{array}$$

where the top line is a complex which is relatively 2-exact in $Q^n(B_\bullet)$, $Q^n(C_\bullet)$, and the bottom line is a complex which is relatively 2-exact in $Z^{n+1}(A_\bullet)$, $Z^{n+1}(B_\bullet)$.

PROOF. If we take the relative cokernels $Q^n(A_\bullet)$, $Q^n(B_\bullet)$ and $Q^n(C_\bullet)$ of the complex diagram

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & & \curvearrowright & & & & \\
 A_{n-2} & \xrightarrow{d_{n-2}^A} & A_{n-1} & \xrightarrow{d_{n-1}^A} & A_n & \xrightarrow{d_n^A} & A_{n+1} \\
 f_{n-2} \downarrow & \lambda_{n-2} \Downarrow & \downarrow f_{n-1} & \Downarrow \lambda_{n-1} & \downarrow f_n & \Downarrow \lambda_n & \downarrow f_{n+1} \\
 B_{n-2} & \xrightarrow{d_{n-2}^B} & B_{n-1} & \xrightarrow{d_{n-1}^B} & B_n & \xrightarrow{d_n^B} & B_{n+1} \\
 g_{n-2} \downarrow & \kappa_{n-2} \Downarrow & \downarrow g_{n-1} & \Downarrow \kappa_{n-1} & \downarrow g_n & \Downarrow \kappa_n & \downarrow g_{n+1} \\
 C_{n-2} & \xrightarrow{d_{n-2}^C} & C_{n-1} & \xrightarrow{d_{n-1}^C} & C_n & \xrightarrow{d_n^C} & C_{n+1} \\
 & & \curvearrowleft & & & & \\
 & & 0 & & & &
 \end{array}$$

then by (the dual of) Proposition 3.20, Proposition 5.4 and Proposition 5.5, we obtain a factorization diagram

$$\begin{array}{ccccccc}
 A_{n-1} & \xrightarrow{d_{n-1}^A} & A_n & \xrightarrow{q_n^A} & Q^n(A_\bullet) & \xrightarrow{\ell_n^A} & A_{n+1} \\
 f_{n-1} \downarrow & \lambda_{n-1} \Downarrow & f_n \downarrow & \bar{\lambda}_{n-1,1} \Downarrow & Q^n(f_\bullet) & \Downarrow \bar{\lambda}_{n-1,2} & \downarrow f_{n+1} \\
 B_{n-1} & \xrightarrow{d_{n-1}^B} & B_n & \xrightarrow{q_n^B} & Q^n(B_\bullet) & \xrightarrow{\ell_n^B} & B_{n+1} \\
 g_{n-1} \downarrow & \kappa_{n-1} \Downarrow & g_n \downarrow & \bar{\kappa}_{n-1,1} \Downarrow & Q^n(g_\bullet) & \Downarrow \bar{\kappa}_{n-1,2} & \downarrow g_{n+1} \\
 C_{n-1} & \xrightarrow{d_{n-1}^C} & C_n & \xrightarrow{q_n^C} & Q^n(C_\bullet) & \xrightarrow{\ell_n^C} & C_{n+1}
 \end{array}$$

and a 2-cell $Q^n(\varphi_\bullet) \in \mathbf{S}^2(Q^n(f_\bullet) \circ Q^n(g_\bullet), 0)$, which satisfy compatibility conditions in Proposition 5.4 and Proposition 5.5. It is also easy to see by the universality of the relative cokernels that

$$(\ell_n^A \circ \lambda_{n+1}) \cdot (\bar{\lambda}_{n-1,2} \circ d_{n+1}^B) \cdot (Q^n(f_\bullet) \circ \mu_{n,2}^B) \cdot (Q^n(f_\bullet))_I^\sharp = (\mu_{n,2}^A \circ f_{n+2}) \cdot (f_{n+2})_I^\flat.$$

Now, since

$$\begin{array}{ccccc}
 & & 0 & & \\
 & & \curvearrowright & & \\
 A_n & \xrightarrow{f_n} & B_n & \xrightarrow{g_n} & C_n \xrightarrow{0} 0
 \end{array}$$

is relatively 2-exact in B_n and C_n , we have $\text{Cok}(f_n) = [C_n, g_n, \varphi_n]$. So, from $\text{Cok}(f_n) = [C_n, g_n, \varphi_n]$ and $\text{Cok}(f_{n-1}) = [C_{n-1}, g_{n-1}, \varphi_{n-1}]$, by Proposition 5.10 we obtain

$$\text{Cok}(Q^n(f_\bullet)) = [Q^n(C_\bullet), Q^n(g_\bullet), Q^n(\varphi_\bullet)],$$

i.e. the complex

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & & \curvearrowright & & & & \\
 Q^n(A_\bullet) & \xrightarrow{Q^n(f_\bullet)} & Q^n(B_\bullet) & \xrightarrow{Q^n(g_\bullet)} & Q^n(C_\bullet) & \xrightarrow{0} & 0
 \end{array}$$

is relatively 2-exact in $Q^n(B_\bullet), Q^n(C_\bullet)$. Dually, we obtain a factorization diagram

$$\begin{array}{ccccc}
 A_n & \xrightarrow{k_{n+1}^A} & Z^{n+1}(A_\bullet) & \xrightarrow{z_{n+1}^A} & A_{n+1} & \xrightarrow{d_{n+1}^A} & A_{n+2} \\
 f_n \downarrow & \lambda_{n+1,2} \Downarrow & Z^{n+1}(f_\bullet) & \Downarrow \lambda_{n+1,1} & f_{n+1} & \Downarrow \lambda_{n+1} & f_{n+2} \\
 B_n & \xrightarrow{k_{n+1}^B} & Z^{n+1}(B_\bullet) & \xrightarrow{z_{n+1}^B} & B_{n+1} & \xrightarrow{d_{n+1}^B} & B_{n+2} \\
 g_n \downarrow & \kappa_{n+1,2} \Downarrow & Z^{n+1}(g_\bullet) & \Downarrow \kappa_{n+1,1} & g_{n+1} & \Downarrow \kappa_{n+1} & g_{n+2} \\
 C_n & \xrightarrow{k_{n+1}^C} & Z^{n+1}(C_\bullet) & \xrightarrow{z_{n+1}^C} & C_{n+1} & \xrightarrow{d_{n+1}^C} & C_{n+2}
 \end{array}$$

such that

$$\begin{aligned}
 (z_{n+1}^A \circ \lambda_{n+1}) \cdot (\lambda_{n+1,1} \circ d_{n+1}^B) \cdot (Z^{n+1}(f_\bullet) \circ \zeta_{n+1}^B) \cdot Z^{n+1}(f_\bullet)_I^\sharp \\
 = (\zeta_{n+1}^A \circ f_{n+2}) \cdot (f_{n+2})_I^b.
 \end{aligned}$$

Then, it can be shown that each of the factorizations

$$\begin{aligned}
 Q^n(f_\bullet) \circ \xi_n^B & : Q^n(f_\bullet) \circ x_n^B \circ z_{n+1}^B \implies Q^n(f_\bullet) \circ \ell_n^B \\
 (x_n^A \circ \lambda_{n+1,1}^{-1}) \cdot (\xi_n^A \circ f_{n+1}) \cdot \bar{\lambda}_{n-1,2} & : x_n^A \circ Z^{n+1}(f_\bullet) \circ z_{n+1}^B \implies Q^n(f_\bullet) \circ \ell_n^B
 \end{aligned}$$

are compatible with ζ_{n+1}^B and $(Q^n(f_\bullet) \circ \mu_{n,2}^B) \cdot (Q^n(f_\bullet))_I^\sharp$.

The image shows two commutative diagrams. The left diagram has nodes $Q^n(A_\bullet)$, $Q^n(B_\bullet)$, $Z^{n+1}(B_\bullet)$, B_{n+1} , and B_{n+2} . Arrows include $Q^n(f_\bullet) \circ \ell_n^B$, $Q^n(f_\bullet) \circ x_n^B$, d_{n+1}^B , ζ_{n+1}^B , and z_{n+1}^B . 2-cells are labeled with 0. The right diagram has nodes $Q^n(A_\bullet)$, $Z^{n+1}(B_\bullet)$, B_{n+1} , and B_{n+2} . Arrows include $Q^n(f_\bullet) \circ \ell_n^B$, $x_n^A \circ Z^{n+1}(f_\bullet)$, d_{n+1}^B , ζ_{n+1}^B , and z_{n+1}^B . 2-cells are labeled with 0. A large 2-cell $(x_n^A \circ \lambda_{n+1,1}^{-1}) \cdot (\xi_n^A \circ f_{n+1}) \cdot \bar{\lambda}_{n-1,2}$ connects the two diagrams.

So, by the universality of the relative kernel, there exists a unique 2-cell $\tilde{\lambda}_n \in \mathbf{S}^2(Q^n(f_\bullet) \circ x_n^B, x_n^A \circ Z^{n+1}(f_\bullet))$ such that

$$(\tilde{\lambda}_n \circ z_{n+1}^B) \cdot (x_n^A \circ \lambda_{n+1,1}^{-1}) \cdot (\xi_n^A \circ f_{n+1}) \cdot \bar{\lambda}_{n-1,2} = Q^n(f_\bullet) \circ \xi_n^B.$$

This $\tilde{\lambda}_n$ also satisfies $(q_n^A \circ \tilde{\lambda}_n) \cdot (\eta_n^A \circ Z^{n+1}(f_\bullet)) \cdot \lambda_{n+1,2} = (\bar{\lambda}_{n-1,1} \circ x_n^B) \cdot (f_n \circ \eta_n^B)$ (see Remark 6.17). Similarly, we obtain a 2-cell $\tilde{\kappa}_n \in \mathbf{S}^2(Q^n(g_\bullet) \circ x_n^C, x_n^B \circ Z^{n+1}(g_\bullet))$ such that

$$(\tilde{\kappa}_n \circ z_{n+1}^C) \cdot (x_n^B \circ \kappa_{n+1,1}^{-1}) \cdot (\xi_n^B \circ g_{n+1}) \cdot \bar{\kappa}_{n-1,2} = Q^n(g_\bullet) \circ \xi_n^C.$$

In the rest, we show the following:

$$(Q^n(f_\bullet) \circ \tilde{\kappa}_n) \cdot (\tilde{\lambda}_n \circ Z^{n+1}(g_\bullet)) \cdot (x_n^A \circ Z^{n+1}(\varphi_\bullet)) \cdot (x_n^A)_I^\sharp = (Q^n(\varphi_\bullet) \circ x_n^C) \cdot (x_n^C)_I^\flat. \quad (52)$$

We have the following equalities:

$$\begin{aligned} & (Q^n(f_\bullet) \circ \tilde{\kappa}_n \circ z_{n+1}^C) \cdot (\tilde{\lambda}_n \circ Z^{n+1}(g_\bullet) \circ z_{n+1}^C) \\ &= (Q^n(f_\bullet) \circ Q^n(g_\bullet) \circ \xi_n^C) \cdot (Q^n(f_\bullet) \circ \bar{\kappa}_{n-1,2}^{-1}) \cdot (\bar{\lambda}_{n-1,2}^{-1} \circ g_{n+1}) \\ & \quad \cdot ((\xi_n^A)^{-1} \circ f_{n+1} \circ g_{n+1}) \cdot (x_n^A \circ \underline{\lambda}_{n+1,1} \circ g_{n+1}) \cdot (x_n^A \circ Z^{n+1}(f_\bullet) \circ \underline{\kappa}_{n+1,1}), \\ & (Q^n(\varphi_\bullet) \circ x_n^C \circ z_{n+1}^C) \cdot (x_n^C \circ z_{n+1}^C)_I^\flat \cdot (x_n^A \circ z_{n+1}^A)_I^{\sharp-1} \cdot (x_n^A \circ z_{n+1}^A \circ \varphi_{n+1}^{-1}) \\ &= (Q^n(f_\bullet) \circ Q^n(g_\bullet) \circ \xi_n^C) \cdot (Q^n(f_\bullet) \circ \bar{\kappa}_{n-1,2}^{-1}) \cdot (\bar{\lambda}_{n-1,2}^{-1} \circ g_{n+1}) \\ & \quad \cdot ((\xi_n^A)^{-1} \circ f_{n+1} \circ g_{n+1}), \\ & (z_{n+1}^C)_I^{\flat-1} = (x_n^A \circ z_{n+1}^A)_I^{\sharp-1} \cdot (x_n^A \circ z_{n+1}^A \circ \varphi_{n+1}^{-1}) \cdot (x_n^A \circ \underline{\lambda}_{n+1,1} \circ g_{n+1}) \\ & \quad \cdot (x_n^A \circ Z^{n+1}(f_\bullet) \circ \underline{\kappa}_{n+1,1}) \cdot (x_n^A \circ Z^{n+1}(\varphi_\bullet) \circ z_{n+1}^C) \cdot ((x_n^A)_I^\sharp \circ z_{n+1}^C). \end{aligned}$$

From these equalities and the faithfulness of z_{n+1}^C , we obtain (52). ■

6.17. REMARK. It can be also shown that $\tilde{\lambda}_n$ in the proof of Lemma 6.16 satisfies

$$(x_n^A \circ \tilde{\lambda}_n) \cdot (\eta_n^A \circ Z^{n+1}(f_\bullet)) \cdot \underline{\lambda}_{n+1,2} = (\bar{\lambda}_{n-1,1} \circ x_n^B) \cdot (f_n \circ \eta_n^B).$$

By Lemma 6.15 and Lemma 6.16, Theorem 6.13 is reduced to the following Proposition:

6.18. PROPOSITION. Consider the following diagram in \mathbf{S} , where $(A_\bullet, d_\bullet^A, \delta_\bullet^A)$ is a complex which is relatively 2-exact in A_2 and A_3 , and $(B_\bullet, d_\bullet^B, \delta_\bullet^B)$ is a complex which is relatively 2-exact in B_1 and B_2 .

$$\begin{array}{ccccccc} & & & 0 & & & \\ & & & \uparrow \delta_2^A & & & \\ & & & \uparrow \delta_2^A & \xrightarrow{d_2^A} & & \\ A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{0} & 0 \\ & f_1 \downarrow & \lambda_1 \downarrow & f_2 \downarrow & \lambda_2 \downarrow & f_3 \downarrow & \\ 0 & \xrightarrow{0} & B_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{d_2^B} & B_3 \\ & & & \delta_2^B \downarrow & & & \\ & & & 0 & & & \end{array}$$

Assume $f_\bullet : A_\bullet \rightarrow B_\bullet$ is a complex morphism. Then there exist $d \in \mathbf{S}^1(\text{Ker}(f_3), \text{Cok}(f_1))$, $\alpha \in \mathbf{S}^2(d_2^A \circ d, 0)$ and $\beta \in \mathbf{S}^2(d \circ \bar{d}_1^B, 0)$ such that the sequence

$$\begin{array}{ccccccc} & & & 0 & & & \\ & & & \uparrow \delta_2^A & & & \\ \text{Ker}(f_1) & \xrightarrow{d_1^A} & \text{Ker}(f_2) & \xrightarrow{d_2^A} & \text{Ker}(f_3) & \xrightarrow{d} & \text{Cok}(f_1) & \xrightarrow{\bar{d}_1^B} & \text{Cok}(f_2) & \xrightarrow{\bar{d}_2^B} & \text{Cok}(f_3) \\ & & & \alpha \downarrow & & & \beta \downarrow & & & & \\ & & & 0 & & & 0 & & & & \end{array} \quad (53)$$

is 2-exact in $\text{Ker}(f_2), \text{Ker}(f_3), \text{Cok}(f_1), \text{Cok}(f_2)$.

6.19. **REMARK.** This sequence does not necessarily become a complex. Indeed, for a relatively exact 2-category \mathbf{S} , the following are shown to be equivalent by an easy diagrammatic argument:

- (i) Any (53) obtained in Proposition 6.18 becomes a complex.
- (ii) For any $f \in \mathbf{S}^1(A, B)$,

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \curvearrowright & \uparrow \varepsilon_f & \curvearrowleft & \\
 \text{Ker}(f) & \xrightarrow{k(f)} & A & \xrightarrow{f} & B \xrightarrow{c(f)} \text{Cok}(f) \\
 & & \downarrow \pi_f & & \\
 & & 0 & &
 \end{array} \tag{54}$$

is a complex.

(Indeed, if (54) is a complex for each of f_1, f_2 and f_3 , then (53) becomes a complex.)

Thus if \mathbf{S} satisfies (ii), then the long cohomology sequence in Theorem 6.13 becomes a complex. But this assumption is a bit too strong, since it is not satisfied by SCG. This is pointed out by the referee.

PROOF OF PROPOSITION 6.18. Put $\text{Ker}(d_2^A \circ f_3) = [K, k, \zeta]$. If we take the kernel of the diagram

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & \curvearrowright & \uparrow \delta_2^A & \curvearrowleft & & & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{0} & 0 \\
 & \downarrow \xi_0 & \downarrow d_2^A \circ f_3 & \downarrow \circ & \downarrow f_3 & & \\
 0 & \xrightarrow{0} & B_3 & \xrightarrow{\text{id}} & B_3 & \xrightarrow{0} & 0
 \end{array} \tag{55}$$

where $\xi_0 := (\delta_2^A \circ f_3) \cdot (f_3)_I^\flat$, then by Proposition 5.5 we obtain a diagram

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & \curvearrowright & \uparrow \alpha_2 & \curvearrowleft & & & \\
 A_1 & \xrightarrow{k_1} & K & \xrightarrow{k_2} & \text{Ker}(f_3) & & \\
 \text{id} \parallel & \downarrow \xi_1 & \downarrow k & \downarrow \xi_2 & \downarrow k(f_3) & & \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{0} & 0 \\
 & \downarrow \delta_2^A & \downarrow & \downarrow & & & \\
 & & 0 & & & &
 \end{array} \tag{56}$$

which satisfies

$$\begin{aligned}
 (k_2 \circ \varepsilon_{f_3}) \cdot (k_2)_I^\# &= (\xi_2 \circ f_3) \cdot \zeta \\
 (\xi_1 \circ d_2^A \circ f_3) \cdot \xi_0 &= (k_1 \circ \zeta) \cdot (k_1)_I^\# \\
 (k_1 \circ \xi_2) \cdot (\xi_1 \circ d_2^A) \cdot \delta_2^A &= (\alpha_2 \circ k(f_3)) \cdot k(f_3)_I^\flat.
 \end{aligned}$$

By Lemma 6.6,

$$\begin{array}{ccc}
 & & 0 \\
 & \curvearrowright & \uparrow \alpha_2 \\
 A_1 & \xrightarrow{k_1} & K \xrightarrow{k_2} \text{Ker}(f_3)
 \end{array}$$

is 2-exact in K . On the other hand, by (the dual of) Proposition 5.11,

$$\begin{array}{ccc} K & \xrightarrow{k_2} & \text{Ker}(f_3) \\ k \downarrow & \xi_2 \Downarrow & \downarrow f_3 \\ A_2 & \xrightarrow{d_2^A} & A_3 \end{array}$$

is a pullback diagram, and k_2 becomes cofaithful since d_2^A is cofaithful. Thus, we have $\text{Cok}(k_1) = [\text{Ker}(f_3), k_2, \alpha_2]$ by Lemma 6.4. Dually, if we put $\text{Cok}(f_1 \circ d_1^B) = [Q, q, \rho]$, then we obtain the following diagram

$$\begin{array}{ccccccc} 0 & \xrightarrow{0} & A_1 & \xrightarrow{\text{id}} & A_1 & \xrightarrow{0} & 0 \\ & & f_1 \downarrow & \circlearrowleft & f_1 \circ d_1^B \downarrow & \Downarrow \eta_0 & \downarrow 0 \\ 0 & \xrightarrow{0} & B_1 & \xrightarrow{d_1^B} & B_2 & \xrightarrow{d_2^B} & B_3 \\ & & c(f_1) \downarrow & \eta_1 \Downarrow & q \downarrow & \eta_2 \Downarrow & \parallel \text{id} \\ & & \text{Cok}(f_1) & \xrightarrow{q_1} & Q & \xrightarrow{q_2} & B_3 \\ & & & \searrow \beta_2 \Downarrow & \nearrow & & \\ & & & & 0 & & \end{array}$$

which satisfies

$$\begin{aligned} \eta_0 &= (f_1)_I^{\sharp-1} \cdot (f_1 \circ \delta_2^{B-1}) \\ \rho &= (f_1 \circ \eta_1) \cdot (\pi_{f_1} \circ q_1) \circ (q_1)_I^b \\ \text{id}_0 &= \eta_0 \cdot (f_1 \circ d_1^B \circ \eta_2) \cdot (\rho \circ q_2) \cdot (q_2)_I^b \\ \delta_2^B &= (d_1^B \circ \eta_2) \cdot (\eta_1 \circ q_2) \cdot (c(f_1) \circ \beta_2) \cdot c(f_1)_I^{\sharp}, \end{aligned}$$

and we have $\text{Ker}(q_2) = [\text{Cok}(f_1), q_1, \beta_2]$. (The “un-duality” in appearance is simply because of the direction of the 2-cells.) Thus, we obtain complex morphisms:

$$\begin{array}{ccccccc} A_1 & \xrightarrow{\text{id}} & A_1 & \xrightarrow{f_1} & B_1 & \xrightarrow{c(f_1)} & \text{Cok}(f_1) \\ k_1 \downarrow & \xRightarrow{\xi_1} & d_1^A \downarrow & \xRightarrow{\lambda_1} & d_1^B \downarrow & \xRightarrow{\eta_1} & \downarrow q_1 \\ K & \xrightarrow{k} & A_2 & \xrightarrow{f_2} & B_2 & \xrightarrow{q} & Q \\ k_2 \downarrow & \xRightarrow{\xi_2} & d_2^A \downarrow & \xRightarrow{\lambda_2} & d_2^B \downarrow & \xRightarrow{\eta_2} & \downarrow q_2 \\ \text{Ker}(f_3) & \xrightarrow{k(f_3)} & A_3 & \xrightarrow{f_3} & B_3 & \xrightarrow{\text{id}} & B_3 \end{array}$$

If we put

$$\begin{aligned} c &:= k \circ f_2 \circ q \\ \alpha_K &:= (\xi_1 \circ f_2 \circ q) \cdot (\lambda_1 \circ q) \cdot \rho \\ \beta_Q &:= (k \circ f_2 \circ \eta_2^{-1}) \cdot (k \circ \lambda_2^{-1}) \cdot \zeta, \end{aligned}$$

then, it can be shown that the following diagram is a complex.

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \curvearrowright & \alpha_K \uparrow & \curvearrowright & \\
 A_1 & \xrightarrow{k_1} & K & \xrightarrow{c} & Q \xrightarrow{q_2} B_3 \\
 & & & & \downarrow \beta_Q \\
 & & & & 0
 \end{array}$$

Since $\text{Cok}(k_1) = [\text{Ker}(f_3), k_2, \alpha_2]$ as already shown, we have a factorization diagram

$$\begin{array}{ccccc}
 & & \text{Ker}(f_3) & & \\
 & \curvearrowright & & \curvearrowright & \\
 A_1 & \xrightarrow{k_1} & K & \xrightarrow{c} & Q \xrightarrow{q_2} B_3 \\
 & & \alpha_K \downarrow & & \downarrow \beta_Q \\
 & & & & 0
 \end{array}$$

which satisfies

$$\begin{aligned}
 (k_1 \circ \bar{\alpha}_K) \cdot \alpha_K &= (\alpha_2 \circ \bar{c}) \cdot (\bar{c})_I^\flat \\
 (\bar{\alpha}_K \circ q_2) \cdot \beta_Q &= (k_2 \circ \bar{\beta}_Q) \cdot (k_2)_I^\sharp.
 \end{aligned}$$

Similarly, since $\text{Ker}(q_2) = [\text{Cok}(f_1), q_1, \beta_2]$, we have a factorization diagram

$$\begin{array}{ccccc}
 & & 0 & & 0 \\
 & \curvearrowright & \alpha_K \uparrow & \curvearrowright & \beta_Q \uparrow \\
 A_1 & \xrightarrow{k_1} & K & \xrightarrow{c} & Q \xrightarrow{q_2} B_3 \\
 & & \alpha_K \downarrow & & \downarrow \beta_2 \\
 & & & & \text{Cok}(f_1)
 \end{array}$$


which satisfies

$$\begin{aligned}
 (\underline{\beta}_Q \circ q_2) \cdot \beta_Q &= (\underline{c} \circ \beta_2) \cdot (\underline{c})_I^\sharp \\
 (k_1 \circ \underline{\beta}_Q) \cdot \alpha_K &= (\underline{\alpha}_K \circ q_1) \cdot (q_1)_I^\flat.
 \end{aligned}$$

Then, there exist $d \in \mathbf{S}^1(\text{Ker}(f_3), \text{Cok}(f_1))$, $\alpha^\dagger \in \mathbf{S}^2(k_2 \circ d, \underline{c})$ and $\beta^\dagger \in \mathbf{S}^2(d \circ q_1, \bar{c})$ such that

$$\begin{aligned}
 (k_1 \circ \alpha^\dagger) \cdot \underline{\alpha}_K &= (\alpha_2 \circ d) \cdot d_I^\flat \\
 (\beta^\dagger \circ q_2) \cdot \bar{\beta}_Q &= (d \circ \beta_2) \cdot d_I^\sharp \\
 (k_2 \circ \beta^\dagger) \cdot \bar{\alpha}_K &= (\alpha^\dagger \circ q_1) \cdot \underline{\beta}_Q
 \end{aligned}$$

(note that $\text{Cok}(k_1) = [\text{Ker}(f_3), k_2, \alpha_2]$ and $\text{Ker}(q_2) = [\text{Cok}(f_1), q_1, \beta_2]$ (cf. Lemma 6.3)):



Applying (the dual of) Proposition 5.8 to the diagram

$$\begin{array}{ccccc}
 \text{Ker}(f_1) & \xrightarrow{d_1^A} & \text{Ker}(f_2) & \xrightarrow{d_2^A} & \text{Ker}(f_3) \\
 k(f_1) \downarrow & \lambda_1 \downarrow & k(f_2) \downarrow & \lambda_2 \downarrow & \downarrow k(f_3) \\
 A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 \\
 0 \downarrow & \xi_0 \downarrow & d_2^A \circ f_3 \downarrow & \circ & \downarrow f_3 \\
 0 & \xrightarrow{0} & B_3 & \xrightarrow{\text{id}} & B_3,
 \end{array}$$

we see that there exist $k' \in \mathbf{S}^1(\text{Ker}(f_2), K)$, $\xi'_1 \in \mathbf{S}^2(\underline{d}_1^A \circ k', k(f_1) \circ k_1)$, $\xi'_2 \in \mathbf{S}^2(\underline{d}_2^A, k' \circ k_2)$ and $\xi \in \mathbf{S}^2(k' \circ k, k(f_2))$ such that

$$\begin{aligned}
 \underline{\delta}_2^A &= (\underline{d}_1^A \circ \xi'_2) \cdot (\xi'_1 \circ k_2) \cdot (k(f_1) \circ \alpha_2) \cdot k(f_1)_I^\sharp \\
 \lambda_2 &= (\xi'_2 \circ k(f_3)) \cdot (k' \circ \xi_2) \cdot (\xi \circ d_2^A) \\
 (\underline{d}_1^A \circ \xi) \cdot \lambda_1 &= (\xi'_1 \circ k) \cdot (k(f_1) \circ \xi_1).
 \end{aligned}$$

Similarly, there exist $q' \in \mathbf{S}^1(Q, \text{Cok}(f_2))$, $\eta'_1 \in \mathbf{S}^2(q_1 \circ q', \bar{d}_1^B)$, $\eta'_2 \in \mathbf{S}^2(q_2 \circ c(f_3), q' \circ \bar{d}_2^B)$ and $\eta \in \mathbf{S}^2(q \circ q', c(f_2))$ such that

$$\begin{aligned}
 (\beta_2 \circ c(f_3)) \cdot c(f_3)_I^b &= (q_1 \circ \eta'_2) \cdot (\eta'_1 \circ \bar{d}_2^B) \cdot \bar{\delta}_2^B \\
 (\eta_1 \circ q') \cdot (c(f_1) \circ \eta'_1) &= (d_1^B \circ \eta) \cdot \bar{\lambda}_1 \\
 (\eta_2 \circ c(f_3)) \cdot (q \circ \eta'_2) \cdot (\eta \circ \bar{d}_2^B) &= \bar{\lambda}_2.
 \end{aligned}$$

If we put

$$\alpha_0 := (\underline{d}_2^A \circ \beta^\dagger) \cdot (\xi'_2 \circ \bar{c}) \cdot (k' \circ \bar{\alpha}_K) \cdot (\xi \circ f_2 \circ q) \cdot (\varepsilon_{f_2} \circ q) \cdot q_I^b,$$

then it can be shown that $\alpha_0 : \underline{d}_2^A \circ d \circ q_1 \implies 0$ is compatible with β_2 .

$$\begin{array}{ccccc}
 & & 0 & & \\
 & & \alpha_0 \uparrow & & \\
 \text{Ker}(f_2) & \xrightarrow{\underline{d}_2^A \circ d} & \text{Cok}(f_1) & \xrightarrow{q_1} & Q & \xrightarrow{q_2} & B_3 \\
 & & & & \downarrow \beta_2 & & \\
 & & & & 0 & &
 \end{array}$$

So by Lemma 3.19, there exists $\alpha \in \mathbf{S}^2(d_2^A \circ d, 0)$ such that

$$(\alpha \circ q_1) \cdot (q_1)_I^b = \alpha_0.$$

Dually, if we put

$$\beta_0 := (\alpha^\dagger \circ \bar{d}_1^B) \cdot (\underline{c} \circ \eta_1^{-1}) \cdot (\underline{\beta}_Q \circ q') \cdot (k \circ f_2 \circ \eta) \cdot (k \circ \pi_{f_2}) \cdot k_I^\sharp,$$

then $\beta_0 : k_2 \circ d \circ \bar{d}_1^B \Longrightarrow 0$ is compatible with α_2 , and there exists $\beta \in \mathbf{S}^2(d \circ \bar{d}_1^B, 0)$ such that

$$(k_2 \circ \beta) \cdot (k_2)_I^\sharp = \beta_0.$$

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & \curvearrowright & \uparrow \delta_2^A & \curvearrowright & \uparrow \beta & \curvearrowright & \\ \text{Ker}(f_1) & \xrightarrow{d_1^A} & \text{Ker}(f_2) & \xrightarrow{d_2^A} & \text{Ker}(f_3) & \xrightarrow{d} & \text{Cok}(f_1) & \xrightarrow{\bar{d}_1^B} & \text{Cok}(f_2) & \xrightarrow{\bar{d}_2^B} & \text{Cok}(f_3) \\ & & \downarrow \alpha & & \downarrow \delta_2^B & & \downarrow \delta_2^B & & & & \\ & & 0 & & 0 & & 0 & & & & \end{array}$$

In the rest, we show that this is 2-exact in $\text{Ker}(f_2), \text{Ker}(f_3), \text{Cok}(f_1), \text{Cok}(f_2)$. We show only the 2-exactness in $\text{Ker}(f_2)$ and $\text{Ker}(f_3)$, since the rest can be shown dually. The 2-exactness in $\text{Ker}(f_2)$ follows immediately from Lemma 6.6. So, we show the 2-exactness in $\text{Ker}(f_3)$. Since we have $\text{Cok}(d_1^A) = [A_3, d_2^A, \delta_2^A]$ and $\text{Cok}(f_1 \circ d_2^B) = [Q, q, \rho]$, there exists a factorization (ℓ, ϖ_1)

$$\begin{array}{ccccccc} & & 0 & & & & \\ & \curvearrowright & \uparrow \delta_2^A & \curvearrowright & & & \\ A_1 & \xrightarrow{d_1^A} & A_2 & \xrightarrow{d_2^A} & A_3 & \xrightarrow{0} & 0 \\ \text{id} \parallel & \lambda_1 \downarrow & f_2 \downarrow & \varpi_1 \downarrow & \downarrow \ell & & \\ A_1 & \xrightarrow{f_1 \circ d_1^B} & B_2 & \xrightarrow{q} & Q & \xrightarrow{0} & 0 \\ & & \rho \downarrow & & & & \\ & & 0 & & & & \end{array} \quad (57)$$

such that


$$(d_1^A \circ \varpi_1) \cdot (\lambda_1 \circ q) \cdot \rho = (\delta_2^A \circ \ell) \cdot \ell_I^b.$$

Applying Lemma 6.5 to diagram (57), we see that the following diagram becomes 2-exact in A_3 :

$$\begin{array}{ccc} & & 0 \\ & \curvearrowright & \uparrow \text{---} \\ \text{Ker}(f_2) & \xrightarrow{k(f_2) \circ d_2^A} & A_3 & \xrightarrow{\ell} & Q \\ & & \uparrow \text{---} & \nearrow & \\ & & & & (k(f_2) \circ \varpi_1) \cdot (\varepsilon_{f_2 \circ q}) \cdot q_I^b \end{array} \quad (58)$$

Then it can be shown that $(\varpi_1 \circ q_2) \cdot (f_2 \circ \eta_2^{-1}) : d_2^A \circ \ell \circ q_2 \Longrightarrow f_2 \circ d_2^B$ is compatible with

δ_2^A and $(\lambda_1 \circ d_2^B) \cdot (f_1 \circ d_2^B) \cdot (f_1)_I^\sharp$. So, comparing the following two factorizations




we see there exists a unique 2-cell $\varpi_2 \in \mathbf{S}^2(\ell \circ q_2, f_3)$ such that

$$(d_2^A \circ \varpi_2) \cdot \lambda_2 = (\varpi_1 \circ q_2) \cdot (f_2 \circ \eta_2^{-1}).$$

Then it can be shown that each of the two factorizations

- (1) $(\alpha^\dagger \circ q_1) \cdot \underline{\beta}_Q : k_2 \circ d \circ q_1 \implies c$
- (2) $(\xi_2 \circ \ell) \cdot (k \circ \varpi_1) : k_2 \circ k(f_3) \circ \ell \implies k \circ f_2 \circ q = c$

is compatible with α_2 and α_K .



So there exists a unique 2-cell $\varpi_3 \in \mathbf{S}^2(d \circ q_1, k(f_3) \circ \ell)$ such that

$$(k_2 \circ \varpi_3) \cdot (\xi_2 \circ \ell) \cdot (k \circ \varpi_1) = (\alpha^\dagger \circ q_1) \cdot \underline{\beta}_Q$$

(recall that $\text{Cok}(k_1) = [\text{Ker}(f_3), k_2, \alpha_2]$). Then we have $(\varpi_3 \circ q_2) \cdot (k(f_3) \circ \varpi_2) \cdot \varepsilon_{f_3} = (d \circ \beta_2) \cdot d_7^\sharp$.

$$\begin{array}{ccc}
 \text{Ker}(f_3) & \xrightarrow{d} & \text{Cok}(f_1) \\
 \left(\begin{array}{ccc}
 \downarrow k(f_3) & \varpi_3 \downarrow & \downarrow q_1 \\
 \varepsilon_{f_3} \downarrow & \downarrow \ell & \downarrow \beta_2 \\
 A_3 & \xrightarrow{\ell} & Q \\
 \downarrow f_3 & \varpi_2 \downarrow & \downarrow q_2 \\
 B_3 & \xlongequal{\text{id}} & B_3
 \end{array} \right) & & 0
 \end{array} \tag{59}$$

By taking kernels of d, ℓ and id_{B_3} in (59), we obtain the following diagram.

$$\begin{array}{ccccc}
 \text{Ker}(d) & \xrightarrow{k(d)} & \text{Ker}(f_3) & \xrightarrow{d} & \text{Cok}(f_1) \\
 \downarrow k(f_3) & & \downarrow \varpi_3 & & \downarrow q_1 \\
 \text{Ker}(\ell) & \xrightarrow{k(\ell)} & A_3 & \xrightarrow{\ell} & Q \\
 \downarrow 0 & & \downarrow f_3 & & \downarrow q_2 \\
 0 & \xrightarrow{0} & B_3 & \xrightarrow{\text{id}} & B_3
 \end{array}$$

Since $\text{Ker}(0 : \text{Ker}(\ell) \rightarrow 0) = \text{Ker}(d)$ by (the dual of) Proposition 5.7, so $\underline{k(f_3)}$ becomes an equivalence. On the other hand, the following is a complex morphism, where $s := k(f_2) \circ d_2^A$.

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \xrightarrow{d_2^A} & \uparrow \alpha & \xrightarrow{d} & \\
 \text{Ker}(f_2) & \xrightarrow{d_2^A} & \text{Ker}(f_3) & \xrightarrow{d} & \text{Cok}(f_1) \\
 \text{id} \parallel & \downarrow \lambda_2 & \downarrow k(f_3) & \downarrow \varpi_3 & \downarrow q_1 \\
 \text{Ker}(f_2) & \xrightarrow{s} & A_3 & \xrightarrow{\ell} & Q \\
 & \searrow & \downarrow & \nearrow & \\
 & & 0 & & (k(f_2) \circ \varpi_1) \cdot (\varepsilon_{f_2} \circ q) \cdot q_1^b
 \end{array} \tag{60}$$

Thus by taking kernels of d and ℓ in diagram (60), we obtain the following factorization by (the dual of) Proposition 5.8.

$$\begin{array}{ccccc}
 & & d_2^A & & \\
 & \xrightarrow{\exists (d_2^A)^\dagger} & \uparrow \exists \alpha & \xrightarrow{k(d)} & \\
 \text{Ker}(f_2) & \xrightarrow{\exists (d_2^A)^\dagger} & \text{Ker}(d) & \xrightarrow{k(d)} & \text{Ker}(f_3) \\
 \text{id} \parallel & \downarrow \exists & \downarrow k(f_3) & \downarrow \varpi_3 & \downarrow k(f_3) \\
 \text{Ker}(f_2) & \xrightarrow{\exists s} & \text{Ker}(\ell) & \xrightarrow{k(\ell)} & A_3 \\
 & \searrow & \downarrow \exists & \nearrow & \\
 & & s & &
 \end{array}$$

Since (58) is 2-exact in A_3 , so \underline{s} becomes fully cofaithful. Since $\underline{k(f_3)}$ is an equivalence, this means $(\underline{d_2^A})^\dagger$ is fully cofaithful, and

$$\begin{array}{ccccc}
 & & 0 & & \\
 & \xrightarrow{\underline{d_2^A}} & \uparrow \alpha & \xrightarrow{d} & \\
 \text{Ker}(f_2) & \xrightarrow{\underline{d_2^A}} & \text{Ker}(f_3) & \xrightarrow{d} & \text{Cok}(f_1)
 \end{array}$$

becomes 2-exact in $\text{Ker}(f_3)$. ■

References

[1] D. Bourn, E. M. Vitale, Extensions of symmetric cat-groups, *Homol. Homotopy Appl.* **4** (2002), 103-162.

- [2] M. Dupont, E. M. Vitale, Proper factorization systems in 2-categories, *J. Pure Applied Algebra* **179** (2003), 65-86.
- [3] S. Kasangian, E. M. Vitale, Factorization systems for symmetric cat-groups, *Theory and Application of Categories* **7** (2000), 47-70.
- [4] S. Mac Lane, "Categories for the working mathematician", Graduate Texts in Math., Vol. 5, Springer-Verlag, New York, 1998.
- [5] B. Pareigis, Non-additive ring and module theory I, General theory of monoids, *Publ. Math. Debrecen* **24** (1977), 189-204.
- [6] B. Pareigis, Non-additive ring and module theory IV, Brauer group of a symmetric monoidal category, in "Brauer groups", Lecture Notes in Math., Vol. 549, pp. 112-133, Springer-Verlag, Berlin, 1976.
- [7] A. del Río, J. Martínez-Moreno, E. M. Vitale, Chain complexes of symmetric categorical groups, *J. Pure Applied Algebra* **196** (2004), 279-312.
- [8] E. M. Vitale, The Brauer and Brauer-Taylor groups of a symmetric monoidal category, *Cahiers de Topologie Géométrie Différentielle Categoricals* **37** (1996), 91-122.
- [9] E. M. Vitale, A Picard-Brauer exact sequence of categorical groups, *J. Pure Applied Algebra* **175** (2002), 383-408.

*Graduate School of Mathematical Sciences,
The University of Tokyo 3-8-1 Komaba,
Meguro, Tokyo, 153-8914 Japan*
Email: deutsche@ms.u-tokyo.ac.jp

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at <ftp://ftp.tac.mta.ca/pub/tac/html/volumes/20/16/20-16.{dvi,ps,pdf}>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS. The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\text{e}$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

$\text{T}_{\text{E}}\text{X}$ TECHNICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT $\text{T}_{\text{E}}\text{X}$ EDITOR. Gavin Seal, McGill University: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown \(at\) btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Aurelio Carboni, Università dell Insubria: aurelio.carboni@uninsubria.it

Valeria de Paiva, Cuill Inc.: valeria@cuill.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Martin Hyland, University of Cambridge: M.Hyland@dpms.cam.ac.uk

P. T. Johnstone, University of Cambridge: ptj@dpms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, University of Western Sydney: s.lack@uws.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@acsu.buffalo.edu

Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Brooke Shipley, University of Illinois at Chicago: bshipley@math.uic.edu

James Stasheff, University of North Carolina: jds@math.unc.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca