

HOPF MONOIDAL COMONADS

DIMITRI CHIKHLADZE, STEPHEN LACK, AND ROSS STREET

ABSTRACT. We generalize to the context internal to an autonomous monoidal bicategory the work of Bruguières, Virelizier, and the second-named author on lifting closed structure on a monoidal category to the category of Eilenberg-Moore algebras for an opmonoidal monad. The result then applies to quantum categories and bialgebroids.

1. Introduction

Early workers on monoidal categories recognized the importance of functors $F : \mathcal{V} \rightarrow \mathcal{W}$ not preserving the tensor, rather, having only coherent natural families of morphisms

$$\varphi : FX \otimes FY \longrightarrow F(X \otimes Y).$$

There is also a morphism $\varphi_0 : I \rightarrow FI$ connecting the tensor units. We follow Eilenberg and Kelly [3] in using the term *monoidal functor* for this structure on F . A monoidal category \mathcal{V} is *right closed* when each pair of objects X and Z has a right internal hom Z^X ; that is, there is an evaluation morphism $\text{ev} : X \otimes Z^X \rightarrow Z$ determining a bijection

$$\mathcal{V}(Y, Z^X) \cong \mathcal{V}(X \otimes Y, Z), \quad f \mapsto \text{ev}(1_X \otimes f).$$

The monoidal structure φ on F can be expressed in terms of right closed structure

$$\varphi^r : F(Z^X) \longrightarrow (FZ)^{FX}$$

defined by the equality

$$\begin{aligned} FX \otimes F(Z^X) &\xrightarrow{1 \otimes \varphi^r} FX \otimes (FZ)^{FX} \xrightarrow{\text{ev}} FZ \\ &= FX \otimes F(Z^X) \xrightarrow{\varphi} F(X \otimes Z^X) \xrightarrow{F(\text{ev})} FZ . \end{aligned}$$

It was also recognized that tensor preserving (up to isomorphism) was an important special case; these are the *strong monoidal functors* where “strong” means that φ and φ_0

The first-named author is grateful for an International Macquarie University Research Scholarship, and Faculty of Science postgraduate research funds; all acknowledge the support of the Australian Research Council through Discovery Grant DP0771252.

Received by the editors 2010-10-14 and, in revised form, 2010-11-26.

Transmitted by R. J. Wood. Published on 2010-11-30.

2000 Mathematics Subject Classification: 18D10.

Key words and phrases: quantum category; monoidal comonad; monoidal bicategory; quantum groupoid; bialgebroid; Hopf monoid; closed category; pseudomonoid.

© Dimitri Chikhladze, Stephen Lack, and Ross Street, 2010. Permission to copy for private use granted.

are invertible. Internal hom preserving was considered too rare to warrant terminology. However, it does occur in interesting cases such as the forgetful functor from G -sets to sets where G is a group, a fact that distinguishes groups G from monoids. This observation was important to both the papers [10] and [19] in defining “Hopf concepts” (in the sense of existence of antipodes). Following [12], we call a monoidal functor *strong right closed* when φ^r and φ_0 are invertible. While strong monoidal functors do preserve duals, they are generally not strong right closed.

Bruguières and Virelizier [4] began a program to generalize the theory of Hopf monoids in a braided monoidal category to appropriate monads on a not-necessarily-braided monoidal category. That paper looks at the case where the monoidal category is autonomous (duals exist). They define the concept of (right) antipode for an opmonoidal monad on a (right) autonomous monoidal category; such an opmonoidal monad with an antipode they call a Hopf monad. In later work [1, 2] with the second author of the present paper they found a characterization of the existence of an antipode on the opmonoidal monad purely in terms of the monoidal structure of the autonomous monoidal category. (This corresponds to the fact for bimonoids that an antipode exists if and only if the fusion map is invertible.) The antipode-free formulation allowed them to extend the Hopf concept to opmonoidal monads on closed monoidal categories. For a right closed monoidal category they then showed that this Hopf condition on an opmonoidal monad was equivalent to the right closedness of the monoidal category of Eilenberg-Moore algebras together with the strong right closedness of the forgetful functor.

Day and the third author [12] defined a concept of quantum category in terms of monoidal comonads. They then defined a quantum groupoid to be a quantum category equipped in a certain way with star-autonomous structure. Pastro and the third author [5] characterized when star-autonomous structure could be lifted to monoidal categories of Eilenberg-Moore coalgebras for a monoidal comonad.

The work of [1] immediately suggested the possibility of strengthening the quantum groupoid definition to a property of quantum category rather than extra structure. To prepare for this, in the present paper, we generalize the work of [1] to the level of pseudomonoids in a monoidal bicategory. To suit our purpose we work with monoidal comonads on the pseudomonoids rather than opmonoidal monoids. The effect is that we need to take the concept of “coclosed” seriously since it is this structure that we wish to lift to the Eilenberg-Moore coalgebra pseudomonoid of the comonad.

We use the term “monoidale” rather than “pseudomonoid” or “monoidal object”. After two sections establishing the context, in Section 4 we study the Hopf condition on monoidal comonads on monoidales in an autonomous monoidal bicategory. The revised definition of quantum groupoid in a reasonably general braided monoidal category appears in Section 5.

2. The bicategory of comonads

Let \mathcal{M} denote a bicategory. We often ignore the associativity and unitivity constraints, writing as if \mathcal{M} were a 2-category.

Essentially following [6] and [7], we consider a bicategory $\text{Comnd}\mathcal{M}$ whose objects are pairs (A, g) where $g = (g : A \rightarrow A, \delta : g \Rightarrow gg, \varepsilon : g \Rightarrow 1_A)$ is a comonad on the object A of \mathcal{M} . A morphism $(k, \kappa) : (A, g) \rightarrow (A', g')$ in $\text{Comnd}\mathcal{M}$ consists of a morphism $k : A \rightarrow A'$ and 2-cell $\kappa : kg \Rightarrow g'k$ in \mathcal{M} satisfying the two conditions

$$\begin{aligned} \left(kg \xrightarrow{\kappa} g'k \xrightarrow{\delta k} g'g'k \right) &= \left(kg \xrightarrow{k\delta} kgg \xrightarrow{\kappa g} g'kg \xrightarrow{g'\kappa} g'g'k \right), \\ \left(kg \xrightarrow{\kappa} g'k \xrightarrow{\varepsilon k} k \right) &= \left(kg \xrightarrow{k\varepsilon} k \right). \end{aligned}$$

A 2-cell $\sigma : (k, \kappa) \Rightarrow (l, \lambda) : (A, g) \rightarrow (A', g')$ in $\text{Comnd}\mathcal{M}$ is a 2-cell $\sigma : k \Rightarrow l : A \rightarrow A'$ satisfying the condition

$$\left(kg \xrightarrow{\kappa} g'k \xrightarrow{g'\sigma} g'l \right) = \left(kg \xrightarrow{\sigma g} lg \xrightarrow{\lambda} g'l \right).$$

Here is an example of “doctrinal adjunction” [8] based on Theorem 1 of [9].

2.1. PROPOSITION. *A morphism $(k, \kappa) : (A, g) \rightarrow (A', g')$ in $\text{Comnd}\mathcal{M}$ has a left adjoint if and only if k has a left adjoint and κ is invertible in \mathcal{M} .*

PROOF. Pseudofunctors, in particular the forgetful $\text{Comnd}\mathcal{M} \rightarrow \mathcal{M}$, preserve adjunction so k has a left adjoint if (k, κ) does. The structure making this left adjoint into a morphism in $\text{Comnd}\mathcal{M}$ is the mate of the inverse of κ . ■

Recall [7] that \mathcal{M} admits the Eilenberg-Moore construction for comonads when the pseudofunctor $\mathcal{M} \rightarrow \text{Comnd}\mathcal{M}$, taking X to $(X, 1_X)$, has a right biadjoint $\text{Comnd}\mathcal{M} \rightarrow \mathcal{M}$ whose value at (A, g) is denoted by A^g . There is a pseudonatural equivalence

$$\mathcal{M}(X, A^g) \simeq \mathcal{M}(X, A)^{\mathcal{M}(X, g)}$$

where the right side is the category of Eilenberg-Moore coalgebras for the comonad $\mathcal{M}(X, g)$ on the category $\mathcal{M}(X, A)$; equally, the right side is the category

$$(\text{Comnd}\mathcal{M})((X, 1_X), (A, g))$$

whose objects we call *g-coalgebras*. Taking $X = A^g$ and evaluating the pseudonatural equivalence at the identity, we obtain the *universal g-coalgebra* $(u, \gamma) : (A^g, 1_{A^g}) \rightarrow (A, g)$ and remind the reader that this $u : A^g \rightarrow A$ has a right adjoint u^* in \mathcal{M} .

2.2. COROLLARY. *If the morphism $(k, \kappa) : (A, g) \rightarrow (A', g')$ has a left adjoint in $\text{Comnd}\mathcal{M}$ then the morphism $k^\kappa : A^g \rightarrow A'^{g'}$ has a left adjoint in \mathcal{M} , assuming the Eilenberg-Moore constructions exist.* ■

3. The bicategory of monoidales

Let \mathcal{M} denote a monoidal bicategory [10]. We often ignore the various constraint equivalences, writing as if \mathcal{M} were a Gray monoid; this is justified by [11].

A *monoidale* E in \mathcal{M} is what has previously been called a “pseudomonoid” or “monoidal object” (for example, see [12, 13]). It consists of an object E , morphisms $p : E \otimes E \rightarrow E$ and $j : I \rightarrow E$, and invertible 2-cells $\alpha : p(p \otimes 1_E) \Rightarrow p(1_E \otimes p)$, $\lambda : p(j \otimes 1_E) \Rightarrow 1_E$ and $\rho : p(1_E \otimes j) \Rightarrow 1_E$ satisfying two axioms. For example, a monoidale in the cartesian monoidal bicategory Cat is a monoidal category.

For monoidales E and F , a *monoidal morphism* $f = (f, \varphi, \varphi_0) : E \rightarrow F$ consists of a morphism $f : E \rightarrow F$ and 2-cells $\varphi : p(f \otimes f) \Rightarrow fp$, $\varphi_0 : j \Rightarrow fj$ in \mathcal{M} satisfying three axioms. For example, a monoidal morphism $f : 1 \rightarrow F$ in Cat is a monoid in the monoidal category F .

The composite of monoidal morphisms $f = (f, \varphi, \varphi_0) : E \rightarrow F$ and $g = (g, \varphi, \varphi_0) : F \rightarrow G$ is defined to be $gf = (gf, \varphi, \varphi_0) : E \rightarrow F$ where

$$\varphi = \left(p(gf \otimes gf) \cong p(g \otimes g)(f \otimes f) \xrightarrow{\varphi(f \otimes f)} gp(f \otimes f) \xrightarrow{g\varphi} gfp \right)$$

and $\varphi_0 = \left(j \xrightarrow{j\varphi_0} gj \xrightarrow{g\varphi_0} gfj \right).$

A monoidal 2-cell $\sigma : f \Rightarrow f' : E \rightarrow F$ is a 2-cell in \mathcal{M} satisfying $(\sigma p)\varphi = \varphi(p(\sigma \otimes \sigma))$ and $(\sigma j)\varphi_0 = \varphi_0$.

With obvious 2-cell compositions, this defines a bicategory $\text{Mon}\mathcal{M}$ of monoidales in \mathcal{M} .

A monoidale in \mathcal{M}^{co} is a monoidale in \mathcal{M} since the 2-cells α , λ and ρ are replaceable by their inverses. However, an *opmonoidal morphism* $w = (w, \psi, \psi_0) : E \rightarrow F$ in \mathcal{M} is a monoidal morphism in \mathcal{M}^{co} ; so we have a morphism $w : E \rightarrow F$ and 2-cells $\psi : wp \Rightarrow p(w \otimes w)$, $\psi_0 : wj \Rightarrow j$ in \mathcal{M} satisfying three axioms.

A monoidal morphism $f = (f, \varphi, \varphi_0) : E \rightarrow F$ is called *strong* when φ_2 and φ_0 are invertible. Notice that a strong monoidal morphism yields an opmorphism by taking $\psi = \varphi^{-1}$ and $\psi_0 = \varphi_0^{-1}$. For a morphism w with a right adjoint w^* , mateship provides a bijection between monoidal structures on w^* and opmonoidal structures on w . Here is another example of “doctrinal adjunction” [8].

3.1. PROPOSITION. *A monoidal morphism $f = (f, \varphi, \varphi_0) : E \rightarrow F$ has a right adjoint in $\text{Mon}\mathcal{M}$ if and only if it is strong and $f : E \rightarrow F$ has a right adjoint in \mathcal{M} . ■*

Suppose \mathcal{M} is right autonomous: for each object A there is an object A° and an equivalence

$$\mathcal{M}(A \otimes B, C) \simeq \mathcal{M}(B, A^\circ \otimes C), \quad h \mapsto \hat{h}$$

pseudonatural in B and C . It follows that the object assignment $A \mapsto A^\circ$ becomes a pseudofunctor $(-)^\circ : \mathcal{M}^{\text{op}} \rightarrow \mathcal{M}$ and there are morphisms $e : A \otimes A^\circ \rightarrow I$ and

$n : I \rightarrow A^\circ \otimes A$, extraordinary pseudonatural, and so on (for example, see [10]). It also follows that we have a pseudonatural equivalence

$$\mathcal{M}(B \otimes A^\circ, C) \simeq \mathcal{M}(B, C \otimes A).$$

A monoidale E is called *right coclosed* when the composite

$$\hat{p} = \left(E \xrightarrow{n \otimes 1} E^\circ \otimes E \otimes E \xrightarrow{1 \otimes p} E^\circ \otimes E \right)$$

has a left adjoint.

3.2. EXAMPLE. Let \mathcal{V} be a complete cocomplete symmetric monoidal closed category. Let $\mathcal{V}\text{-Mod}$ denote the autonomous monoidal bicategory whose objects are \mathcal{V} -categories and whose hom categories are defined by

$$\mathcal{V}\text{-Mod}(A, B) = [B^{\text{op}} \otimes A, \mathcal{V}]$$

in the \mathcal{V} -functor-category notation of [14]. We call the objects of $\mathcal{V}\text{-Mod}(A, B)$ *modules* from the \mathcal{V} -category A to the \mathcal{V} -category B (they are also called “bimodules”, “profunctors” and “distributors”). Each \mathcal{V} -functor $t : A \rightarrow B$ determines a module $t_* : A \rightarrow B$ defined by $t_*(b, a) = B(b, t(a))$. In fact, t_* has a right adjoint t^* defined by $t^*(a, b) = B(t(a), b)$. Moreover, autonomy of $\mathcal{V}\text{-Mod}$ is clear by taking $E^\circ = E^{\text{op}}$. Suppose C is a right-closed monoidal \mathcal{V} -category; so we have

$$C(x \otimes y, z) \cong C(y, z^x).$$

We can regard the right internal hom as a \mathcal{V} -functor $h : C \otimes C^{\text{op}} \rightarrow C^{\text{op}}$, $(x, z) \mapsto z^x$. Then our example is that $E = C^{\text{op}}$ becomes a coclosed monoidale in $\mathcal{V}\text{-Mod}$ with the module $p : E \otimes E \rightarrow E$ defined by

$$p(z, x, y) = E(z, x \otimes y) = C(x \otimes y, z).$$

This is because $\hat{p}(x, z, y) = p(z, x, y) \cong E(z^x, y) = h^*(x, z, y)$, so that $\hat{p} = h^*$ has the left adjoint h_* .

Suppose A and E are monoidales in \mathcal{M} . Dualizing a concept from Section 9 of [12] we say that an opmonoidal morphism $w : A \rightarrow E$, with a right adjoint w^* in \mathcal{M} , is *strong right coclosed* when $\psi : wp \Rightarrow p(w \otimes w)$ has an invertible mate:

$$\psi^r : p(1 \otimes w^*) \Rightarrow w^*p(w \otimes 1).$$

4. Monoidal comonads

When \mathcal{M} is a monoidal bicategory, so too is $\text{Comnd}\mathcal{M}$ where

$$(A, g) \otimes (B, h) = (A \otimes B, g \otimes h).$$

A *monoidal comonad* in \mathcal{M} is a comonad $g = (g, \varphi, \varphi_0) : E \rightarrow E$ in the bicategory $\text{Mon}\mathcal{M}$. Equally, it is a monoidale in the monoidal bicategory $\text{Comnd}\mathcal{M}$. It consists of a monoidale E , a comonad g on E , and 2-cells $\varphi : p(g \otimes g) \Rightarrow gp$, $\varphi_0 : j \Rightarrow gj$ such that $(p, \varphi) : (E \otimes E, g \otimes g) \rightarrow (E, g)$ and $(j, \varphi_0) : (I, 1) \rightarrow (E, g)$ are morphisms in $\text{Comnd}\mathcal{M}$.

If \mathcal{M} admits the Eilenberg-Moore construction for comonads then so too does $\text{Mon}\mathcal{M}$; see [15] and [16]. We now generalize a definition due to Bruguières-Lack-Virelizier announced in [1]. All of Proposition 4.2, Theorem 4.3 and Proposition 4.4 below are generalizations to our setting of the results forming part of [1, 2].

4.1. DEFINITION. A *monoidal comonad* g on a monoidale E is *right Hopf* when the pasted 2-cell 4.1 is invertible.

$$\begin{array}{ccccc}
 E \otimes E & \xrightarrow{g \otimes 1} & E \otimes E & \xrightarrow{p} & E \\
 1 \otimes g \downarrow & \xrightarrow{\delta \otimes 1 g} & g \otimes g \downarrow & \xrightarrow{\varphi} & \downarrow g \\
 E \otimes E & \xrightarrow{g \otimes 1} & E \otimes E & \xrightarrow{p} & E
 \end{array} \tag{4.1}$$

We henceforth assume that the monoidal bicategory \mathcal{M} is right autonomous and admits the Eilenberg-Moore construction for comonads.

4.2. PROPOSITION. A *monoidal comonad* g on E is *right Hopf* if and only if the 2-cell 4.2 is invertible.

$$\begin{array}{ccccc}
 E^g \otimes E & \xrightarrow{u \otimes 1} & E \otimes E & \xrightarrow{p} & E \\
 1 \otimes g \downarrow & \xrightarrow{\gamma \otimes 1 g} & g \otimes g \downarrow & \xrightarrow{\varphi} & \downarrow g \\
 E^g \otimes E & \xrightarrow{u \otimes 1} & E \otimes E & \xrightarrow{p} & E
 \end{array} \tag{4.2}$$

PROOF. Consider the following 3×3 diagram.

$$\begin{array}{ccccc}
 u \otimes g & \xrightarrow{\gamma \otimes 1} & gu \otimes g & \xrightarrow{\varphi(u \otimes 1)} & g(u \otimes 1) \\
 \gamma \otimes 1 \downarrow & & g\gamma \otimes 1 \downarrow & & g\gamma \otimes 1 \downarrow \\
 gu \otimes g & \xrightarrow{\delta u \otimes 1} & g^2u \otimes g & \xrightarrow{\varphi(gu \otimes 1)} & g(gu \otimes 1) \\
 g\gamma \otimes 1 \downarrow & \delta u \otimes 1 \downarrow & g^2\gamma \otimes 1 \downarrow & g\delta u \otimes 1 \downarrow & g(g\gamma \otimes 1) \downarrow \\
 & & & & g(\delta u \otimes 1) \downarrow \\
 g^2u \otimes g & \xrightarrow{\delta gu \otimes 1} & g^3u \otimes g & \xrightarrow{\varphi(g^2u \otimes 1)} & g(g^2u \otimes 1)
 \end{array} \tag{4.3}$$

Since the pseudofunctor $- \otimes E : \mathcal{M} \rightarrow \mathcal{M}$ has a left biadjoint $- \otimes E^\circ$, it preserves Eilenberg-Moore constructions (as they are weighted limits). It follows that $u \otimes E : E^g \otimes E \rightarrow E \otimes E$ is comonadic and then that the columns of 4.3 are absolute equalizers. If 4.1 is invertible then the composites in the second and third row of 4.3 are invertible. Therefore the composite in the first row is invertible yielding the invertibility of 4.2. For the converse, note that precomposing 4.2 with $u^* \otimes E$ yields 4.1. ■

4.3. THEOREM. *Suppose E is a right coclosed monoidal in \mathcal{M} . A monoidal comonad g on E is right Hopf if and only if the monoidal E^g is right coclosed and $u : E^g \rightarrow E$ is strong right coclosed.*

PROOF. Suppose g is a Hopf monoidal comonad on E . Consider the following diagram.

$$\begin{array}{ccccccc}
 E & \xrightarrow{n \otimes 1} & E^{g^\circ} \otimes E^g \otimes E & \xrightarrow{1 \otimes u \otimes 1} & E^{g^\circ} \otimes E \otimes E & \xrightarrow{1 \otimes p} & E^{g^\circ} \otimes E \\
 g \downarrow & \cong & 1 \otimes 1 \otimes g \downarrow & \xrightarrow{1 \otimes \gamma \otimes 1} & 1 \otimes g \otimes g \downarrow & \xrightarrow{1 \otimes \varphi} & \downarrow 1 \otimes g \\
 E & \xrightarrow{n \otimes 1} & E^{g^\circ} \otimes E^g \otimes E & \xrightarrow{1 \otimes u \otimes 1} & E^{g^\circ} \otimes E \otimes E & \xrightarrow{1 \otimes p} & E^{g^\circ} \otimes E
 \end{array} \tag{4.4}$$

The pasted 2-cell is invertible by Proposition 4.2 while the top and bottom composites are isomorphic to the composite

$$E \xrightarrow{p} E^\circ \otimes E \xrightarrow{u^\circ \otimes 1} E^{g^\circ} \otimes E. \tag{4.5}$$

Now \hat{p} has a left adjoint since E is right coclosed and $u^\circ \otimes 1$ has the left adjoint $u^{\circ*} \otimes 1$. By Corollary 2.2 the induced morphism $q : E^g \rightarrow E^{g^\circ} \otimes E^g$ on the Eilenberg-Moore constructions of the comonads at the left and right ends of diagram 4.4 has a left adjoint. In fact, $q = \hat{p}$ where this p is that of the monoidal E^g . To see this we must see that pasting the triangle containing the universal g -coalgebra γ to the left of diagram 4.4 conjugated by the isomorphisms of the top and bottom with the morphism 4.5, is equal to \hat{p} pasted on the right with the triangle for the universal $(E^{g^\circ} \otimes g)$ -coalgebra $E^{g^\circ} \otimes \gamma$. The morphism part of this calculation is:

$$\begin{aligned}
 (E^{g^\circ} \otimes u) \hat{p} &= (E^{g^\circ} \otimes u) (E^{g^\circ} \otimes p) (n \otimes E^g) \\
 &\cong (E^{g^\circ} \otimes p) (u \otimes u) (n \otimes E^g) \cong (E^{g^\circ} \otimes p) (u^\circ \otimes E \otimes E) (n \otimes E) u \\
 &\cong (u^\circ \otimes E) (E^\circ \otimes p) (n \otimes E) u \cong (u^\circ \otimes E) \hat{p} u.
 \end{aligned}$$

For the 2-cell part of the calculation notice that monoidalness of γ implies the equality of the composite 2-cells

$$\begin{aligned}
 &(1 \otimes p)(1 \otimes u \otimes u)(n \otimes 1) \\
 &\quad \downarrow (1 \otimes p)(1 \otimes \gamma \otimes \gamma)(n \otimes 1) \\
 (1 \otimes p)(1 \otimes gu \otimes gu)(n \otimes 1) &= (1 \otimes p)(1 \otimes g \otimes g)(1 \otimes u \otimes u)(n \otimes 1) \\
 &\quad \downarrow \psi(1 \otimes u \otimes u)(n \otimes 1) \\
 (1 \otimes g)(1 \otimes u)(1 \otimes p)(n \otimes 1) &\cong (1 \otimes g)(1 \otimes p)(1 \otimes u \otimes u)(n \otimes 1)
 \end{aligned}$$

and

$$\begin{aligned}
 (1 \otimes p)(1 \otimes u \otimes u)(n \otimes 1) &\cong (1 \otimes u)(1 \otimes p)(n \otimes 1) \\
 &\quad \downarrow \gamma(1 \otimes p)(n \otimes 1) \\
 (1 \otimes g)(1 \otimes u)(1 \otimes p)(n \otimes 1) &= (1 \otimes gu)(1 \otimes p)(n \otimes 1)
 \end{aligned}$$

Hence it follows that E^g is right coclosed.

We need to see that $u : E^g \rightarrow E$ is strong right coclosed. That is, we need to show $p(1 \otimes u^*) \Rightarrow u^*p(u \otimes 1)$ is invertible. As the unit for the adjunction $u \dashv u^*$ is an equalizer, composition with u is conservative (invertibility reflecting). So it suffices to show that $up(1 \otimes u^*) \Rightarrow uu^*p(u \otimes 1)$ is invertible. But, using the invertibility of 4.2, we have

$$up(1 \otimes u^*) \cong p(u \otimes u)(1 \otimes u^*) \cong p(u \otimes g) \cong gp(u \otimes 1) \cong uu^*p(u \otimes 1).$$

The converse follows from the more general fact expressed in Proposition 4.4. ■

4.4. PROPOSITION. *If $w : A \rightarrow E$ is a strong right coclosed, strong monoidal morphism with a right adjoint w^* in \mathcal{M} then the generated monoidal comonad $g = ww^*$ on E is Hopf.*

PROOF. We have

$$p(g \otimes 1)(1 \otimes g) \cong p(g \otimes g) \cong p(w \otimes w)(w^* \otimes w^*)$$

and, using the fact that w is strong monoidal, we continue with

$$\cong wp(w^* \otimes w^*) \cong wp(1 \otimes w^*)(w^* \otimes 1)$$

and, using the fact that w is strong right closed, we continue with

$$\cong ww^*p(w \otimes 1)(w^* \otimes 1) \cong ww^*p(ww^* \otimes 1) \cong gp(g \otimes 1).$$

■

4.5. EXAMPLE. In the setting of Example 3.2, suppose $t = (t, \mu, \eta, \psi, \psi_0)$ is an opmonoidal monad on a monoidal \mathcal{V} -category C . We may consider t equally to be a monoidal comonad, which we will write as $g = (g, \delta, \varepsilon, \varphi, \varphi_0)$ to mark the distinction, on $E = C^{\text{op}}$. Then t is a Hopf opmonoidal monad on C in the sense of [1] (actually, they say more briefly that t is a ‘‘Hopf monad’’) if and only if g_* is a Hopf monoidal comonad on $E = C^{\text{op}}$ in $\mathcal{V}\text{-Mod}$. To be explicit, the right Hopf condition is, for all objects x and y in C , the invertibility of the composite

$$t(t(x) \otimes y) \xrightarrow{\psi_{t(x),y}} t^2(x) \otimes t(y) \xrightarrow{\mu_x \otimes 1_{t(x)}} t(x) \otimes t(y). \tag{4.6}$$

It should be noted that, if C^t denotes the \mathcal{V} -category of Eilenberg-Moore t -algebras, then $E^g = (C^t)^{\text{op}}$ in $\mathcal{V}\text{-Mod}$. By Proposition 4.2, the invertibility of the arrows 4.6 is equivalent to the invertibility of the arrows

$$t(a \otimes y) \xrightarrow{\psi_{a,y}} t(a) \otimes t(y) \xrightarrow{\alpha \otimes 1} a \otimes t(y) \tag{4.7}$$

for all t -algebras (a, α) and all objects y . If C is right closed, it follows that, for all t -algebras (a, α) and (b, β) , there is a unique t -action $\rho : t(b^a) \rightarrow b^a$ on b^a such that

$$\left(t(a \otimes b^a) \xrightarrow{t(\text{ev})} t(b) \xrightarrow{\beta} b \right) = \left(t(a \otimes b^a) \xrightarrow{t(\text{ev})} t(a) \otimes t(b^a) \longrightarrow a \otimes b^a \xrightarrow{\text{ev}} b \right).$$

As pointed out in [2], this means that $\text{ev} : a \otimes b^a \rightarrow b$ becomes a t -algebra morphism. Hence (b^a, ρ) is a right internal hom of (a, α) and (b, β) in C^t , all in accord with our Theorem 4.3.

5. Quantum groupoids

Let \mathcal{V} be a braided monoidal category with coreflexive equalizers preserved by the functors $X \otimes - : \mathcal{V} \rightarrow \mathcal{V}$. Following [17], we consider the right autonomous monoidal bicategory $\text{Comod}(\mathcal{V})$ whose objects are comonoids $C = (C, \delta)$ (sometimes called “coalgebras”) in \mathcal{V} and whose hom $\text{Comod}(\mathcal{V})(C, D)$ is the Eilenberg-Moore category for the comonad $C \otimes - \otimes D$ on \mathcal{V} . Eilenberg-Moore coalgebras M for $C \otimes - \otimes D$ are called *comodules from C to D* and depicted by $M : C \rightarrow D$. The composite $N \circ M = M \diamond_D N$ of comodules $M : C \rightarrow D$ and $N : D \rightarrow E$ is defined as a coreflexive equalizer

$$M \diamond_D N \longrightarrow M \otimes N \begin{array}{c} \longrightarrow \\ \longrightarrow \end{array} M \otimes D \otimes N.$$

The monoidal structure on $\text{Comod}(\mathcal{V})$ simply extends the tensor product on \mathcal{V} . The right dual C° of the comonoid C has comultiplication defined using the braiding thus:

$$\left(C^\circ \xrightarrow{\delta} C^\circ \otimes C^\circ \right) = \left(C \xrightarrow{\delta} C \otimes C \xrightarrow{c_{C,C}} C \otimes C \right).$$

It follows that the biduality $C \dashv_b C^\circ$ generates a monoidal structure $p = 1 \otimes e \otimes 1$, $j = n$ on $C^\circ \otimes C$ in $\text{Comod}(\mathcal{V})$.

Recall that a *quantum category* $A = (C, G)$ in \mathcal{V} is defined in [12] to be a comonoid C in \mathcal{V} equipped with a monoidal comonad G on the monoidal $C^\circ \otimes C$ in $\text{Comod}(\mathcal{V})$. The authors go on to add conditions, involving star-autonomy, in order for a quantum category to be a quantum groupoid. Now we shall suggest a more general definition which is in the spirit of [10], [18] and [19]. A virtue is that it is a property, rather than extra structure, that a quantum category may have.

5.1. DEFINITION. *A quantum groupoid in \mathcal{V} is a quantum category $A = (C, G)$ where the monoidal comonad G is Hopf.*

References

- [1] Alain Bruguières, “Hopf monads on monoidal categories, and modularity of the centre” *Australian Category Seminar* (Macquarie University, 10 December 2008); <http://www.maths.usyd.edu.au/u/AusCat/abstracts/081210ab.html>
- [2] A. Bruguières, S. Lack and A. Virelizier, Hopf monads on monoidal categories, *Advances in Mathematics*, to appear; also <http://au.arxiv.org/abs/1003.1920>.
- [3] S. Eilenberg and G. M. Kelly, Closed categories, *Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965)*, Springer-Verlag, New York, 1966, 421–562.

- [4] A. Bruguières and A. Virelizier, Hopf monads, *Advances in Mathematics* **215** (2007), 679–733.
- [5] C. Pastro and R. Street, Closed categories, star-autonomy, and monoidal comonads, *Journal of Algebra* **321** (2009), 3494–3520.
- [6] J. Bénabou, Introduction to bicategories, *Lecture Notes in Mathematics* **47** (Springer-Verlag, Berlin, 1967), 1–77.
- [7] R. Street, The formal theory of monads, *Journal of Pure and Applied Algebra* **2** (1972), 149–168.
- [8] G. M. Kelly, Doctrinal adjunction, *Category Seminar (Proc. Sem., Sydney, 1972/1973) Lecture Notes in Math.*, **420** (Springer, Berlin, 1974), 257–280.
- [9] R. Street, Two constructions on lax functors, *Cahiers topologie et géométrie différentielle* **13** (1972), 217–264.
- [10] B. Day and R. Street, Monoidal bicategories and Hopf algebroids, *Advances in Mathematics* **129** (1997), 99–157.
- [11] R. Gordon, A. J. Power and R. Street, *Coherence for tricategories*, *Memoirs of the American Math. Society*, Volume 117, Number 558, 1995.
- [12] B. Day and R. Street, Quantum categories, star autonomy, and quantum groupoids, *Fields Institute Communications (American Math. Soc.)* **43** (2004), 187–226.
- [13] I. Lopez-Franco, R. Street and R. J. Wood, Duals invert, *Applied Categorical Structures*, to appear; published online 26 September 2009 at <http://www.springerlink.com/content/u517174h46wm2313/>
- [14] G. M. Kelly, *Basic concepts of enriched category theory*, London Mathematical Society Lecture Note Series, 64. Cambridge University Press, Cambridge, 1982. <http://www.tac.mta.ca/tac/reprints/articles/10/tr10.pdf>
- [15] I. Moerdijk, Monads on tensor categories, *Journal of Pure Appl. Algebra* **168** (2002), 189–208.
- [16] P. McCrudden, Opmonoidal monads, *Theory and Applications of Categories* **10** (2002), 469–485. <http://www.tac.mta.ca/tac/volumes/10/19/10-19.pdf>
- [17] B. Day, P. McCrudden and R. Street, Dualizations and antipodes, *Applied Categorical Structures* **11** (2003), 229–260.
- [18] R. Street, Fusion operators and cocycloids in monoidal categories, *Applied Categorical Structures* **6** (1998), 177–191.
- [19] P. Schauenburg, Duals and doubles of quantum groupoids (\times $_R$ -Hopf algebras), *Contemporary Mathematics (Amer. Math. Soc., Providence, RI)* **267** (2000), 273–299.

Department of Mathematics, Macquarie University, NSW 2109, Australia

Email: d.chikhladze@gmail.com, steve.lack@mq.edu.au, ross.street@mq.edu.au

This article may be accessed at <http://www.tac.mta.ca/tac/> or by anonymous ftp at [{dvi,ps,pdf}](ftp://ftp.tac.mta.ca/pub/tac/html/volumes/24/19/24-19)

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION. Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS. The typesetting language of the journal is \TeX , and $\text{\LaTeX}2\epsilon$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

\TeX NICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT \TeX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis, cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Aurelio Carboni, Università dell' Insubria: aurelio.carboni@uninsubria.it

Valeria de Paiva: valeria.depaiiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Martin Hyland, University of Cambridge: M.Hyland@dpms.cam.ac.uk

P. T. Johnstone, University of Cambridge: ptj@dpms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Glasgow, Tom.Leinster@glasgow.ac.uk

Jean-Louis Loday, Université de Strasbourg: loday@math.u-strasbg.fr

Ieke Moerdijk, University of Utrecht: moerdijk@math.uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Brooke Shipley, University of Illinois at Chicago: bshipley@math.uic.edu

James Stasheff, University of North Carolina: jds@math.unc.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca