

COMPLETION, CLOSURE, AND DENSITY RELATIVE TO A MONAD, WITH EXAMPLES IN FUNCTIONAL ANALYSIS AND SHEAF THEORY

RORY B. B. LUCYSHYN-WRIGHT

ABSTRACT. Given a monad \mathbb{T} on a suitable enriched category \mathcal{B} equipped with a proper factorization system $(\mathcal{E}, \mathcal{M})$, we define notions of \mathbb{T} -completion, \mathbb{T} -closure, and \mathbb{T} -density. We show that not only the familiar notions of completion, closure, and density in normed vector spaces, but also the notions of sheafification, closure, and density with respect to a Lawvere-Tierney topology, are instances of the given abstract notions. The process of \mathbb{T} -completion is equally the enriched idempotent monad associated to \mathbb{T} (which we call the *idempotent core of \mathbb{T}*), and we show that it exists as soon as every morphism in \mathcal{B} factors as a \mathbb{T} -dense morphism followed by a \mathbb{T} -closed \mathcal{M} -embedding. The latter hypothesis is satisfied as soon as \mathcal{B} has certain pullbacks as well as wide intersections of \mathcal{M} -embeddings. Hence the resulting theorem on the existence of the idempotent core of an enriched monad entails Fakir’s existence result in the non-enriched case, as well as adjoint functor factorization results of Applegate-Tierney and Day.

1. Introduction

Examples of monads abound throughout mathematics, particularly since every adjunction determines one, yet monads play also a seemingly more narrow role as theories of algebraic structure, each monad \mathbb{T} on a category \mathcal{B} determining a category $\mathcal{B}^{\mathbb{T}}$ of \mathbb{T} -algebras. Working in the context of the theory of categories enriched over a symmetric monoidal closed category \mathcal{V} , we show herein that, on the other hand, every monad \mathbb{T} on any suitable category \mathcal{B} also gives rise to concepts that are seemingly more ‘topological’ in nature, namely, canonical notions of *closure*, *density*, *completeness*, *completion*, and *separatedness* with respect to \mathbb{T} . As a guiding example, we show that when \mathcal{B} is the category of normed or semi-normed vector spaces and \mathbb{T} is the monad given by taking the double-dual, the resulting notions with respect to \mathbb{T} coincide with the familiar notions. As an example of a very different sort, we show that when \mathcal{B} is an (elementary) topos equipped with a

The author gratefully acknowledges financial support in the form of an NSERC Postdoctoral Fellowship, as well as earlier, partial financial support in the form of an Ontario Graduate Scholarship.

Received by the editors 2014-06-09 and, in revised form, 2014-12-16.

Transmitted by F. W. Lawvere. Published on 2014-12-19.

2010 Mathematics Subject Classification: 18A20, 18A22, 18A30, 18A32, 18A40, 18B25, 18C15, 18D15, 18D20, 18F10, 18F20, 46B04, 46B10, 46B28, 46M99.

Key words and phrases: completion; closure; density; monad; idempotent monad; idempotent core; idempotent approximation; normed vector space; adjunction; reflective subcategory; enriched category; factorization system; orthogonal subcategory; sheaf; sheafification; Lawvere-Tierney topology; monoidal category; closed category.

© Rory B. B. Lucyshyn-Wright, 2014. Permission to copy for private use granted.

Lawvere-Tierney topology j and we take \mathbb{T} to be the double-dualization monad for Ω_j , we recover the notions of j -closure, j -density, j -sheaf, j -sheafification, and j -separatedness.

In more detail, let \mathbb{T} be a monad on a \mathcal{V} -category \mathcal{B} equipped with a proper \mathcal{V} -enriched factorization system $(\mathcal{E}, \mathcal{M})$ ([21]), and refer to the morphisms in \mathcal{M} as *embeddings*. Using techniques of enriched factorization systems and orthogonality, we define notions of \mathbb{T} -dense morphism, \mathbb{T} -closed embedding, and \mathbb{T} -complete object. Next, we make the following assumption, which is satisfied as soon as \mathcal{B} is complete and well-powered with respect to \mathcal{M} :

$$\begin{aligned} \text{Every morphism in } \mathcal{B} \text{ factors as a } \mathbb{T}\text{-dense} \\ \text{morphism followed by a } \mathbb{T}\text{-closed embedding.} \end{aligned} \tag{1.0.i}$$

Under this assumption, the \mathbb{T} -dense morphisms and \mathbb{T} -closed embeddings constitute an enriched factorization system, and we obtain an operation of \mathbb{T} -closure of \mathcal{M} -subobjects. We then show that the full subcategory $\mathcal{B}_{(\mathbb{T})}$ of \mathcal{B} consisting of the \mathbb{T} -complete objects is reflective in \mathcal{B} , and the resulting idempotent monad $\tilde{\mathbb{T}}$ on \mathcal{B} we call the \mathbb{T} -completion monad. The \mathbb{T} -completion $\tilde{T}B$ of an object $B \in \mathcal{B}$ is gotten as in the example of normed vector spaces: We take $\tilde{T}B$ to be the \mathbb{T} -closure of the $(\mathcal{E}, \mathcal{M})$ -image of the unit morphism $\eta_B : B \rightarrow TB$.

The resulting \mathbb{T} -completion monad $\tilde{\mathbb{T}}$ is an idempotent \mathcal{V} -enriched monad on \mathcal{B} that inverts (i.e., sends to isomorphisms) exactly the same morphisms as \mathbb{T} and so is *the idempotent core*¹ of \mathbb{T} , studied in the context of non-enriched categories by Casacuberta and Frei [3] (under the name of the *idempotent approximation* of \mathbb{T}), by Hébert [14, Proposition 14], and earlier by Fakir [11], who had shown that the idempotent core of a monad on an ordinary category \mathcal{B} exists as soon as \mathcal{B} is complete and well-powered. Our construction of the \mathbb{T} -completion monad shows that the \mathcal{V} -enriched idempotent core $\tilde{\mathbb{T}}$ of \mathbb{T} exists as soon as the factorization assumption (1.0.i) is satisfied, and Fakir’s result is recovered as a corollary. Note that our result applies even in the absence of set-indexed limits and so applies, for example, in constructing the j -sheafification monad for a Lawvere-Tierney topology j on an arbitrary (elementary) topos.

After recalling some preliminary material on monads and adjunctions in 2-categories (§2.1), enriched categories (§2.2), enriched factorization systems (§2.3), and closure operators (§2.4), we treat aspects of enriched orthogonal subcategories needed in the sequel (§3). Next, we treat the basic theory of the idempotent core of an enriched monad (§4), as no such treatment exists in the literature; in particular, we consider the universal property of the idempotent core $\tilde{\mathbb{T}}$, we establish several equivalent characterizations of $\tilde{\mathbb{T}}$ and of its existence, and we examine the relation of the enriched idempotent core to the ordinary. Next we show that the completion monad on normed (resp. seminormed) vector spaces is the enriched idempotent core of the double-dualization monad (§5); as a corollary, we show that the full subcategory consisting of all Banach spaces is the enriched reflective hull of the space of scalars (\mathbb{R} or \mathbb{C}). In §6 we then define the notions of \mathbb{T} -density, \mathbb{T} -closure, etc., and we prove our general result on the existence of the idempotent \mathcal{V} -core $\tilde{\mathbb{T}}$,

¹Terminology suggested by F. W. Lawvere.

Theorem 6.25. In §7 we return to the example of normed and seminormed vector spaces and show that the notions of \mathbb{T} -density and \mathbb{T} -closure there coincide with the familiar notions. In §8 we treat the example of j -sheafification, j -closure, and j -density for a Lawvere-Tierney topology j on a topos \mathcal{X} , and we show that the category of j -sheaves is the \mathcal{X} -enriched reflective hull of Ω_j in \mathcal{X} ; note that Lambek and Rattray had shown in [19] that the j -sheafification monad can be obtained by Fakir’s construction.

In treating the basic theory of the idempotent core $\tilde{\mathbb{T}}$ of an enriched monad \mathbb{T} on a \mathcal{V} -category \mathcal{B} , we show in 4.15 that if $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ is an arbitrary \mathcal{V} -adjunction inducing \mathbb{T} , then $\tilde{\mathbb{T}}$ exists if and only if $F \dashv G$ factors as a composite \mathcal{V} -adjunction

$$\mathcal{B} \begin{array}{c} \xleftarrow{J} \\ \xrightarrow{\mathbb{T}} \\ \xrightarrow{K} \end{array} \mathcal{B}' \begin{array}{c} \xleftarrow{G'} \\ \xrightarrow{\mathbb{T}} \\ \xrightarrow{F'} \end{array} \mathcal{C}$$

with J a \mathcal{V} -reflective full subcategory inclusion and F' conservative. Hence our Theorem 6.25 on the existence of $\tilde{\mathbb{T}}$ yields a generalization on results of Applegate and Tierney [1] and Day [6] to the effect that any adjunction $F \dashv G$ on a suitable category factors in such a way. An important point that was not made by these authors is that the resulting reflective subcategory \mathcal{B}' depends only on the *monad* \mathbb{T} induced by $F \dashv G$. Cassidy, Hébert, and Kelly later proved in the non-enriched context a variant of Day’s adjoint-factorization result ([5, Theorem 3.3]), and in the second paragraph of their proof they make use of an instance of what we now call here the (\mathbb{T} -dense, \mathbb{T} -closed embedding)-factorization system, there written as $(\mathcal{N}^\uparrow, \mathcal{N})$.

We consider also the following refinement of the theory of \mathbb{T} -completeness, \mathbb{T} -density, etc.: Given an enriched monad \mathbb{T} on a \mathcal{V} -category \mathcal{B} equipped with a proper \mathcal{V} -prefactorization-system $(\mathcal{E}, \mathcal{M})$ ([21]), together with a given class Σ consisting of morphisms inverted by T , we define notions of Σ -dense morphism, Σ -closed embedding, and Σ -complete object, and again assuming that every morphism in \mathcal{B} factors as a Σ -dense morphism followed by a Σ -closed embedding, we show that those Σ -complete objects $B \in \mathcal{B}$ that are also \mathbb{T} -separated (meaning that $\eta_B : B \rightarrow TB$ is an embedding) constitute a \mathcal{V} -reflective subcategory $\mathcal{B}_{(\mathbb{T}, \Sigma)}$ of \mathcal{B} (6.20), so that we obtain an idempotent monad $\tilde{\mathbb{T}}_\Sigma$, the \mathbb{T} -separated Σ -completion monad. This reflectivity is related to Day’s result [6, Corollary 2.3]. The ‘ \mathbb{T} -’ rather than ‘ Σ -’ notions are recovered when $\Sigma := T^{-1}(\text{Iso})$, and in the latter case it is notable that every \mathbb{T} -complete object is necessarily \mathbb{T} -separated, provided $(\mathcal{E}, \mathcal{M})$ -factorizations exist.

The given notions of completeness, closure, and density with respect to an enriched monad \mathbb{T} and/or a class of morphisms Σ were employed in the author’s recent Ph.D. thesis [20], in which they were applied with regard to R -module objects in a cartesian closed category (and generalizations thereupon) in providing a basis for abstract functional analysis in a closed category. In such a context, the usual notions of completeness, closure, and density familiar from functional analysis are not typically available, and so one may instead employ the above ‘ \mathbb{T} -’ notions, with respect to the double-dualization monad \mathbb{T} , and these we call *functional completeness*, *functional closure*, and *functional density*.

2. Preliminaries

2.1. 2-CATEGORICAL PRELIMINARIES.

2.1.1. Given an object \mathcal{B} in a 2-category \mathcal{K} , there is a category $\text{Mnd}_{\mathcal{K}}(\mathcal{B})$ whose objects are monads on \mathcal{B} and whose morphisms $\theta : (T, \eta, \mu) \rightarrow (T', \eta', \mu')$ (called *maps of monads* in [18]) consist of a 2-cell $\theta : T \rightarrow T'$ such that $\theta \cdot \eta = \eta'$ and $\mu' \cdot (\theta \circ \theta) = \theta \cdot \mu$. The identity monad $\mathbb{1}_{\mathcal{B}}$ is an initial object in $\text{Mnd}_{\mathcal{K}}(\mathcal{B})$, since for each monad $\mathbb{T} = (T, \eta, \mu)$ on \mathcal{B} , the 2-cell η is the unique monad morphism $\eta : \mathbb{1}_{\mathcal{B}} \rightarrow \mathbb{T}$.

2.1.2. Given a monad $\mathbb{T} = (T, \eta, \mu)$ on \mathcal{B} in a 2-category \mathcal{K} , a \mathbb{T} -algebra (B, β) in \mathcal{K} ([18, 3.1]) consists of a 1-cell $B : \mathcal{A} \rightarrow \mathcal{B}$ equipped with an *action* of \mathbb{T} on B , i.e. a 2-cell $\beta : TB \rightarrow B$ with $\beta \cdot \eta B = 1_B$ and $\beta \cdot T\beta = \beta \cdot \mu B$. Given a morphism of monads $\theta : \mathbb{T} \rightarrow \mathbb{T}' = (T', \eta', \mu')$, it is shown in [18, (3.8), (3.9)] that the composite $\beta := (TT' \xrightarrow{\theta T'} T'T' \xrightarrow{\mu'} T')$ is an action of \mathbb{T} on T' and that θ can be expressed in terms of β as the composite $T \xrightarrow{T\eta'} TT' \xrightarrow{\beta} T'$.

2.1.3. Recall that a monad $\mathbb{S} = (S, \rho, \lambda)$ on \mathcal{B} in \mathcal{K} is said to be *idempotent* if $\lambda : SS \rightarrow S$ is an isomorphism (from which it then follows that $\lambda^{-1} = S\rho = \rho S$). If \mathbb{S} is idempotent, then for any \mathbb{S} -algebra (B, β) (2.1.2), the 1-cell $\beta : SB \rightarrow B$ is an isomorphism with inverse ρB .

2.1.4. Given objects \mathcal{A}, \mathcal{B} in a 2-category \mathcal{K} , there is a category $\text{Adj}_{\mathcal{K}}(\mathcal{A}, \mathcal{B})$ whose objects are adjunctions $F \frac{\eta}{\varepsilon} \Big| G : \mathcal{B} \rightarrow \mathcal{A}$ in \mathcal{K} and whose morphisms $(\phi, \psi) : (F \frac{\eta}{\varepsilon} \Big| G) \rightarrow (F' \frac{\eta'}{\varepsilon'} \Big| G')$ consist of 2-cells $\phi : F \rightarrow F'$ and $\psi : G \rightarrow G'$ such that $(\psi \circ \phi) \cdot \eta = \eta'$ and $\varepsilon' \cdot (\phi \circ \psi) = \varepsilon$.

There is a functor $\text{Adj}_{\mathcal{K}}(\mathcal{A}, \mathcal{B}) \rightarrow \text{Mnd}_{\mathcal{K}}(\mathcal{A})$ sending an adjunction to its induced monad and a morphism $(\phi, \psi) : (F \frac{\eta}{\varepsilon} \Big| G) \rightarrow (F' \frac{\eta'}{\varepsilon'} \Big| G')$ to the morphism $\psi \circ \phi : \mathbb{T} \rightarrow \mathbb{T}'$ between the induced monads. Hence, in particular, isomorphic adjunctions induce isomorphic monads.

2.1.5. PROPOSITION. *Let $F \frac{\eta}{\varepsilon} \Big| G : \mathcal{B} \rightarrow \mathcal{A}$ be an adjunction in a 2-category \mathcal{K} . Then there is an associated monoidal functor $[F, G] := \mathcal{K}(F, G) : \mathcal{K}(\mathcal{B}, \mathcal{B}) \rightarrow \mathcal{K}(\mathcal{A}, \mathcal{A})$ with the following property: For any adjunction $F' \frac{\eta'}{\varepsilon'} \Big| G' : \mathcal{C} \rightarrow \mathcal{B}$ with induced monad \mathbb{T}' on \mathcal{B} , the monad $[F, G](\mathbb{T}')$ on \mathcal{A} is equal to the monad induced by the composite adjunction*

$$\mathcal{A} \begin{array}{c} \xleftarrow{G} \\ \eta \top \varepsilon \\ \xrightarrow{F} \end{array} \mathcal{B} \begin{array}{c} \xleftarrow{G'} \\ \eta' \top \varepsilon' \\ \xrightarrow{F'} \end{array} \mathcal{C} .$$

PROOF. The monoidal structure on the functor $[F, G]$ consists of the morphisms $GH\varepsilon KF : GHFGKF \rightarrow GHKF$ in $\mathcal{K}(\mathcal{A}, \mathcal{A})$ (for all objects H, K in $\mathcal{K}(\mathcal{B}, \mathcal{B})$) and the morphism $\eta : 1_{\mathcal{A}} \rightarrow GF$ in $\mathcal{K}(\mathcal{A}, \mathcal{A})$. The verification is straightforward. ■

2.1.6. PROPOSITION. Let $F \xleftarrow[\varepsilon]{\eta} G$ and $F' \xleftarrow[\varepsilon']{\eta'} G$ be adjunctions, having the same right adjoint $G : \mathcal{B} \rightarrow \mathcal{A}$, in a 2-category \mathcal{K} . Then these adjunctions are isomorphic and hence induce isomorphic monads on \mathcal{A} .

PROOF. The 2-cell $\phi := \varepsilon F' \cdot F \eta' : F \rightarrow F'$ has inverse $\varepsilon' F \cdot F' \eta$ ([13], I,6.3), and one checks that $(\phi, 1_G)$ serves as the needed isomorphism of adjunctions. ■

2.1.7. PROPOSITION. Let $\mathcal{A} \xleftarrow[\underset{F}{\eta \top \varepsilon}]{G} \mathcal{B} \xleftarrow[\underset{F'}{\eta' \top \varepsilon'}]{G'} \mathcal{C}$ and $\mathcal{A} \xleftarrow[\underset{F''}{\eta'' \top \varepsilon''}]{G''} \mathcal{C}$ be adjunctions in a 2-category \mathcal{K} , with respective induced monads $\mathbb{T}, \mathbb{T}', \mathbb{T}''$, and suppose that $GG' = G''$. Then there is an associated monad morphism $\mathbb{T} \rightarrow \mathbb{T}''$.

PROOF. Let $F_c \xleftarrow[\varepsilon_c]{\eta_c} G''$ be the composite adjunction, and let \mathbb{T}_c be its induced monad. By 2.1.5, we have that $\mathbb{T}_c = [F, G](\mathbb{T}')$, whereas $\mathbb{T} = [F, G](1_{\mathcal{B}})$. By applying $[F, G]$ to the monad morphism $\eta' : 1_{\mathcal{B}} \rightarrow \mathbb{T}'$, we obtain a monad morphism

$$G\eta'F = [F, G](\eta') : \mathbb{T} = [F, G](1_{\mathcal{B}}) \rightarrow [F, G](\mathbb{T}') = \mathbb{T}_c .$$

Also, by 2.1.6, there is an isomorphism of monads $\xi : \mathbb{T}_c \rightarrow \mathbb{T}''$, and we obtain a composite morphism of monads $\mathbb{T} \xrightarrow{G\eta'F} \mathbb{T}_c \xrightarrow{\xi} \mathbb{T}''$. ■

2.2. PRELIMINARIES ON ENRICHED CATEGORIES.

In what follows, we work in the context of the theory of categories enriched in a symmetric monoidal category \mathcal{V} , as documented in the seminal paper [10] and the comprehensive references [17], [9]. We shall include an explicit indication of \mathcal{V} when employing notions such as \mathcal{V} -category, \mathcal{V} -functor, and so on, omitting the prefix \mathcal{V} only when concerned with the corresponding notions for non-enriched or *ordinary* categories. When such ordinary notions and terminology are applied to a given \mathcal{V} -category \mathcal{A} , they should be interpreted relative to the underlying ordinary category of \mathcal{A} . In the absence of any indication to the contrary, we will assume throughout that \mathcal{V} is a *closed* symmetric monoidal category, and in this case we denote by $\underline{\mathcal{V}}$ the \mathcal{V} -category canonically associated to \mathcal{V} , whose underlying ordinary category is isomorphic to \mathcal{V} ; in particular, the internal homs in \mathcal{V} will therefore be denoted by $\underline{\mathcal{V}}(V_1, V_2)$. We do not assume that any limits or colimits exist in \mathcal{V} .

2.2.1. The ordinary categories \mathcal{C} considered in this paper are *not* assumed *locally small*—that is, they are not necessarily Set-enriched categories. Rather, we assume that for each category \mathcal{C} under consideration, there is a category SET of *classes* in which lie the hom-classes of \mathcal{C} , so that \mathcal{C} is SET-enriched, but SET is not assumed cartesian closed.

2.2.2. A morphism $m : B_1 \rightarrow B_2$ in a \mathcal{V} -category \mathcal{B} (i.e., in the underlying ordinary category of \mathcal{B}) is a \mathcal{V} -*mono*(morphism) if $\mathcal{B}(A, m) : \mathcal{B}(A, B_1) \rightarrow \mathcal{B}(A, B_2)$ is a monomorphism in \mathcal{V} for each object $A \in \mathcal{B}$. A \mathcal{V} -*epi*(morphism) is a \mathcal{V} -mono in \mathcal{B}^{op} . We denote the classes of all \mathcal{V} -monos and \mathcal{V} -epis in \mathcal{B} by $\text{Mono}_{\mathcal{V}} \mathcal{B}$ and $\text{Epi}_{\mathcal{V}} \mathcal{B}$, respectively. ■

2.2.3. A \mathcal{V} -limit in a \mathcal{V} -category \mathcal{B} is a conical \mathcal{V} -enriched limit in the sense of [17] §3.8, equivalently a limit (in the underlying ordinary category of \mathcal{B}) that is preserved by each (ordinary) functor $\mathcal{B}(A, -) : \mathcal{B} \rightarrow \mathcal{V}$ ($A \in \mathcal{B}$). In particular, we obtain the notions of \mathcal{V} -product, \mathcal{V} -fibre-product, etc. \mathcal{V} -colimits in \mathcal{B} are \mathcal{V} -limits in \mathcal{B}^{op} . Any \mathcal{V} -fibre-product $m : A \rightarrow B$ of a family of \mathcal{V} -monos $(m_i : A_i \rightarrow B)_{i \in I}$ is again a \mathcal{V} -mono ([21] 2.7); we then say that m is a \mathcal{V} -intersection of the m_i .

2.2.4. PROPOSITION. Let $G : \mathcal{C} \rightarrow \mathcal{B}$ be a \mathcal{V} -functor, and suppose that for each $B \in \mathcal{B}$ we are given an object FB in \mathcal{C} and a morphism $\eta_B : B \rightarrow GFB$ in \mathcal{B} such that for each $C \in \mathcal{C}$, the composite

$$\phi_{BC} := \left(\mathcal{C}(FB, C) \xrightarrow{G_{FBC}} \mathcal{B}(GFB, GC) \xrightarrow{\mathcal{B}(\eta_B, GC)} \mathcal{B}(B, GC) \right)$$

is an isomorphism in \mathcal{V} . Then the given morphisms η_B constitute the unit η of a \mathcal{V} -adjunction $F \dashv \eta \mid G$ in which F acts in the given way on objects and is given on homs by formula (6) of Ch. 0 of [9].

PROOF. Fixing an object $B \in \mathcal{B}$, the weak Yoneda lemma ([17], 1.9) yields a bijection between morphisms $B \rightarrow GFB$ and \mathcal{V} -natural transformations $\mathcal{C}(FB, -) \rightarrow \mathcal{B}(B, G-)$, under which η_B corresponds to $\phi_{B-} := (\phi_{BC})_{C \in \mathcal{C}}$. In particular, $\phi_{B-} : \mathcal{C}(FB, -) \rightarrow \mathcal{B}(B, G-)$ is thus a \mathcal{V} -natural isomorphism, showing that $\mathcal{B}(B, G-) : \mathcal{C} \rightarrow \mathcal{V}$ is a representable \mathcal{V} -functor. Since this holds for all $B \in \mathcal{B}$, the result follows from Proposition 0.2 of [9]. ■

2.2.5. Let \mathcal{B} be a \mathcal{V} -category. A \mathcal{V} -reflective-subcategory of \mathcal{B} is a full replete sub- \mathcal{V} -category \mathcal{B}' of \mathcal{B} for which the inclusion \mathcal{V} -functor $J : \mathcal{B}' \hookrightarrow \mathcal{B}$ has a left \mathcal{V} -adjoint K . Any such \mathcal{V} -adjunction $K \dashv \rho \mid J : \mathcal{B}' \hookrightarrow \mathcal{B}$ is called a \mathcal{V} -reflection (on \mathcal{B}). Given an idempotent \mathcal{V} -monad $\mathbb{S} = (S, \rho, \lambda)$ (2.1.3) on \mathcal{B} , we let $\mathcal{B}^{(\mathbb{S})}$ denote the \mathcal{V} -reflective-subcategory of \mathcal{B} consisting of those objects B for which ρ_B is iso.

2.2.6. PROPOSITION. There is a bijection $\text{Refl}_{\mathcal{V}}(\mathcal{B}) \cong \text{IdmMnd}_{\mathcal{V}\text{-CAT}}(\mathcal{B})$ between the class $\text{Refl}_{\mathcal{V}}(\mathcal{B})$ of all \mathcal{V} -reflections on \mathcal{B} and the class $\text{IdmMnd}_{\mathcal{V}\text{-CAT}}(\mathcal{B})$ of all idempotent \mathcal{V} -monads on \mathcal{B} , which associates to each \mathcal{V} -reflection on \mathcal{B} its induced \mathcal{V} -monad. The \mathcal{V} -reflective-subcategory associated to a given idempotent \mathcal{V} -monad \mathbb{S} via this bijection is $\mathcal{B}^{(\mathbb{S})}$.

2.3. ENRICHED FACTORIZATION SYSTEMS.

Given morphisms $e : A_1 \rightarrow A_2$, $m : B_1 \rightarrow B_2$ in a \mathcal{V} -category \mathcal{B} , we say that e is \mathcal{V} -orthogonal to m , written $e \downarrow_{\mathcal{V}} m$, if the commutative square

$$\begin{array}{ccc} \mathcal{B}(A_2, B_1) & \xrightarrow{\mathcal{B}(A_2, m)} & \mathcal{B}(A_2, B_2) \\ \mathcal{B}(e, B_1) \downarrow & & \downarrow \mathcal{B}(e, B_2) \\ \mathcal{B}(A_1, B_1) & \xrightarrow{\mathcal{B}(A_1, m)} & \mathcal{B}(A_1, B_2) \end{array} \tag{2.3.i}$$

is a pullback in \mathcal{V} . Given classes \mathcal{E}, \mathcal{M} of morphisms in \mathcal{B} , we define associated classes of morphisms as follows:

$$\mathcal{E}^{\downarrow \mathcal{V}} := \{m \mid \forall e \in \mathcal{E} : e \downarrow_{\mathcal{V}} m\}, \quad \mathcal{M}^{\uparrow \mathcal{V}} := \{e \mid \forall m \in \mathcal{M} : e \downarrow_{\mathcal{V}} m\}.$$

The pair $(\mathcal{E}, \mathcal{M})$ is called a \mathcal{V} -prefactorization-system on \mathcal{B} if condition 1 below holds, and $(\mathcal{E}, \mathcal{M})$ is called a \mathcal{V} -factorization system if conditions 1 and 2 both hold:

1. $\mathcal{E}^{\downarrow \mathcal{V}} = \mathcal{M}$ and $\mathcal{M}^{\uparrow \mathcal{V}} = \mathcal{E}$.
2. Each morphism in \mathcal{B} factors as a morphism in \mathcal{E} followed by a morphism in \mathcal{M} .

For ordinary categories, with $\mathcal{V} = \text{SET}$ one obtains the familiar notion of *factorization system*; in this case, we drop the indication of \mathcal{V} from the notation. The relation of \mathcal{V} -factorization-systems to ordinary factorization systems is elaborated in [21], where theorems on the existence of \mathcal{V} -factorization-systems are proved as well.

2.3.1. Given a \mathcal{V} -prefactorization-system $(\mathcal{E}, \mathcal{M})$ on \mathcal{B} , the following stability properties of \mathcal{E} and \mathcal{M} are established in [21, 4.4]. The class \mathcal{M} is closed under composition, cotensors, arbitrary \mathcal{V} -fibre-products, and \mathcal{V} -pullbacks along arbitrary morphisms in \mathcal{B} . Further, if $g \cdot f \in \mathcal{M}$ and $g \in \mathcal{M}$, then $f \in \mathcal{M}$. Also, if $\mathcal{E} = \mathcal{H}^{\uparrow \mathcal{V}}$ for some class of \mathcal{V} -epimorphisms \mathcal{H} , then whenever $g \cdot f \in \mathcal{M}$, it follows that $f \in \mathcal{M}$. Since $(\mathcal{M}, \mathcal{E})$ is a \mathcal{V} -prefactorization-system in \mathcal{B}^{op} , one obtains stability properties for \mathcal{E} that are exactly dual to the above properties of \mathcal{M} . Analogous stability properties hold for a prefactorization system on an ordinary category \mathcal{B} , even when \mathcal{B} is not locally small; cf. [12, 2.1.1].

2.3.2. Given a \mathcal{V} -prefactorization-system $(\mathcal{E}, \mathcal{M})$ on \mathcal{B} with \mathcal{M} a class of \mathcal{V} -monos in \mathcal{B} , we say that \mathcal{B} is \mathcal{M} -subobject-complete (as a \mathcal{V} -category) if \mathcal{B} is cotensored and has \mathcal{V} -intersections (2.2.3) of arbitrary (class-indexed) families of \mathcal{M} -morphisms, as well as \mathcal{V} -pullbacks of \mathcal{M} -morphisms along arbitrary morphisms. By [21, 7.4], if \mathcal{B} is \mathcal{M} -subobject-complete, then the following hold:

1. $(\mathcal{E}, \mathcal{M})$ is a \mathcal{V} -factorization-system on \mathcal{B} .
2. For any class Σ of morphisms in \mathcal{B} , if we let $\mathcal{N} := \Sigma^{\downarrow \mathcal{V}} \cap \mathcal{M}$, then $(\mathcal{N}^{\uparrow \mathcal{V}}, \mathcal{N})$ is a \mathcal{V} -factorization-system on \mathcal{B} .

2.3.3. A \mathcal{V} -prefactorization-system $(\mathcal{E}, \mathcal{M})$ on \mathcal{B} is said to be \mathcal{V} -proper if every morphism in \mathcal{E} is a \mathcal{V} -epimorphism in \mathcal{B} and every morphism in \mathcal{M} is a \mathcal{V} -monomorphism. A \mathcal{V} -strong-mono(morphism) in \mathcal{B} is a \mathcal{V} -mono to which each \mathcal{V} -epi in \mathcal{B} is \mathcal{V} -orthogonal, and a \mathcal{V} -strong-epi(morphism) in \mathcal{B} is a \mathcal{V} -strong-mono in \mathcal{B}^{op} . We denote the classes of all such by $\text{StrMono}_{\mathcal{V}} \mathcal{B}$ and $\text{StrEpi}_{\mathcal{V}} \mathcal{B}$, respectively. In a tensored and cotensored \mathcal{V} -category, these notions reduce (by [21, 6.8]) to the familiar notions of strong monomorphism (resp. strong epimorphism), applied to the underlying ordinary category of \mathcal{B} . For any \mathcal{V} -proper \mathcal{V} -prefactorization-system $(\mathcal{E}, \mathcal{M})$, we have $\text{StrMono}_{\mathcal{V}} \mathcal{B} \subseteq \mathcal{E}^{\downarrow \mathcal{V}} = \mathcal{M}$ and similarly $\text{StrEpi}_{\mathcal{V}} \mathcal{B} \subseteq \mathcal{E}$; hence, since every section is a \mathcal{V} -strong-mono ([21, 6.3]), every section therefore lies in \mathcal{M} , and dually, every retraction lies in \mathcal{E} .

2.3.4. PROPOSITION. *If either of the following conditions holds, then $(\mathbf{Epi}_{\mathcal{V}} \mathcal{B}, \mathbf{StrMono}_{\mathcal{V}} \mathcal{B})$ is a \mathcal{V} -factorization-system on \mathcal{B} and \mathcal{B} is \mathcal{M} -subobject-complete for $\mathcal{M} = \mathbf{StrMono}_{\mathcal{V}} \mathcal{B}$.*

1. \mathcal{B} is cotensored, well-powered with respect to \mathcal{V} -strong-monos, and has small \mathcal{V} -limits and \mathcal{V} -cokernel-pairs.
2. \mathcal{B} is cotensored and tensored, well-powered with respect to strong monos, and has small limits.

If either of the following conditions holds, then $(\mathbf{StrEpi}_{\mathcal{V}} \mathcal{B}, \mathbf{Mono}_{\mathcal{V}} \mathcal{B})$ is a \mathcal{V} -factorization-system and \mathcal{B} is \mathcal{M} -subobject-complete for $\mathcal{M} = \mathbf{Mono}_{\mathcal{V}} \mathcal{B}$.

3. \mathcal{B} is cotensored, well-powered with respect to \mathcal{V} -monos, and has small \mathcal{V} -limits and \mathcal{V} -cokernel pairs.
4. \mathcal{B} is cotensored and tensored, well-powered, and has small limits.

PROOF. In each case, the statement that the pair $(\mathcal{E}, \mathcal{M})$ in question is a \mathcal{V} -factorization-system under the given condition is part of Theorem 7.14 of [21]. In cases 1 and 3, it is clear that \mathcal{B} is \mathcal{M} -subobject-complete. In cases 2 and 4, we deduce that \mathcal{B} is \mathcal{M} -subobject-complete by [21, 2.4, 6.8]. ■

2.4. CLOSURE OPERATORS IN CATEGORIES.

Let $(\mathcal{E}, \mathcal{M})$ be a prefactorization system on a category \mathcal{B} with $\mathcal{M} \subseteq \mathbf{Mono} \mathcal{B}$. For each object B of \mathcal{B} , denote by $\mathbf{Sub}_{\mathcal{M}}(B)$ the preordered class of all \mathcal{M} -morphisms with codomain B . Suppose that for each $f : A \rightarrow B$ in \mathcal{B} and $m \in \mathbf{Sub}_{\mathcal{M}}(B)$, the pullback $f^{-1}(m)$ of m along f exists; by 2.3.1, $f^{-1}(m)$ then lies in $\mathbf{Sub}_{\mathcal{M}}(A)$. Under these assumptions, we shall recall some basic results concerning the notion of *idempotent closure operator* defined in [7] and, in more general settings, in [8, 25].

2.4.1. An *idempotent closure operator* on \mathcal{M} in \mathcal{B} is an assignment to each object B of \mathcal{B} a monotone map $\overline{(-)} : \mathbf{Sub}_{\mathcal{M}}(B) \rightarrow \mathbf{Sub}_{\mathcal{M}}(B)$ such that

1. $m \leq \overline{m}$ and $\overline{\overline{m}} \leq \overline{m}$ for each $m \in \mathcal{M}$, and
2. for each $f : A \rightarrow B$ in \mathcal{B} and each $n \in \mathbf{Sub}_{\mathcal{M}}(B)$, $\overline{f^{-1}(n)} \leq f^{-1}(\overline{n})$.

An \mathcal{M} -morphism $m \in \mathbf{Sub}_{\mathcal{M}}(B)$ is said to be *closed* with respect to $\overline{(-)}$ if $\overline{m} \cong m$ in $\mathbf{Sub}_{\mathcal{M}}(B)$, whereas m is said to be *dense* if $\overline{m} \cong 1_B$. By 1, each \mathcal{M} -morphism $m : M \rightarrow B$ factors as $M \xrightarrow{d_m} \overline{M} \xrightarrow{\overline{m}} B$ for a unique morphism d_m , and by 2.3.1, $d_m \in \mathcal{M}$. We say that $\overline{(-)}$ is *weakly hereditary* if for each $m \in \mathcal{M}$, d_m is dense.

2.4.2. Suppose that $(\mathcal{E}, \mathcal{M})$ -factorizations exist. For all morphisms $f : A \rightarrow B$ and $m : M \rightarrow A$ with $m \in \mathcal{M}$, denote by $f(m)$ the second factor of the $(\mathcal{E}, \mathcal{M})$ -factorization of the composite $M \xrightarrow{m} A \xrightarrow{f} B$. Then condition 2 above is equivalent to the following condition:

$$2'. f(\overline{m}) \leq \overline{f(m)}.$$

Given a weakly hereditary idempotent closure operator $\overline{(-)}$ on \mathcal{M} in \mathcal{B} , we obtain an associated factorization system $(\text{Dense}, \text{CIEmb})$ on \mathcal{B} , where CIEmb is the class of all closed \mathcal{M} -morphisms and Dense is the class of all *dense* morphisms, i.e. those $f : A \rightarrow B$ in \mathcal{B} whose *image* $f(1_A)$ is a dense \mathcal{M} -morphism.

2.4.3. Given a factorization system $\mathcal{F} = (\mathcal{D}, \mathcal{C})$ with $\mathcal{C} \subseteq \mathcal{M}$, we obtain a weakly hereditary idempotent closure operator $\overline{(-)}^{\mathcal{F}}$ on \mathcal{M} in \mathcal{B} by defining the closure $\overline{m}^{\mathcal{F}}$ of each $m \in \mathcal{M}$ to be the second factor of the $(\mathcal{D}, \mathcal{C})$ -factorization of m . Supposing as in 2.4.2 that $(\mathcal{E}, \mathcal{M})$ -factorizations exist, the class of closed \mathcal{M} -morphisms (resp. dense morphisms) determined by $\overline{(-)}^{\mathcal{F}}$ is then equal to \mathcal{C} (resp. \mathcal{D}). In the case that $\mathcal{F} = (\mathcal{D}, \mathcal{C})$ is the factorization system associated to a given closure operator $\overline{(-)}$ on \mathcal{M} (2.4.2), we find that $\overline{m}^{\mathcal{F}} \cong \overline{m}$ in $\text{Sub}_{\mathcal{M}}(B)$ for all $m \in \text{Sub}_{\mathcal{M}}(B)$, $B \in \mathcal{B}$.

2.4.4. PROPOSITION. *Let $(\text{Dense}, \text{CIEmb})$ be the factorization system determined by a weakly hereditary idempotent closure operator $\overline{(-)}$ on \mathcal{M} in \mathcal{B} . Then*

$$\text{CIEmb} = \text{DenseEmb}^\downarrow \cap \mathcal{M},$$

where $\text{DenseEmb} := \mathcal{M} \cap \text{Dense}$.

PROOF. Since $(\text{Dense}, \text{CIEmb})$ is a prefactorization system and $\text{DenseEmb} \subseteq \text{Dense}$ we know that $\text{CIEmb} = \text{Dense}^\downarrow \subseteq \text{DenseEmb}^\downarrow$ and hence $\text{CIEmb} \subseteq \text{DenseEmb}^\downarrow \cap \mathcal{M}$. For the converse inclusion, suppose that $m : M \rightarrow B$ lies in $\text{DenseEmb}^\downarrow \cap \mathcal{M}$. Then, taking the $(\text{Dense}, \text{CIEmb})$ -factorization $M \xrightarrow{d} C \xrightarrow{c} B$ of m , we find by 2.3.1 that d lies in \mathcal{M} and hence lies in DenseEmb , so $d \downarrow m$. Therefore, there is a unique morphism k such that the diagram

$$\begin{array}{ccc} M & \xlongequal{\quad} & M \\ d \downarrow & \nearrow k & \downarrow m \\ C & \xrightarrow{c} & B \end{array}$$

commutes, whence $c \cong m$ in $\text{Sub}_{\mathcal{M}}(B)$, so that m is closed as c is so. ■

3. Orthogonality, adjunctions, and reflections

The following notion of orthogonality in the enriched context was employed in [6].

3.1. DEFINITION. Let \mathcal{B} be a \mathcal{V} -category, let Σ be a class of morphisms in \mathcal{B} , and let \mathcal{C} be a class of objects in \mathcal{B} .

1. For a morphism $f : A_1 \rightarrow A_2$ in \mathcal{B} and an object B in \mathcal{B} , we say that f is \mathcal{V} -orthogonal to B , written $f \perp_{\mathcal{V}} B$, if $\mathcal{B}(f, B) : \mathcal{B}(A_2, B) \rightarrow \mathcal{B}(A_1, B)$ is an isomorphism in \mathcal{V} .
2. We define $\Sigma^{\perp_{\mathcal{V}}} := \{C \in \text{Ob } \mathcal{B} \mid \forall f \in \Sigma : f \perp_{\mathcal{V}} C\}$. We let \mathcal{B}_{Σ} be the full sub- \mathcal{V} -category of \mathcal{B} whose objects are those in $\Sigma^{\perp_{\mathcal{V}}}$.
3. We define $\mathcal{C}^{\top_{\mathcal{V}}} := \{f \in \text{Mor } \mathcal{B} \mid \forall C \in \mathcal{C} : f \perp_{\mathcal{V}} C\}$.
4. We say that (Σ, \mathcal{C}) is a \mathcal{V} -orthogonal-pair in \mathcal{B} if $\Sigma^{\perp_{\mathcal{V}}} = \mathcal{C}$ and $\mathcal{C}^{\top_{\mathcal{V}}} = \Sigma$.
5. Given a functor $F : \mathcal{B} \rightarrow \mathcal{C}$, we denote by Σ_F the class of all morphisms in \mathcal{B} inverted by F (i.e. sent to isomorphisms in \mathcal{C}).

3.2. REMARK. For an ordinary category \mathcal{B} , with $\mathcal{V} = \text{SET}$ we obtain the familiar notions of orthogonality [12] and orthogonal pair [4], for which we omit the indication of \mathcal{V} and employ the unadorned symbols \perp, \top . Enriched orthogonality clearly implies ordinary orthogonality.

3.3. REMARK. For any class of morphisms Σ in \mathcal{B} , $(\Sigma^{\perp_{\mathcal{V}} \top_{\mathcal{V}}}, \Sigma^{\perp_{\mathcal{V}}})$ is a \mathcal{V} -orthogonal-pair in \mathcal{B} . For any class of objects \mathcal{C} in \mathcal{B} , $(\mathcal{C}^{\top_{\mathcal{V}}}, \mathcal{C}^{\top_{\mathcal{V}} \perp_{\mathcal{V}}})$ is a \mathcal{V} -orthogonal-pair in \mathcal{B} .

3.4. REMARK. If \mathcal{B} has a \mathcal{V} -terminal object 1 , then it is easy to show that $f \perp_{\mathcal{V}} B$ iff $f \downarrow_{\mathcal{V}} !_B$, where $!_B : B \rightarrow 1$.

3.5. PROPOSITION. Let (Σ, \mathcal{C}) be a \mathcal{V} -orthogonal-pair in a \mathcal{V} -category \mathcal{B} . Then

1. Σ is closed under \mathcal{V} -enriched weighted colimits in \mathcal{B} . I.e., given a \mathcal{V} -natural-transformation $h : B \Rightarrow B' : \mathcal{J} \rightarrow \mathcal{B}$ whose components lie in Σ , if $W : \mathcal{J}^{\text{op}} \rightarrow \underline{\mathcal{V}}$ is a \mathcal{V} -functor for which weighted colimits $W \otimes B$ and $W \otimes B'$ exist in \mathcal{B} , then the induced morphism $W \otimes h : W \otimes B \rightarrow W \otimes B'$ lies in Σ .
2. \mathcal{C} is closed under \mathcal{V} -enriched weighted limits in \mathcal{B} . I.e., given \mathcal{V} -functors $C : \mathcal{J} \rightarrow \mathcal{B}$ and $W : \mathcal{J} \rightarrow \underline{\mathcal{V}}$ for which a weighted limit $[W, C]$ in \mathcal{B} exists, if $C_j \in \mathcal{C}$ for all $j \in \mathcal{J}$, then $[W, C] \in \mathcal{C}$.

PROOF. 1. For each object $C \in \mathcal{C}$, we have an isomorphism

$$\mathcal{B}(W \otimes h, C) \cong [\mathcal{J}^{\text{op}}, \underline{\mathcal{V}}](W, \mathcal{B}(h_-, C)) = \int_{j \in \mathcal{J}} \underline{\mathcal{V}}(Wj, \mathcal{B}(h_j, C))$$

in the arrow category of \mathcal{V} , but each $\mathcal{B}(h_j, C)$ is iso and hence $\mathcal{B}(W \otimes h, C)$ is iso, showing that $W \otimes h \in \mathcal{C}^{\top_{\mathcal{V}}} = \Sigma$. 2. For each $h \in \Sigma$, we have an isomorphism

$$\mathcal{B}(h, [W, C]) \cong [\mathcal{J}, \underline{\mathcal{V}}](W, \mathcal{B}(h, C_-)) = \int_{j \in \mathcal{J}} \underline{\mathcal{V}}(Wj, \mathcal{B}(h, C_j))$$

in the arrow category of \mathcal{V} , but each $\mathcal{B}(h, C_j)$ is iso and hence $\mathcal{B}(h, [W, C])$ is iso, showing that $[W, C] \in \Sigma^{\perp_{\mathcal{V}}} = \mathcal{C}$. ■

3.6. PROPOSITION. *Let \mathcal{B} be a \mathcal{V} -category, $\Sigma \subseteq \text{Mor } \mathcal{B}$, $\mathcal{C} \subseteq \text{Ob } \mathcal{B}$.*

1. *If \mathcal{B} is tensored and Σ is closed under tensors in \mathcal{B} , then $\Sigma^{\perp_{\mathcal{V}}} = \Sigma^{\perp}$.*

2. *If \mathcal{B} is cotensored and \mathcal{C} is closed under cotensors in \mathcal{B} , then $\mathcal{C}^{\top_{\mathcal{V}}} = \mathcal{C}^{\top}$.*

PROOF. 1. Given $C \in \Sigma^{\perp}$ and $h \in \Sigma$, it suffices to show that $h \perp_{\mathcal{V}} C$. Letting \mathcal{B}_0 denote the underlying ordinary category of \mathcal{B} , we have that for each $V \in \mathcal{V}$, $\mathcal{V}(V, \mathcal{B}(h, C)) \cong \mathcal{B}_0(V \otimes h, C)$ in the arrow category of SET, and the latter morphism is iso. Hence $\mathcal{B}(h, C)$ is an isomorphism in \mathcal{V} . 2 is proved analogously. ■

3.7. PROPOSITION. *For a \mathcal{V} -category \mathcal{B} , $(\text{Ob } \mathcal{B})^{\top_{\mathcal{V}}} = \text{Iso } \mathcal{B} = (\text{Mor } \mathcal{B})^{\uparrow_{\mathcal{V}}}$.*

PROOF. $\text{Iso } \mathcal{B}$ is clearly included in both the rightmost and leftmost classes. Also, the inclusion $(\text{Mor } \mathcal{B})^{\uparrow_{\mathcal{V}}} \subseteq \text{Iso } \mathcal{B}$ follows from [21, 3.7]. Lastly, if $h : B_1 \rightarrow B_2$ lies in $(\text{Ob } \mathcal{B})^{\top_{\mathcal{V}}}$, then the \mathcal{V} -natural transformation $\mathcal{B}(h, -) : \mathcal{B}(B_2, -) \rightarrow \mathcal{B}(B_1, -)$ is an isomorphism; but by the weak Yoneda lemma ([17, 1.9]), the (ordinary) functor $Y : \mathcal{B}^{\text{op}} \rightarrow \mathcal{V}\text{-CAT}(\mathcal{B}, \mathcal{V})$ given by $YB = \mathcal{B}(B, -)$ is fully faithful, so h is iso. ■

3.8. PROPOSITION. *Let \mathcal{B} be a \mathcal{V} -category and Σ a class of morphisms in \mathcal{B} . Then for any morphisms $e : A_1 \rightarrow A_2$ in Σ and $m : B_1 \rightarrow B_2$ in \mathcal{B}_{Σ} , we have that $e \downarrow_{\mathcal{V}} m$ in \mathcal{B} .*

PROOF. Since $e \perp_{\mathcal{V}} B_1$ and $e \perp_{\mathcal{V}} B_2$, the left and right sides of the commutative square (2.3.i) are isomorphisms, so the square is a pullback. ■

In the non-enriched context, the first of the following equivalences appears in the proof of Lemma 4.2.1 of [12], and variants of both equivalences are given in [23].

3.9. PROPOSITION. *Let $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ be a \mathcal{V} -adjunction, $f : B_1 \rightarrow B_2$ a morphism in \mathcal{B} , $g : C_1 \rightarrow C_2$ a morphism in \mathcal{C} , and C an object of \mathcal{C} .*

1. $Ff \downarrow_{\mathcal{V}} g \iff f \downarrow_{\mathcal{V}} Gg$.

2. $Ff \perp_{\mathcal{V}} C \iff f \perp_{\mathcal{V}} GC$.

PROOF. 1. Via the given \mathcal{V} -adjunction, the commutative diagram

$$\begin{array}{ccc} \mathcal{C}(FB_2, C_1) & \xrightarrow{\mathcal{C}(FB_2, g)} & \mathcal{C}(FB_2, C_2) \\ \mathcal{C}(Ff, C_1) \downarrow & & \downarrow \mathcal{C}(Ff, C_2) \\ \mathcal{C}(FB_1, C_1) & \xrightarrow{\mathcal{C}(FB_1, g)} & \mathcal{C}(FB_1, C_2) \end{array}$$

is isomorphic to the commutative diagram

$$\begin{array}{ccc} \mathcal{B}(B_2, GC_1) & \xrightarrow{\mathcal{B}(B_2, Gg)} & \mathcal{B}(B_2, GC_2) \\ \mathcal{B}(f, GC_1) \downarrow & & \downarrow \mathcal{B}(f, GC_2) \\ \mathcal{B}(B_1, GC_1) & \xrightarrow{\mathcal{B}(B_1, Gg)} & \mathcal{B}(B_1, GC_2) . \end{array}$$

2. $\mathcal{C}(Ff, C) \cong \mathcal{B}(f, GC)$ in the arrow category $[2, \mathcal{V}]$. ■

In the non-enriched setting, the first of the following equations is noted in [5, 3.3] and specializes [12, 4.2.1].

3.10. COROLLARY. *Let $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ be a \mathcal{V} -adjunction. Then*

$$(G(\text{Mor } \mathcal{C}))^{\uparrow \nu} = \Sigma_F = (G(\text{Ob } \mathcal{C}))^{\top \nu} .$$

Hence, in particular, $(\Sigma_F, \Sigma_F^{\downarrow \nu})$ is a \mathcal{V} -prefactorization-system on \mathcal{B} , and $(\Sigma_F, \Sigma_F^{\downarrow \nu})$ is a \mathcal{V} -orthogonal-pair in \mathcal{B} .

PROOF. By 3.9 and 3.7, we may compute as follows:

$$(G(\text{Mor } \mathcal{C}))^{\uparrow \nu} = F^{-1}((\text{Mor } \mathcal{C})^{\uparrow \nu}) = F^{-1}(\text{Iso } \mathcal{C}) = \Sigma_F;$$

$$(G(\text{Ob } \mathcal{C}))^{\top \nu} = F^{-1}((\text{Ob } \mathcal{C})^{\top \nu}) = F^{-1}(\text{Iso } \mathcal{C}) = \Sigma_F.$$

■

Clearly any sub- \mathcal{V} -category of \mathcal{B} of the form \mathcal{B}_Σ is replete. The following proposition shows that every \mathcal{V} -reflective-subcategory of \mathcal{B} is of the form \mathcal{B}_Σ for each of two canonical choices of Σ :

3.11. PROPOSITION. *Let $K \overset{\rho}{\dashv} J : \mathcal{C} \hookrightarrow \mathcal{B}$ be a \mathcal{V} -reflection.*

1. $\mathcal{C} = \mathcal{B}_{\Sigma_K} = \mathcal{B}_\Sigma$, where $\Sigma := \{\rho_B \mid B \in \mathcal{B}\}$.
2. $(\text{Mor } \mathcal{C})^{\uparrow \nu} = \Sigma_K = (\text{Ob } \mathcal{C})^{\top \nu}$, where the right- and leftmost expressions are evaluated with respect to \mathcal{B} .
3. $(\Sigma_K, \Sigma_K^{\downarrow \nu})$ is a \mathcal{V} -prefactorization-system on \mathcal{B} , and (Σ_K, \mathcal{C}) is a \mathcal{V} -orthogonal-pair in \mathcal{B} .

PROOF. 2 follows from 3.10. Regarding 1, first observe that $\Sigma \subseteq \Sigma_K$, so that $\mathcal{B}_{\Sigma_K} \subseteq \mathcal{B}_\Sigma$. Also, by 2, $\text{Ob } \mathcal{B}_{\Sigma_K} = (\Sigma_K)^{\perp \nu} = (\text{Ob } \mathcal{C})^{\top \nu \perp \nu} \supseteq \text{Ob } \mathcal{C}$. Hence it now suffices to show $\mathcal{B}_\Sigma \subseteq \mathcal{C}$. Suppose $B \in \mathcal{B}_\Sigma$. Since we also have that $KB \in \mathcal{C} \subseteq \mathcal{B}_\Sigma$, the morphism $\rho_B : B \rightarrow KB$ lies in \mathcal{B}_Σ , so since $\rho_B \in \Sigma$ we deduce by 3.8 that $\rho_B \downarrow_\nu \rho_B$. Hence by [21, 3.7], ρ_B is iso, so $B \in \mathcal{C}$. Lastly, observe that 3 follows from 3.10 and 1. ■

3.12. PROPOSITION. *Let \mathcal{C} be a reflective subcategory of the underlying ordinary category of a \mathcal{V} -category \mathcal{B} , and suppose that \mathcal{B} is cotensored and \mathcal{C} is closed under cotensors in \mathcal{B} . Then \mathcal{C} is a \mathcal{V} -reflective-subcategory of \mathcal{B} .*

PROOF. For all $B \in \mathcal{B}$, the reflection morphism $\rho_B : B \rightarrow KB$ lies in \mathcal{C}^\top , but by 3.6, $\mathcal{C}^\top = \mathcal{C}^{\top \nu}$, so $\rho_B \perp_\nu C$ for all $C \in \mathcal{C}$ and the result follows by 2.2.4. ■

3.13. DEFINITION. Given a full sub- \mathcal{V} -category \mathcal{C} of a \mathcal{V} -category \mathcal{B} , the \mathcal{V} -reflective hull of \mathcal{C} (in \mathcal{B}), if it exists, is the smallest \mathcal{V} -reflective-subcategory of \mathcal{B} containing \mathcal{C} .

3.14. PROPOSITION. Let \mathcal{C} be a full sub- \mathcal{V} -category a \mathcal{V} -category \mathcal{B} .

1. Any \mathcal{V} -reflective-subcategory of \mathcal{B} that contains \mathcal{C} must also contain $\mathcal{C}^{\top_{\mathcal{V}} \perp_{\mathcal{V}}}$.
2. Hence, if $\mathcal{C}^{\top_{\mathcal{V}} \perp_{\mathcal{V}}}$ is a \mathcal{V} -reflective-subcategory of \mathcal{B} , then the \mathcal{V} -reflective-hull of \mathcal{C} in \mathcal{B} exists and equals $\mathcal{C}^{\top_{\mathcal{V}} \perp_{\mathcal{V}}}$.

PROOF. If $\mathcal{C} \hookrightarrow \mathcal{D} \hookrightarrow \mathcal{B}$ and the latter inclusion has left \mathcal{V} -adjoint $K : \mathcal{B} \rightarrow \mathcal{D}$, then $\mathcal{C}^{\top_{\mathcal{V}} \perp_{\mathcal{V}}} \subseteq \mathcal{D}^{\top_{\mathcal{V}} \perp_{\mathcal{V}}} = \Sigma_K^{\perp_{\mathcal{V}}} = \mathcal{D}$ by 3.11 2 & 1. \blacksquare

3.15. COROLLARY. Let $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ be a \mathcal{V} -adjunction. Then if $\mathcal{B}_{\Sigma_F} \hookrightarrow \mathcal{B}$ is a \mathcal{V} -reflective-subcategory, it follows that the \mathcal{V} -reflective-hull of $G(\text{Ob } \mathcal{C}) \hookrightarrow \mathcal{B}$ exists and equals \mathcal{B}_{Σ_F} .

PROOF. $(G(\text{Ob } \mathcal{C}))^{\top_{\mathcal{V}} \perp_{\mathcal{V}}} = \Sigma_F^{\perp_{\mathcal{V}}} = \mathcal{B}_{\Sigma_F}$ by 3.10, so the result follows from 3.14 2. \blacksquare

4. Definition and characterizations of the idempotent core

4.1. LEMMA. Let $\mathbb{S} = (S, \rho, \lambda)$ and $\mathbb{T} = (T, \eta, \mu)$ be monads on an object \mathcal{B} of 2-category \mathcal{X} , and suppose that \mathbb{S} is idempotent. Then

1. A 2-cell $\alpha : S \rightarrow T$ is a morphism of monads $\mathbb{S} \rightarrow \mathbb{T}$ if and only if $\alpha \cdot \rho = \eta$.
2. If a morphism of monads $\mathbb{S} \rightarrow \mathbb{T}$ exists, then it is unique.

PROOF. One of the implications in 1 is trivial; for the other, suppose that $\alpha \cdot \rho = \eta$. We must show that $\mu \cdot (\alpha \circ \alpha) = \alpha \cdot \lambda$. But λ is an isomorphism with $\lambda^{-1} = \rho S$, and $\mu \cdot (\alpha \circ \alpha) \cdot \rho S = \mu \cdot \alpha T \cdot S \alpha \cdot \rho S = \mu \cdot \alpha T \cdot \rho T \cdot \alpha = \mu \cdot \eta T \cdot \alpha = \alpha$. Regarding 2, let $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ be a morphism of monads. By 2.1.2, the 1-cell T carries the structure of an \mathbb{S} -algebra (T, β) where β is the composite $ST \xrightarrow{\alpha T} TT \xrightarrow{\mu} T$. But by 2.1.3, since \mathbb{S} is idempotent, β is an isomorphism with inverse $\rho T : T \rightarrow ST$. Hence by 2.1.2, $\alpha = \beta \cdot S \eta = (\rho T)^{-1} \cdot S \eta$, so we have expressed α in terms of \mathbb{S} and \mathbb{T} , showing that α is the unique monad morphism $\mathbb{S} \rightarrow \mathbb{T}$. \blacksquare

4.2. DEFINITION. Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} .

1. If $\tilde{\mathbb{T}}$ is an idempotent \mathcal{V} -monad on \mathcal{B} for which there exists a (necessarily unique, 4.1) morphism $\iota_{\mathbb{T}} : \tilde{\mathbb{T}} \rightarrow \mathbb{T}$ satisfying the following condition, then we say that $\tilde{\mathbb{T}}$ is a *terminal idempotent \mathcal{V} -monad over \mathbb{T}* :

$$\text{For each morphism of } \mathcal{V}\text{-monads } \alpha : \mathbb{S} \rightarrow \mathbb{T} \text{ with } \mathbb{S} \text{ idempotent, there is a unique morphism } \alpha_{\sharp} : \mathbb{S} \rightarrow \tilde{\mathbb{T}} \text{ with } \iota_{\mathbb{T}} \cdot \alpha_{\sharp} = \alpha.$$

2. If $\tilde{\mathbb{T}}$ is an idempotent \mathcal{V} -monad on \mathcal{B} whose underlying endofunctor \tilde{T} inverts the same morphisms as T , then we say that $\tilde{\mathbb{T}}$ is an *idempotent (\mathcal{V} -)core of \mathbb{T}* .

4.3. **REMARK.** A terminal idempotent \mathcal{V} -monad over \mathbb{T} is equally a terminal object in the category of idempotent \mathcal{V} -monads over \mathbb{T} and so is unique, up to isomorphism, if it exists. We will see in 4.15 that any idempotent core of \mathbb{T} is in particular a terminal idempotent \mathcal{V} -monad over \mathbb{T} . Hence, an idempotent core of \mathbb{T} is unique, up to isomorphism, if it exists, in which case any terminal idempotent \mathcal{V} -monad over \mathbb{T} is an idempotent core.

4.4. **REMARK.** If for every \mathcal{V} -monad \mathbb{T} on \mathcal{B} , the terminal idempotent \mathcal{V} -monad over \mathbb{T} exists, then the full subcategory $\text{IdmMnd}_{\mathcal{V}\text{-CAT}}(\mathcal{B})$ of $\text{Mnd}_{\mathcal{V}\text{-CAT}}(\mathcal{B})$ consisting of all idempotent \mathcal{V} -monads is a coreflective subcategory.

4.5. **LEMMA.** *Suppose given a \mathcal{V} -adjunction $F \frac{\eta}{\varepsilon} \Big| G : \mathcal{C} \rightarrow \mathcal{B}$ and a \mathcal{V} -reflection $K \frac{\rho}{\varepsilon} \Big| J : \mathcal{B}' \hookrightarrow \mathcal{B}$ such that the image of G lies in \mathcal{B}' . Then there is a \mathcal{V} -adjunction $F' \frac{\eta'}{\varepsilon'} \Big| G' : \mathcal{C} \rightarrow \mathcal{B}'$ with $JG' = G$, $F'K \cong F$, $F' = FJ$, $J\eta' = \eta J$, and $\varepsilon' = \varepsilon$. Hence by 2.1.6, the adjunction $F \frac{\eta}{\varepsilon} \Big| G$ is isomorphic to the composite of the \mathcal{V} -adjunctions $K \frac{\rho}{\varepsilon} \Big| J$ and $F' \frac{\eta'}{\varepsilon'} \Big| G'$.*

PROOF. G' is just the corestriction of G , the components of η' are just those of η ; the \mathcal{V} -naturality of η' is immediate, and the triangular equations are readily verified. \blacksquare

4.6. **DEFINITION.** Given data as in 4.5, we say that $F \frac{\eta}{\varepsilon} \Big| G$ *factors through* \mathcal{B}' , and, equivalently, that $F \frac{\eta}{\varepsilon} \Big| G$ *factors through* $K \frac{\rho}{\varepsilon} \Big| J$.

4.7. **PROPOSITION.** *Let \mathbb{T} be a \mathcal{V} -monad, and let $F \frac{\eta}{\varepsilon} \Big| G : \mathcal{C} \rightarrow \mathcal{B}$ be any \mathcal{V} -adjunction inducing \mathbb{T} . Let \mathbb{S} be an idempotent \mathcal{V} -monad on \mathcal{B} , with associated \mathcal{V} -reflection $K \frac{\rho}{\varepsilon} \Big| J : \mathcal{B}' \hookrightarrow \mathcal{B}$. Then the following are equivalent:*

1. *There exists a (necessarily unique, 4.1) morphism of \mathcal{V} -monads $\alpha : \mathbb{S} \rightarrow \mathbb{T}$.*
2. *\mathcal{B}' contains each object TB (with $B \in \mathcal{B}$).*
3. *$F \frac{\eta}{\varepsilon} \Big| G$ factors through $K \frac{\rho}{\varepsilon} \Big| J$.*

PROOF. Observe that 2 is equivalent to the statement that the Kleisli \mathcal{V} -adjunction for \mathbb{T} factors through $K \frac{\rho}{\varepsilon} \Big| J$. Hence it suffices to prove that $1 \Leftrightarrow 3$, for then the equivalence $1 \Leftrightarrow 2$ follows as a special case. If 3 holds, then the existence of a morphism of \mathcal{V} -monads $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ is guaranteed by 2.1.7. For the converse implication, let us assume 1 and prove that 3 holds. Working with only the underlying ordinary monad morphism and adjunction, note that the given adjunction determines a comparison functor $\mathcal{C} \rightarrow \mathcal{B}^{\mathbb{T}}$, and we have also a functor $\mathcal{B}^{\alpha} : \mathcal{B}^{\mathbb{T}} \rightarrow \mathcal{B}^{\mathbb{S}}$ induced by α . Both these functors commute with the forgetful functors to \mathcal{B} , and so too does their composite $\mathcal{C} \rightarrow \mathcal{B}^{\mathbb{T}} \rightarrow \mathcal{B}^{\mathbb{S}} \cong \mathcal{B}'$. Hence, applying this composite functor to any given $C \in \mathcal{C}$, we find that the carrier GC of the associated \mathbb{S} -algebra lies in \mathcal{B}' , so 3 holds. \blacksquare

4.8. **REMARK.** Proposition 4.7 shows in particular that the question of whether a \mathcal{V} -adjunction $F \frac{\eta}{\varepsilon} \Big| G$ factors through a given \mathcal{V} -reflection depends only on the \mathcal{V} -monad \mathbb{T} induced by $F \frac{\eta}{\varepsilon} \Big| G$.

4.9. **COROLLARY.** *Suppose $\mathbb{S} = (S, \rho, \lambda)$ and $\mathbb{T} = (T, \eta, \mu)$ are \mathcal{V} -monads on a \mathcal{V} -category \mathcal{B} with \mathbb{S} idempotent, and let $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ be a morphism of \mathcal{V} -monads. Then for each $B \in \mathcal{B}$, $\alpha_B : SB \rightarrow TB$ is the unique morphism such that $\alpha_B \cdot \rho_B = \eta_B$.*

PROOF. By 4.7, we know that TB lies in the reflective subcategory $\mathcal{B}' \hookrightarrow \mathcal{B}$ determined by \mathbb{S} , so α_B is the unique extension of $\eta_B : B \rightarrow TB$ along the reflection unit component $\rho_B : B \rightarrow SB$. ■

4.10. Given a \mathcal{V} -category \mathcal{B} , the class $\text{Refl}_{\mathcal{V}}(\mathcal{B})$ (2.2.6) of all \mathcal{V} -reflections on \mathcal{B} acquires the structure of a preordered class when ordered by inclusion of the associated \mathcal{V} -reflective-subcategories.

4.11. **COROLLARY.** *Let \mathcal{B} be a \mathcal{V} -category.*

1. *The full subcategory $\text{IdmMnd}_{\mathcal{V}\text{-CAT}}(\mathcal{B})$ of $\text{Mnd}_{\mathcal{V}\text{-CAT}}(\mathcal{B})$ is a preordered class isomorphic to $(\text{Refl}_{\mathcal{V}}(\mathcal{B}))^{\text{op}}$ via the bijection given in 2.2.6.*
2. *Given a \mathcal{V} -monad \mathbb{T} on \mathcal{B} , the isomorphism in 1 restricts to an isomorphism between the full subcategories determined by the following objects:*
 - (a) *Idempotent \mathcal{V} -monads \mathbb{S} on \mathcal{B} for which a (necessarily unique, 4.1) morphism of \mathcal{V} -monads $\alpha : \mathbb{S} \rightarrow \mathbb{T}$ exists.*
 - (b) *\mathcal{V} -reflections on \mathcal{B} whose associated \mathcal{V} -reflective-subcategory contains each object TB ($B \in \mathcal{B}$).*

PROOF. We shall prove 1, and then 2 follows by 4.7. By 4.1, $\text{IdmMnd}_{\mathcal{V}}(\mathcal{B})$ is a preorder, and it suffices to show that the bijection $\text{Refl}_{\mathcal{V}}(\mathcal{B}) \rightarrow \text{IdmMnd}_{\mathcal{V}}(\mathcal{B})$ (2.2.6) and its inverse are contravariantly functorial (i.e. order-reversing). But this follows from 4.7, since the given preorder relation on $\text{Refl}_{\mathcal{V}}(\mathcal{B})$ may equally be described as

$$(K' \frac{\rho'}{\varepsilon} \Big| J') \leq (K \frac{\rho}{\varepsilon} \Big| J) \iff K' \frac{\rho'}{\varepsilon} \Big| J' \text{ factors through } K \frac{\rho}{\varepsilon} \Big| J.$$

■

4.12. **THEOREM.** *Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} , and let $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ be any \mathcal{V} -adjunction inducing \mathbb{T} . Then the following are equivalent:*

1. *The terminal idempotent \mathcal{V} -monad over \mathbb{T} exists.*
2. *The full sub- \mathcal{V} -category $T(\text{Ob } \mathcal{B}) \hookrightarrow \mathcal{B}$ has a \mathcal{V} -reflective hull.*
3. *The full sub- \mathcal{V} -category $G(\text{Ob } \mathcal{C}) \hookrightarrow \mathcal{B}$ has a \mathcal{V} -reflective hull.*
4. *There is a smallest \mathcal{V} -reflective-subcategory through which $F \dashv G$ factors.*

Further, given an idempotent \mathcal{V} -monad $\tilde{\mathbb{T}}$ on \mathcal{B} with associated \mathcal{V} -reflective-subcategory \mathcal{B}' , $\tilde{\mathbb{T}}$ is a terminal idempotent \mathcal{V} -monad over \mathbb{T} if and only if \mathcal{B}' is a \mathcal{V} -reflective hull of $T(\mathbf{Ob} \mathcal{B})$, resp. $G(\mathbf{Ob} \mathcal{C})$, equivalently, a smallest \mathcal{V} -reflective-subcategory through which $F \dashv G$ factors.

PROOF. A terminal idempotent \mathcal{V} -monad over \mathbb{T} is by definition a terminal object of the preordered class described in 4.11 2(a), so the result follows from 4.11 2 and 4.7. ■

4.13. LEMMA. *Given an (ordinary) adjunction $F \overset{\eta}{\dashv} G : \mathcal{C} \rightarrow \mathcal{B}$ with induced endofunctor T on \mathcal{B} , $\Sigma_F = \Sigma_T$. Hence all left adjoints inducing a given monad \mathbb{T} invert the same morphisms.*

PROOF. One inclusion is immediate. For the other, suppose $f : B \rightarrow B'$ in \mathcal{B} is inverted by T . Then Tf has an inverse $(Tf)^{-1} : TB' \rightarrow TB$, and one easily shows that the transpose $FB' \rightarrow FB$ of the composite $B' \xrightarrow{\eta_{B'}} GFB' \xrightarrow{(Tf)^{-1}} GFB$ under the given adjunction serves as inverse for Ff . ■

4.14. COROLLARY. *Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} . Then $(\Sigma_T, \Sigma_T^{\perp \mathcal{V}})$ is a \mathcal{V} -prefactorization-system on \mathcal{B} , and $(\Sigma_T, \Sigma_T^{\perp \mathcal{V}})$ is a \mathcal{V} -orthogonal-pair in \mathcal{B} .*

PROOF. Taking any \mathcal{V} -adjunction $F \dashv G$ inducing \mathbb{T} (e.g., the Kleisli \mathcal{V} -adjunction), we have that $\Sigma_T = \Sigma_F$ by 4.13, and the result follows from 3.10. ■

4.15. THEOREM. *Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} , and let $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ be any \mathcal{V} -adjunction inducing \mathbb{T} . Then the following are equivalent:*

1. *The idempotent core of \mathbb{T} exists.*
2. *The terminal idempotent \mathcal{V} -monad over \mathbb{T} exists, and its underlying endofunctor inverts the same morphisms as T .*
3. *$\mathcal{B}_{\Sigma_T} (= \Sigma_T^{\perp \mathcal{V}})$ is a \mathcal{V} -reflective-subcategory of \mathcal{B} .*
4. *$F \dashv G$ factors through a \mathcal{V} -reflection $K \dashv J : \mathcal{B}' \hookrightarrow \mathcal{B}$ in such a way that the induced left \mathcal{V} -adjoint $F' : \mathcal{B}' \rightarrow \mathcal{C}$ (4.6) is conservative (i.e. reflects isomorphisms).*

Further, given an idempotent \mathcal{V} -monad $\tilde{\mathbb{T}}$ on \mathcal{B} with associated \mathcal{V} -reflective-subcategory \mathcal{B}' , the following are equivalent: (i) $\tilde{\mathbb{T}}$ is an idempotent core of \mathbb{T} , (ii) $\tilde{\mathbb{T}}$ is a terminal idempotent \mathcal{V} -monad over \mathbb{T} and inverts the same morphisms as T , (iii) $\mathcal{B}' = \mathcal{B}_{\Sigma_T}$, (iv) $F \dashv G$ factors through $\mathcal{B}' \hookrightarrow \mathcal{B}$ in such a way that the induced left \mathcal{V} -adjoint F' is conservative.

PROOF. We prove the equivalence of (i)-(iv), from which the equivalence of 1-4 follows. Let $K \dashv J : \mathcal{B}' \rightarrow \mathcal{B}$ be the \mathcal{V} -reflection determined by $\widetilde{\mathbb{T}}$. The implication (ii) \Rightarrow (i) is immediate. To see that (i) \Rightarrow (iii), observe that if (i) holds then $\Sigma_T = \Sigma_{\widetilde{\mathbb{T}}} = \Sigma_K$, and so $\mathcal{B}_{\Sigma_T} = \mathcal{B}_{\Sigma_K} = \mathcal{B}'$ by 3.11 1. Regarding the implication (iv) \Rightarrow (iii), we reason that if (iv) holds then we have that $\Sigma_K = \Sigma_{F'K}$ since F' is conservative, but $F'K \cong F$ by 4.5 and $\Sigma_F = \Sigma_T$ by 4.13; therefore $\Sigma_K = \Sigma_{F'K} = \Sigma_F = \Sigma_T$, and using 3.11 1 we deduce that $\mathcal{B}' = \mathcal{B}_{\Sigma_K} = \mathcal{B}_{\Sigma_T}$. It now suffices to prove the implications (iii) \Rightarrow (ii) and (iii) \Rightarrow (iv). Assuming (iii), we have that $\mathcal{B}' = \mathcal{B}_{\Sigma_T} = \mathcal{B}_{\Sigma_F} = \Sigma_F^{\perp \mathcal{V}}$, since $\Sigma_T = \Sigma_F$ by 4.13. Hence since \mathcal{B}' is a \mathcal{V} -reflective-subcategory of \mathcal{B} we deduce by 3.15 that the \mathcal{V} -reflective-hull of $G(\text{Ob } \mathcal{C})$ exists and equals \mathcal{B}' , so by 4.12, $\widetilde{\mathbb{T}}$ is a terminal idempotent \mathcal{V} -monad over \mathbb{T} and $F \dashv G$ factors through $K \dashv J$. Using 3.11 1, we know that $\Sigma_K^{\perp \mathcal{V}} = \mathcal{B}' = \Sigma_F^{\perp \mathcal{V}}$, so since $(\Sigma_K, \Sigma_K^{\perp \mathcal{V}})$ and $(\Sigma_F, \Sigma_F^{\perp \mathcal{V}})$ are \mathcal{V} -orthogonal-pairs by 3.10, we deduce that $\Sigma_{\widetilde{\mathbb{T}}} = \Sigma_K = \Sigma_F = \Sigma_T$. But from this it follows also that $F' : \mathcal{B}' \rightarrow \mathcal{C}$ is conservative, since if $F'f$ is iso (for some morphism f in \mathcal{B}'), then since $F' = FJ$ (4.5) and $\Sigma_F = \Sigma_K$ we find that KJf is iso, but $KJ \cong 1_{\mathcal{B}'}$ and hence f is iso. ■

4.16. PROPOSITION. *Let $\widetilde{\mathbb{T}}$ be an idempotent \mathcal{V} -core of a \mathcal{V} -monad \mathbb{T} on \mathcal{B} . Then the underlying ordinary monad of $\widetilde{\mathbb{T}}$ is an idempotent core of the underlying ordinary monad of \mathbb{T} . Hence, whereas in general, \mathcal{V} -orthogonality implies ordinary orthogonality, we have in this case an equation $\Sigma_T^{\perp \mathcal{V}} = \Sigma_T^{\perp}$.*

PROOF. The underlying ordinary monad of $\widetilde{\mathbb{T}}$ is an idempotent monad which inverts the same morphisms as \mathbb{T} and hence is an ordinary idempotent core of \mathbb{T} . Its associated reflective subcategory is Σ_T^{\perp} and yet has same objects as the \mathcal{V} -reflective-subcategory determined by $\widetilde{\mathbb{T}}$, which is $\Sigma_T^{\perp \mathcal{V}}$. ■

4.17. THEOREM. *Let \mathbb{T} be a \mathcal{V} -monad on a tensored and cotensored \mathcal{V} -category, and suppose that the idempotent core of the underlying ordinary monad of \mathbb{T} exists. Then the idempotent \mathcal{V} -core of \mathbb{T} exists.*

PROOF. By 4.14 and 3.5, Σ_T is closed under tensors in \mathcal{B} , so by 3.6, $\Sigma_T^{\perp \mathcal{V}} = \Sigma_T^{\perp}$. The latter is a reflective subcategory of \mathcal{B} (by an application of 4.15 to the underlying ordinary monad of \mathbb{T}). But $\Sigma_T^{\perp \mathcal{V}}$ is closed under cotensors by 3.5 and hence by 3.12 is a \mathcal{V} -reflective-subcategory of \mathcal{B} , so the result follows by 4.15. ■

4.18. EXAMPLE (Double-dualization monads). Let \mathcal{V} be a symmetric monoidal closed category and $R \in \mathcal{V}$ an object. For each object V of \mathcal{V} , we shall call the internal hom $V^* := \underline{\mathcal{V}}(V, R)$ the *dual* of V with respect to R . We obtain a \mathcal{V} -adjunction

$$\begin{array}{ccc} & (-)^* & \\ \mathcal{V} & \xleftarrow{\quad \mathbb{T} \quad} & \mathcal{V}^{\text{op}} \\ & \xrightarrow{\quad (-)^* \quad} & \end{array}$$

which we call *the dualization \mathcal{V} -adjunction* (for R); it is an instance of a ‘*hom-cotensor*’ \mathcal{V} -adjunction [17, (3.42)]. We call the induced \mathcal{V} -monad \mathbb{T} (on $\underline{\mathcal{V}}$) the *double-dualization*

\mathcal{V} -monad; its underlying \mathcal{V} -functor $(-)^{**}$ sends each $V \in \mathcal{V}$ to the *double-dual* V^{**} of V . Since $\underline{\mathcal{V}}$ is tensored and cotensored, 4.17 entails that the idempotent \mathcal{V} -core $\tilde{\mathbb{T}}$ of \mathbb{T} exists as soon as the idempotent core of the underlying ordinary monad of \mathbb{T} exists. In this case, the \mathcal{V} -reflective-subcategory \mathcal{B}' determined by $\tilde{\mathbb{T}}$ is the \mathcal{V} -reflective-hull of the single object R in $\underline{\mathcal{V}}$. Indeed, by 4.12, \mathcal{B}' is the \mathcal{V} -reflective-hull of $\{\underline{\mathcal{V}}(V, R) \mid V \in \mathcal{V}\}$ in $\underline{\mathcal{V}}$, but any \mathcal{V} -reflective-subcategory of $\underline{\mathcal{V}}$ containing R is closed under cotensors in $\underline{\mathcal{V}}$ and hence contains each cotensor $\underline{\mathcal{V}}(V, R)$ of R .

4.19. EXAMPLE (Completion of normed vector spaces). We shall show in 5 that the double-dualization \mathcal{V} -monad \mathbb{T} on the category \mathcal{V} of normed or seminormed vector spaces (over $R = \mathbb{R}$ or \mathbb{C}) has an idempotent \mathcal{V} -core $\tilde{\mathbb{T}}$ that associates to each (semi-)normed vector space V the (*Cauchy-*)*completion* of V . The \mathcal{V} -reflective-subcategory of $\underline{\mathcal{V}}$ determined by $\tilde{\mathbb{T}}$ is the category of Banach spaces, which therefore is the \mathcal{V} -reflective-hull of R in \mathcal{V} (by 4.18).

4.20. EXAMPLE (Sheafification for a Lawvere-Tierney topology). Given an (elementary) topos \mathcal{X} and a Lawvere-Tierney topology j on \mathcal{X} , let Ω_j be the associated retract of the subobject classifier Ω , and let \mathbb{T} be the double-dualization \mathcal{V} -monad for Ω_j . We show in 8.8 that the idempotent \mathcal{V} -core $\tilde{\mathbb{T}}$ of \mathbb{T} is the *j-sheafification* \mathcal{X} -monad, whose associated \mathcal{X} -reflective-subcategory of $\underline{\mathcal{X}}$ consists of the *j-sheaves*. Hence the \mathcal{X} -category of *j-sheaves* is by 4.18 the \mathcal{X} -reflective-hull of Ω_j in $\underline{\mathcal{X}}$.

5. Example: Completion of normed vector spaces

Let SNorm_1 be the category of seminormed vector spaces over $R = \mathbb{R}$ or \mathbb{C} with nonexpansive linear maps (i.e. bounded linear maps of seminorm ≤ 1), and let Norm_1 be the full subcategory consisting of normed vector spaces.

Letting \mathcal{V} be either SNorm_1 or Norm_1 , it is well-known that \mathcal{V} is symmetric monoidal closed; e.g. see [2] §3.4. Indeed, in both categories, the internal hom $\underline{\mathcal{V}}(V, W)$ ($V, W \in \mathcal{V}$) is the vector space of all bounded linear maps $V \rightarrow W$, equipped with the usual *operator (semi)norm*. Given seminormed (resp. normed) spaces V, W , the monoidal product $V \otimes W$ in SNorm_1 (resp. Norm_1) is the algebraic tensor product, equipped with the *projective seminorm* (resp. norm) $\|x\| = \inf\{\sum_i \|v_i\| \|w_i\| \mid x = \sum_{i=1}^n v_i \otimes w_i\}$. (In particular, the projective seminorm is a norm as soon as V and W are normed [24, Ch. III, Exercise 20].)

The *dualization* \mathcal{V} -functor $(-)^* : \underline{\mathcal{V}}^{\text{op}} \rightarrow \underline{\mathcal{V}}$ (4.18) associates to each $V \in \mathcal{V}$ the space V^* of all bounded linear functionals on V , and assigns to each morphism $h : V_1 \rightarrow V_2$ in \mathcal{V} the map $h : V_2^* \rightarrow V_1^*$ given by $\psi \mapsto \psi \cdot h$. The *double-dualization* \mathcal{V} -monad \mathbb{T} (4.18) on $\underline{\mathcal{V}}$ associates to each $V \in \mathcal{V}$ its *double-dual* $TV = V^{**}$, and the unit morphism $\eta_V : V \rightarrow TV$ is the familiar canonical linear map, which is always *isometric* (i.e. $\|\eta_V(v)\| = \|v\|$ for all $v \in V$), so that η_V is an *isometric embedding* (i.e., isometric and injective) as soon as V is normed.

In the present section, we show that the double-dualization \mathcal{V} -monad \mathbb{T} on $\underline{\mathcal{V}}$ has an idempotent \mathcal{V} -core $\widetilde{\mathbb{T}}$, given by *completion*.

5.1. The full subcategory Ban_1 of \mathcal{V} ($= \text{SNorm}_1$ or Norm_1) consisting of all Banach spaces is a \mathcal{V} -reflective-subcategory of $\underline{\mathcal{V}}$. Indeed, for an arbitrary morphism $\rho_V : V \rightarrow \widetilde{V}$ in \mathcal{V} with \widetilde{V} a Banach space, the following conditions are equivalent, and they characterize (up to isomorphism) the familiar *completion* \widetilde{V} of V :

1. $\rho_V : V \rightarrow \widetilde{V}$ is dense and isometric.
2. For each Banach space B , the morphism $\underline{\mathcal{V}}(\rho_V, B) : \underline{\mathcal{V}}(\widetilde{V}, B) \rightarrow \underline{\mathcal{V}}(V, B)$ in \mathcal{V} is an (isometric) isomorphism.
3. For each morphism $f : V \rightarrow B$ in \mathcal{V} with B a Banach space, there is a unique morphism $f^\# : \widetilde{V} \rightarrow B$ in \mathcal{V} with $f^\# \cdot \rho_V = f$; further, $\|f^\#\| = \|f\|$.

Concretely, we can take \widetilde{V} to be the familiar *Cauchy-completion* of V , or the closure of the image of $\eta_V : V \rightarrow TV = V^{**}$.

5.2. LEMMA. *Let $h : V \rightarrow W$ in \mathcal{V} ($= \text{SNorm}_1$ or Norm_1). Then h is dense if and only if $h^* : W^* \rightarrow V^*$ is injective.*

PROOF. The ‘only if’ part is straightforward. Conversely, suppose h is not dense. Then, letting $C \subseteq W$ be the closure of the image of h , the Hahn-Banach Extension Theorem entails that there is some nonzero $\psi \in W^*$ with $\psi|_C = 0$. But then $h^*(\psi) = \psi \cdot h = 0$ and yet $\psi \neq 0$, showing that h^* is not injective. ■

5.3. PROPOSITION. *For each morphism $h : V \rightarrow W$ in \mathcal{V} ($= \text{SNorm}_1$ or Norm_1), the following are equivalent:*

1. h is dense and isometric.
2. $\widetilde{h} : \widetilde{V} \rightarrow \widetilde{W}$ is an (isometric) isomorphism.
3. $h^* : W^* \rightarrow V^*$ is an (isometric) isomorphism.

PROOF. To show $1 \Rightarrow 2$, suppose that h is dense and isometric. We have a commutative square

$$\begin{array}{ccc}
 V & \xrightarrow{\rho_V} & \widetilde{V} \\
 h \downarrow & & \downarrow \widetilde{h} \\
 W & \xrightarrow{\rho_W} & \widetilde{W}
 \end{array} \tag{5.3.i}$$

in which both h and ρ_W are dense and isometric, so that the composite $\rho_W \cdot h : V \rightarrow \widetilde{W}$ is a dense, isometric morphism into a Banach space and hence satisfies the universal property characterizing the completion of V (5.1). Using this and also the (same) universal property of ρ_V , it follows that \widetilde{h} is an isomorphism.

To show $2 \Rightarrow 3$, suppose that \tilde{h} is an isomorphism. Then, applying $(-)^*$ to the commutative square (5.3.i), we obtain a commutative square

$$\begin{array}{ccc} V^* & \xleftarrow{\rho_V^*} & \tilde{V}^* \\ \uparrow h^* & & \uparrow \tilde{h}^* \\ W^* & \xleftarrow{\rho_W^*} & \tilde{W}^* \end{array}$$

in which the right side is an isomorphism. But the second characterization of the completion in 5.1 entails that the top and bottom faces are isomorphisms as well, so h^* is an isomorphism.

To show $3 \Rightarrow 1$, suppose that h^* is an isomorphism. By 5.2 we deduce that h is dense. Further, we have a commutative square

$$\begin{array}{ccc} V & \xrightarrow{\eta_V} & V^{**} \\ \downarrow h & & \downarrow h^{**} \\ W & \xrightarrow{\eta_W} & W^{**} \end{array}$$

in which the right side is an isomorphism and the top and bottom sides are isometric, and it follows that h is isometric. ■

5.4. Let $\tilde{\mathbb{T}}$ be the idempotent \mathcal{V} -monad induced by the completion \mathcal{V} -adjunction $(-)^{\rho} \dashv J : \text{Ban}_1 \hookrightarrow \underline{\mathcal{V}}$, where $\mathcal{V} = \text{SNorm}_1$ or Norm_1 , so that $\tilde{T}V = \tilde{V}$ is the completion of $V \in \mathcal{V}$.

5.5. THEOREM. *The completion \mathcal{V} -monad $\tilde{\mathbb{T}}$ on the category \mathcal{V} of normed (resp. seminormed) vector spaces is the idempotent \mathcal{V} -core of the double-dualization \mathcal{V} -monad \mathbb{T} on $\underline{\mathcal{V}}$. Moreover, $\Sigma_T = \Sigma_{\tilde{T}}$ is the class of all dense, isometric morphisms in \mathcal{V} .*

PROOF. By 4.13, $\Sigma_T = \Sigma_{(-)^*}$, and by 5.3, the latter class equals $\Sigma_{\tilde{T}}$ and consists of exactly the dense, isometric morphisms. ■

5.6. COROLLARY. *The category Ban_1 of Banach spaces is the \mathcal{V} -reflective-hull of $R (= \mathbb{R}$ or $\mathbb{C})$ in the category \mathcal{V} of normed (resp. seminormed) vector spaces. Moreover, a normed (resp. seminormed) vector space B is a Banach space if and only if the following equivalent conditions hold for each morphism $h : V \rightarrow W$ in \mathcal{V} :*

1. *If $h \perp_{\mathcal{V}} R$, then $h \perp_{\mathcal{V}} B$.*
2. *If $h^* : W^* \rightarrow V^*$ is an isomorphism (in \mathcal{V}), then $\underline{\mathcal{V}}(h, B) : \underline{\mathcal{V}}(W, B) \rightarrow \underline{\mathcal{V}}(V, B)$ is an isomorphism.*

6. Completion, closure, and density relative to a monad. Existence results via factorization.

6.1. GIVEN DATA. In the present section, we work with given data as follows, which we shall later suppose to satisfy Assumption 6.4 below:

1. Let $\mathbb{T} = (T, \eta, \mu)$ be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} .
2. Let $\Sigma \subseteq \Sigma_T$ be a class of morphisms inverted by T .
3. Let $(\mathcal{E}, \mathcal{M})$ be a \mathcal{V} -proper \mathcal{V} -prefactorization-system on \mathcal{B} .

We shall refer to the morphisms in \mathcal{M} as \mathcal{M} -embeddings.

6.2. DEFINITION. Given data as in 6.1, we make the following definitions:

1. An object B of \mathcal{B} is Σ -complete if $B \in \mathcal{B}_\Sigma$.
2. An object B of \mathcal{B} is \mathbb{T} -separated if $\eta_B : B \rightarrow TB$ is an \mathcal{M} -embedding.
3. We denote the full sub- \mathcal{V} -category of \mathcal{B} consisting of the Σ -complete \mathbb{T} -separated objects by $\mathcal{B}_{(\mathbb{T}, \Sigma)}$.
4. We say that an \mathcal{M} -embedding $m : B_1 \rightarrow B_2$ in \mathcal{B} is Σ -closed if $m \in \Sigma^{\downarrow \mathcal{V}}$. We denote by $\Sigma\text{-ClEmb} := \Sigma^{\downarrow \mathcal{V}} \cap \mathcal{M}$ the class of all Σ -closed \mathcal{M} -embeddings in \mathcal{B} .
5. We say that a morphism in \mathcal{B} is Σ -dense if it lies in $\Sigma\text{-Dense} := \Sigma\text{-ClEmb}^{\uparrow \mathcal{V}}$.
6. If $\Sigma = \Sigma_T$, then we replace the prefixes “ Σ -” with “ \mathbb{T} -”, obtaining the notions of \mathbb{T} -complete object, \mathbb{T} -closed \mathcal{M} -embedding, and \mathbb{T} -dense morphism.

6.3. REMARK. Observe that $(\Sigma\text{-Dense}, \Sigma\text{-ClEmb})$ is a \mathcal{V} -prefactorization-system on \mathcal{B} , since $\Sigma\text{-ClEmb} = \Sigma^{\downarrow \mathcal{V}} \cap \mathcal{E}^{\downarrow \mathcal{V}} = (\Sigma \cup \mathcal{E})^{\downarrow \mathcal{V}}$.

6.4. ASSUMPTION. For the remainder of §6, we shall assume that *every morphism in \mathcal{B} factors as a Σ -dense morphism followed by a Σ -closed \mathcal{M} -embedding*.

It then follows by 6.3 that $(\Sigma\text{-Dense}, \Sigma\text{-ClEmb})$ is a \mathcal{V} -factorization-system on \mathcal{B} .

6.5. EXAMPLE. Given data as in 6.1, Assumption 6.4 is satisfied as soon as \mathcal{B} is \mathcal{M} -subobject-complete (2.3.2).

6.6. **EXAMPLE** (Normed and seminormed spaces). As we shall show in 7.2, data satisfying Assumption 6.4 can be obtained as follows. Take $\mathcal{V} := \text{Norm}_1$ or SNorm_1 to be the category of normed or seminormed vector spaces (5), let $\mathcal{B} := \underline{\mathcal{V}}$, let \mathcal{E} consist of all surjective morphisms in \mathcal{V} , and let \mathcal{M} consist of all isometric embeddings in \mathcal{V} . Let \mathbb{T} be the double-dualization \mathcal{V} -monad on $\underline{\mathcal{V}}$ (5), and let $\Sigma := \Sigma_T$. Then

1. a (semi)normed space is \mathbb{T} -complete if and only if it is a Banach space;
2. a seminormed space is \mathbb{T} -separated if and only if it is normed;
3. the \mathbb{T} -closed \mathcal{M} -embeddings are exactly the closed isometric embeddings;
4. the \mathbb{T} -dense morphisms are exactly the dense morphisms in the usual sense;
5. Σ_T consists of exactly the dense isometric morphisms in \mathcal{V} .

6.7. **EXAMPLE** (Sheaves, density, and closed subobjects). Given an (elementary) topos \mathcal{X} and a Lawvere-Tierney topology j on \mathcal{X} , we show in 8.8 that data satisfying Assumption 6.4 can be obtained as follows. Take $\mathcal{V} := \mathcal{X}$, let $\mathcal{B} := \underline{\mathcal{X}}$, and let $(\mathcal{E}, \mathcal{M}) := (\text{Epi } \mathcal{X}, \text{Mono } \mathcal{X})$. Let \mathbb{T} be the double-dualization \mathcal{X} -monad for Ω_j , and let $\Sigma := \Sigma_T$. Then

1. an object of \mathcal{X} is \mathbb{T} -complete if and only if it is a j -sheaf;
2. an object of \mathcal{X} is \mathbb{T} -separated if and only if it is j -separated;
3. the \mathbb{T} -closed \mathcal{M} -embeddings are exactly the j -closed monomorphisms;
4. the \mathbb{T} -dense morphisms are exactly the j -dense morphisms.

Also, given a Grothendieck quasitopos \mathcal{Y} , so that (up to equivalence of categories) \mathcal{Y} is the category of K -separated J -sheaves on a *bisite* (\mathcal{C}, J, K) , we show in 8.9 that data satisfying Assumption 6.4 can be obtained by considering the associated Lawvere-Tierney topologies j, k on the presheaf topos $\mathcal{X} := [\mathcal{C}^{\text{op}}, \text{Set}]$, taking \mathbb{T} to be the double-dualization \mathcal{X} -monad for Ω_k , and letting Σ consist of all morphisms inverted by the double-dualization functor for Ω_j , so that $\underline{\mathcal{X}}_{(\mathbb{T}, \Sigma)} = \mathcal{Y}$.

6.8. **PROPOSITION.**

1. Every morphism $e \in \mathcal{E}$ is Σ -dense.
2. Every morphism in Σ is Σ -dense.
3. If a composite $B_1 \xrightarrow{f} B_2 \xrightarrow{g} B_3$ is Σ -dense, then its second factor g is Σ -dense.
4. If a composite $B_1 \xrightarrow{f} B_2 \xrightarrow{g} B_3$ and its second factor g are both Σ -closed \mathcal{M} -embeddings, then the first factor f is a Σ -closed \mathcal{M} -embedding.

- 5. Every Σ -dense Σ -closed \mathcal{M} -embedding is an isomorphism.
- 6. Σ -CIEmb is closed under composition, cotensors, arbitrary \mathcal{V} -intersections, and \mathcal{V} -pullbacks along arbitrary morphisms in \mathcal{B} .
- 7. Σ -Dense is closed under composition, tensors, arbitrary \mathcal{V} -cofibre-coproducts, and \mathcal{V} -pushouts along arbitrary morphisms in \mathcal{B} .

PROOF. 1. e is \mathcal{V} -orthogonal to every \mathcal{M} -embedding and hence to every Σ -closed \mathcal{M} -embedding. 2. Σ -CIEmb $\subseteq \Sigma^{\downarrow\mathcal{V}}$, so $\Sigma \subseteq \Sigma^{\downarrow\mathcal{V}\uparrow\mathcal{V}} \subseteq \Sigma$ -CIEmb $^{\uparrow\mathcal{V}} = \Sigma$ -Dense. 3-7 follow from 2.3.1. ■

6.9. Since $(\Sigma$ -Dense, Σ -CIEmb) is a \mathcal{V} -factorization-system and hence an ordinary factorization system (by [21, 5.3]), it determines a weakly hereditary idempotent closure operator $\overline{(-)}$ on \mathcal{M} in \mathcal{B} (2.4.3). For each \mathcal{M} -embedding m , we call \overline{m} the Σ -closure of m ; in the case that $\Sigma = \Sigma_T$, we call \overline{m} the \mathbb{T} -closure of m .

6.10. PROPOSITION. Let $m : B' \rightarrow B$ be an \mathcal{M} -embedding in \mathcal{B} , and suppose B is Σ -complete. Then

$$B' \text{ is } \Sigma\text{-complete} \iff m \text{ is } \Sigma\text{-closed}.$$

PROOF. For each $h : B_1 \rightarrow B_2$ in Σ , we have a commutative square

$$\begin{array}{ccc} \mathcal{B}(B_2, B') & \xrightarrow{\mathcal{B}(h, B')} & \mathcal{B}(B_1, B') \\ \mathcal{B}(B_2, m) \downarrow & & \downarrow \mathcal{B}(B_1, m) \\ \mathcal{B}(B_2, B) & \xrightarrow{\mathcal{B}(h, B)} & \mathcal{B}(B_1, B) \end{array}$$

in which $\mathcal{B}(h, B)$ is iso, so the square is a pullback if and only if $\mathcal{B}(h, B')$ is iso. ■

For the remainder of this section, let us fix a \mathcal{V} -adjunction $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ inducing \mathbb{T} ; for example, one can take $F \dashv G$ to be the Kleisli \mathcal{V} -adjunction.

6.11. PROPOSITION. Let C be an object of \mathcal{C} . Then GC is Σ -complete.

PROOF. For each $h : B_1 \rightarrow B_2$ in Σ , we have a commutative square

$$\begin{array}{ccc} \mathcal{B}(B_2, GC) & \xrightarrow{\mathcal{B}(h, GC)} & \mathcal{B}(B_1, GC) \\ \wr \downarrow & & \downarrow \wr \\ \mathcal{C}(FB_2, C) & \xrightarrow{\mathcal{C}(Fh, C)} & \mathcal{C}(FB_1, C) \end{array}$$

whose left and right sides are isomorphisms, and since $h \in \Sigma \subseteq \Sigma_T = \Sigma_F$ by 4.13, the bottom side is iso, so the top side is iso. ■

6.12. PROPOSITION. *Let $B \in \mathcal{B}$ be \mathbb{T} -separated. Then*

$$B \text{ is } \Sigma\text{-complete} \Leftrightarrow \eta_B : B \rightarrow TB \text{ is } \Sigma\text{-closed}.$$

PROOF. Since $TB = GFB$ is Σ -complete by 6.11 and η_B is an \mathcal{M} -embedding, this follows from 6.10. ■

6.13. PROPOSITION. *An object $B \in \mathcal{B}$ is \mathbb{T} -separated (resp. Σ -complete and \mathbb{T} -separated) iff there exists an \mathcal{M} -embedding (resp. Σ -closed \mathcal{M} -embedding) $m : B \rightarrow GC$ for some $C \in \mathcal{C}$.*

PROOF. If B is \mathbb{T} -separated, then $\eta_B : B \rightarrow GFB$ is an \mathcal{M} -embedding; if B is also Σ -complete, then by 6.12, η_B is Σ -closed. Conversely, if $m : B \rightarrow GC$ is an \mathcal{M} -embedding then we have a commutative triangle

$$\begin{array}{ccc} B & \xrightarrow{\eta_B} & GFB \\ & \searrow m & \downarrow Gm^\sharp \\ & & GC \end{array}$$

for a unique morphism m^\sharp in \mathcal{C} , so $\eta_B \in \mathcal{M}$ by 2.3.1 (since $\mathcal{C} \subseteq \text{Epi}_\gamma \mathcal{B}$), so B is \mathbb{T} -separated. If the given \mathcal{M} -embedding m is also Σ -closed, then since GC is Σ -complete by 6.11, we deduce by 6.10 that B is Σ -complete. ■

6.14. COROLLARY. *For each $B \in \mathcal{B}$, TB is Σ -complete and \mathbb{T} -separated.*

PROOF. $1_{TB} : TB \rightarrow TB = GFB$ is a Σ -closed \mathcal{M} -embedding. ■

6.15. DEFINITION. For each $B \in \mathcal{B}$, let

$$\begin{array}{ccc} B & \xrightarrow{\eta_B} & TB \\ & \searrow \rho_B & \nearrow \iota_B \\ & & KB \end{array}$$

be the $(\Sigma\text{-Dense}, \Sigma\text{-CIEmb})$ -factorization of η_B .

6.16. PROPOSITION. *For each $B \in \mathcal{B}$, KB is Σ -complete and \mathbb{T} -separated.*

PROOF. Since $TB = GFB$, this follows from 6.13. ■

The following lemma was inspired by an idea employed in the proof of 3.3 of [5] in the non-enriched context.

6.17. LEMMA. *Let $f : B_1 \rightarrow B_2$ be a Σ -dense morphism for which $Ff : FB_1 \rightarrow FB_2$ is a section. Then Ff is iso.*

PROOF. The periphery of the following diagram commutes.

$$\begin{array}{ccc}
 B_1 & \xrightarrow{\eta_{B_1}} & GF B_1 \\
 f \downarrow & \nearrow k & \downarrow GF f \\
 B_2 & \xrightarrow{\eta_{B_2}} & GF B_2
 \end{array}$$

Also, GFf is a section and hence is an \mathcal{M} -embedding (by 2.3.3). Further, by 3.10, we have that $GFf \in \Sigma_F^{\downarrow\gamma} \subseteq \Sigma^{\downarrow\gamma}$ (using the fact that $\Sigma \subseteq \Sigma_T = \Sigma_F$), so GFf is a Σ -closed \mathcal{M} -embedding. Hence, since f is Σ -dense, there is a unique morphism k making the above diagram commute. In particular, $GFf \cdot k = \eta_{B_2}$; taking the transposes of both sides of this equation, with respect to the adjunction $F \dashv G$, we find that $Ff \cdot k^\sharp = 1_{FB_2}$ where $k^\sharp : FB_2 \rightarrow FB_1$ is the transpose of k . Hence Ff is a split epi and hence, being also a split mono, is iso. ■

6.18. PROPOSITION. For each $B \in \mathcal{B}$, $\rho_B : B \rightarrow KB$ is inverted by F .

PROOF. Taking the transposes of each side of the equation

$$\left(B \xrightarrow{\rho_B} KB \xrightarrow{\iota_B} GF B \right) = \eta_B$$

under the adjunction $F \xrightarrow{\eta} \varepsilon | G$, we obtain

$$\left(FB \xrightarrow{F\rho_B} FKB \xrightarrow{F\iota_B} FGF B \xrightarrow{\varepsilon_{FB}} FB \right) = 1_{FB},$$

so $F\rho_B$ is a section, so since ρ_B is Σ -dense, 6.17 applies, and we deduce that $F\rho_B$ is iso. ■

6.19. PROPOSITION. Let $B, B' \in \mathcal{B}$ and suppose B' is \mathbb{T} -separated and Σ -complete. Then $\rho_B \perp_{\gamma} B'$.

PROOF. We have a commutative diagram as follows.

$$\begin{array}{ccc}
 \mathcal{B}(KB, B') & \xrightarrow{\mathcal{B}(\rho_B, B')} & \mathcal{B}(B, B') \\
 \mathcal{B}(KB, \eta_{B'}) \downarrow & & \downarrow \mathcal{B}(B, \eta_{B'}) \\
 \mathcal{B}(KB, GF B') & \xrightarrow{\mathcal{B}(\rho_B, GF B')} & \mathcal{B}(B, GF B') \\
 \downarrow \wr & & \downarrow \wr \\
 \mathcal{C}(FKB, FB') & \xrightarrow{\mathcal{C}(F\rho_B, FB')} & \mathcal{C}(FB, FB')
 \end{array}$$

Since B' is \mathbb{T} -separated and Σ -complete, we have by 6.12 that $\eta_{B'}$ is a Σ -closed \mathcal{M} -embedding, so since ρ_B is Σ -dense, $\rho_B \downarrow_{\gamma} \eta_{B'}$, so the upper square is a pullback. Also, $F\rho_B$ is iso by 6.18, so the left, bottom, and right sides of the lower square are iso. Therefore $\mathcal{B}(\rho_B, GF B')$ is iso and hence its pullback $\mathcal{B}(\rho_B, B')$ is iso. ■

6.20. THEOREM. Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} equipped with a \mathcal{V} -proper \mathcal{V} -prefactorization-system $(\mathcal{E}, \mathcal{M})$, and let $\Sigma \subseteq \Sigma_{\mathbb{T}}$. Suppose that every morphism in \mathcal{B} factors as a Σ -dense morphism followed by a Σ -closed \mathcal{M} -embedding. Then the morphisms $\rho_B : B \rightarrow KB$ ($B \in \mathcal{B}$) of 6.15 exhibit the \mathcal{V} -category $\mathcal{B}_{(\mathbb{T}, \Sigma)}$ of Σ -complete \mathbb{T} -separated objects (6.2) as a \mathcal{V} -reflective-subcategory of \mathcal{B} .

PROOF. For each $B \in \mathcal{B}$ we have by 6.16 that KB lies in $\mathcal{B}_{(\mathbb{T}, \Sigma)}$, and for each $B' \in \mathcal{B}_{(\mathbb{T}, \Sigma)}$,

$$\mathcal{B}_{(\mathbb{T}, \Sigma)}(KB, B') = \mathcal{B}(KB, B') \xrightarrow{\mathcal{B}(\rho_B, B')} \mathcal{B}(B, B')$$

is iso by 6.19. The result follows by 2.2.4. ■

6.21. REMARK. The hypothesis in 6.20 that $(\Sigma\text{-Dense}, \Sigma\text{-CIEmb})$ -factorizations exist (i.e., Assumption 6.4) is satisfied as soon as \mathcal{B} is \mathcal{M} -subobject-complete (2.3.2).

6.22. DEFINITION. Given data satisfying the hypotheses of 6.20, we call the idempotent \mathcal{V} -monad $\tilde{\mathbb{T}}_{\Sigma}$ on \mathcal{B} induced by the resulting \mathcal{V} -reflection $K \dashv \mid J : \mathcal{B}_{(\mathbb{T}, \Sigma)} \hookrightarrow \mathcal{B}$ the \mathbb{T} -separated Σ -completion \mathcal{V} -monad.

6.23. PROPOSITION. Suppose given data satisfying the hypotheses of 6.20.

1. There is a unique morphism of \mathcal{V} -monads $\iota : \tilde{\mathbb{T}}_{\Sigma} \rightarrow \mathbb{T}$.
2. Each component of ι is a Σ -closed \mathcal{M} -embedding.
3. Each component of the unit $\rho : 1_{\mathcal{B}} \rightarrow \tilde{\mathbb{T}}_{\Sigma}$ of $\tilde{\mathbb{T}}_{\Sigma}$ is a Σ -dense morphism.
4. Each \mathcal{V} -adjunction inducing \mathbb{T} factors through the \mathcal{V} -reflection $K \dashv \mid J : \mathcal{B}_{(\mathbb{T}, \Sigma)} \hookrightarrow \mathcal{B}$ determined by $\tilde{\mathbb{T}}_{\Sigma}$.

PROOF. 1 and 4 follow from 6.14 and 4.7. By 4.9 and 6.15, the components of the resulting morphism of \mathcal{V} -monads are necessarily the Σ -closed \mathcal{M} -embeddings ι_B given in 6.15, so 2 holds. 3 is immediate. ■

6.24. COROLLARY. Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} equipped with a \mathcal{V} -proper \mathcal{V} -factorization-system $(\mathcal{E}, \mathcal{M})$. Then the full sub- \mathcal{V} -category of \mathcal{B} consisting of the \mathbb{T} -separated objects is \mathcal{V} -reflective in \mathcal{B} .

PROOF. Taking $\Sigma = \emptyset$, the objects of $\mathcal{B}_{(\mathbb{T}, \Sigma)}$ are exactly the \mathbb{T} -separated objects of \mathcal{B} , and the hypotheses of 6.20 are satisfied since $\emptyset\text{-CIEmb} = \mathcal{M}$ and hence $\emptyset\text{-Dense} = \mathcal{E}$. ■

6.25. THEOREM. Let \mathbb{T} be a \mathcal{V} -monad on a \mathcal{V} -category \mathcal{B} equipped with a \mathcal{V} -proper \mathcal{V} -factorization-system $(\mathcal{E}, \mathcal{M})$, and suppose that every morphism in \mathcal{B} factors as a \mathbb{T} -dense morphism followed by a \mathbb{T} -closed \mathcal{M} -embedding.

1. The idempotent \mathcal{V} -core $\tilde{\mathbb{T}}$ of \mathbb{T} exists.
2. The \mathcal{V} -reflective-subcategory of \mathcal{B} determined by $\tilde{\mathbb{T}}$ consists of the \mathbb{T} -complete objects.
3. Every \mathbb{T} -complete object of \mathcal{B} is \mathbb{T} -separated.
4. Each component of the unique morphism of \mathcal{V} -monads $\iota : \tilde{\mathbb{T}} \rightarrow \mathbb{T}$ is a \mathbb{T} -closed \mathcal{M} -embedding.
5. Each component of the unit $\rho : 1_{\mathcal{B}} \rightarrow \tilde{T}$ of $\tilde{\mathbb{T}}$ is a \mathbb{T} -dense morphism.
6. Every \mathcal{V} -adjunction $F \dashv G : \mathcal{C} \rightarrow \mathcal{B}$ inducing \mathbb{T} factors through the \mathcal{V} -reflection $K \dashv J : \mathcal{B}_{\Sigma_T} \hookrightarrow \mathcal{B}$ determined by $\tilde{\mathbb{T}}$, in such a way that the induced left \mathcal{V} -adjoint $F' : \mathcal{B}_{\Sigma_T} \rightarrow \mathcal{C}$ (4.6) is conservative.

PROOF. 3. By 6.24 we know that the full sub- \mathcal{V} -category $\mathcal{B}_{(\mathbb{T}, \emptyset)}$ of \mathcal{B} consisting of the \mathbb{T} -separated objects is \mathcal{V} -reflective in \mathcal{B} , and we will denote the components of the unit of the associated \mathcal{V} -reflection by $\sigma_B : B \rightarrow LB$ ($B \in \mathcal{B}$). Hence, σ_B is gotten as the morphism ρ_B of 6.15 in the case that $\Sigma = \emptyset$. By 6.18 we know that each such component σ_B is inverted by F — i.e. $\sigma_B \in \Sigma_F = \Sigma_T$. Hence, given any \mathbb{T} -complete object $B' \in \mathcal{B}_{\Sigma_T}$, we have that $\sigma_B \perp_{\mathcal{V}} B'$ for every $B \in \mathcal{B}$, so by 3.11, $B' \in \mathcal{B}_{(\mathbb{T}, \emptyset)}$.

By 3 we know that $\mathcal{B}_{\Sigma_T} = \mathcal{B}_{(\mathbb{T}, \Sigma_T)}$, and by 6.20 we deduce that $\mathcal{B}_{(\mathbb{T}, \Sigma_T)}$ is a \mathcal{V} -reflective-subcategory of \mathcal{B} . Hence 1, 2, and 6 follow immediately from 4.15.

4 and 5 follow from 6.23. ■

6.26. COROLLARY. Let \mathbb{T} be a \mathcal{V} -monad on an \mathcal{M} -subobject-complete \mathcal{V} -category \mathcal{B} , where $(\mathcal{E}, \mathcal{M})$ is a \mathcal{V} -proper \mathcal{V} -prefactorization-system on \mathcal{B} . Then the idempotent \mathcal{V} -core $\tilde{\mathbb{T}}$ of \mathbb{T} exists.

PROOF. The hypotheses of 6.25 are satisfied (2.3.2, 6.5). ■

In view of 6.25, we shall extend the notation and terminology of 6.2 as follows:

6.27. DEFINITION. Let data satisfying the hypotheses of 6.25 be given.

1. We call $\tilde{\mathbb{T}}$ the \mathbb{T} -completion \mathcal{V} -monad.
2. For each object $B \in \mathcal{B}$, we call $\tilde{T}B = KB$ the \mathbb{T} -completion of B .
3. We denote by $\mathcal{B}_{(\mathbb{T})} := \mathcal{B}_{\Sigma_T} = \mathcal{B}_{(\mathbb{T}, \Sigma_T)}$ the \mathcal{V} -reflective-subcategory of \mathcal{B} consisting of the \mathbb{T} -complete objects.

7. Example: Closure and density in normed spaces

We saw in 5.5 that the completion \mathcal{V} -monad on the category \mathcal{V} of normed (resp. seminormed) vector spaces is the idempotent \mathcal{V} -core of the double-dualization \mathcal{V} -monad on $\underline{\mathcal{V}}$. Further, we shall see that the familiar construction of the completion of a normed or seminormed vector space V as the closure of V in V^{**} is an instance of the general procedure given in 6.25 for forming the idempotent \mathcal{V} -core $\widetilde{\mathbb{T}}$ of a \mathcal{V} -monad \mathbb{T} . Indeed, we will show that the notions of \mathbb{T} -closure and \mathbb{T} -density in this example coincide with the familiar notions of closure and density.

Again as in 5, let \mathcal{V} be either SNorm_1 or Norm_1 , and let \mathbb{T} denote the double-dualization \mathcal{V} -monad on $\underline{\mathcal{V}}$. Further, let \mathcal{E} denote the class of all surjective morphisms in \mathcal{V} , and let \mathcal{M} denote the class of all isometric embeddings in \mathcal{V} .

7.1. LEMMA. *$(\mathcal{E}, \mathcal{M})$ is a \mathcal{V} -proper \mathcal{V} -factorization-system on the \mathcal{V} -category $\underline{\mathcal{V}}$ of normed (resp. seminormed) vector spaces.*

PROOF. Each surjective morphism (resp. each isometric embedding) is clearly an epimorphism (resp. a monomorphism) in \mathcal{V} and hence, by [21, 2.4], is a \mathcal{V} -epi (resp. a \mathcal{V} -mono) in $\underline{\mathcal{V}}$. Since $\underline{\mathcal{V}}$ is a cotensored \mathcal{V} -category, it suffices by [21, 5.7] to show that $(\mathcal{E}, \mathcal{M})$ is an ordinary factorization system on \mathcal{V} and that \mathcal{M} is closed under cotensors in $\underline{\mathcal{V}}$. Clearly every morphism in \mathcal{V} factors as a morphism in \mathcal{E} followed by a morphism in \mathcal{M} . Moreover, it is easy to check that each morphism $e \in \mathcal{E}$ is orthogonal to each morphism $m \in \mathcal{M}$. Hence, since \mathcal{E} and \mathcal{M} are also closed under composition with isomorphisms, we deduce that $(\mathcal{E}, \mathcal{M})$ is a factorization system on \mathcal{V} (e.g., by [21, 5.2]). Further \mathcal{M} is closed under cotensors in \mathcal{V} , since for any isometric embedding $m : W_1 \rightarrow W_2$ in \mathcal{V} , it is readily verified that the induced morphism $\underline{\mathcal{V}}(V, m) : \underline{\mathcal{V}}(V, W_1) \rightarrow \underline{\mathcal{V}}(V, W_2)$ (given by composing with m) is injective and isometric. ■

7.2. THEOREM. *With respect to the double-dualization \mathcal{V} -monad \mathbb{T} on the category \mathcal{V} of normed (resp. seminormed) vector spaces and the \mathcal{V} -factorization-system $(\mathcal{E}, \mathcal{M})$, the following hold:*

1. *A morphism in \mathcal{V} is \mathbb{T} -dense if and only if it is dense.*
2. *A morphism in \mathcal{V} is a \mathbb{T} -closed \mathcal{M} -embedding if and only if it is a closed isometric embedding.*
3. *Every morphism in \mathcal{V} factors as a \mathbb{T} -dense morphism followed by a \mathbb{T} -closed \mathcal{M} -embedding.*
4. *The \mathbb{T} -closure of a subspace $V \hookrightarrow W$ of a normed (resp. seminormed) vector space W is the usual closure $\overline{V} \hookrightarrow W$ of V in W .*

PROOF. One sees immediately that the familiar process of taking the closure of a subspace defines a weakly hereditary idempotent closure operator $\overline{(-)}$ on \mathcal{M} in \mathcal{V} (2.4.1). By 2.4.2, $\overline{(-)}$ determines an associated factorization system $(\text{Dense}, \text{CIEmb})$, consisting of the dense morphisms and closed \mathcal{M} -embeddings, respectively.

Recall from 5.5 that the class Σ_T of all morphisms inverted by $T = (-)^{**}$ consists of exactly the dense isometric morphisms. By definition, an \mathcal{M} -embedding m is \mathbb{T} -closed iff $m \in \Sigma_T^\perp$. But since $(\Sigma_T, \Sigma_T^\perp)$ is a \mathcal{V} -orthogonal-pair in $\underline{\mathcal{V}}$ (4.14), we deduce by 3.5 that Σ_T is closed under tensors in $\underline{\mathcal{V}}$, so by [21, 5.4], $\Sigma_T^\perp = \Sigma_T^\perp$, whence $\mathbb{T}\text{-CIEmb} = \Sigma_T^\perp \cap \mathcal{M}$.

To prove 2, first observe that since $\Sigma_T \subseteq \text{Dense}$, $\text{CIEmb} = \text{Dense}^\perp \subseteq \Sigma_T^\perp$ and hence $\text{CIEmb} \subseteq \mathbb{T}\text{-CIEmb}$. Also, by 2.4.4 we know that $\text{CIEmb} = \text{DenseEmb}^\perp \cap \mathcal{M}$, where DenseEmb is the set of all dense isometric embeddings, so that since $\text{DenseEmb} \subseteq \Sigma_T$, $\Sigma_T^\perp \subseteq \text{DenseEmb}^\perp$ and hence $\mathbb{T}\text{-CIEmb} \subseteq \text{CIEmb}$.

Since both Σ_T^\perp and \mathcal{M} are closed under cotensors in $\underline{\mathcal{V}}$ (by 2.3.1 and 4.14), their intersection $\mathbb{T}\text{-CIEmb} = \text{CIEmb}$ is closed under cotensors in $\underline{\mathcal{V}}$, so the factorization system $(\text{Dense}, \text{CIEmb})$ is a \mathcal{V} -factorization-system, by [21, 5.7]. Hence 1 and 3 follow, since $\text{Dense} = \text{CIEmb}^{\uparrow\mathcal{V}} = (\mathbb{T}\text{-CIEmb})^{\uparrow\mathcal{V}} = \mathbb{T}\text{-Dense}$. Further, 4 follows as well, since the \mathbb{T} -closure operator is by definition the closure operator determined by $(\mathbb{T}\text{-Dense}, \mathbb{T}\text{-CIEmb}) = (\text{Dense}, \text{CIEmb})$ (2.4.3), which coincides with $\overline{(-)}$. ■

7.3. REMARK. By 7.1 and 7.2 3, the data $\mathbb{T}, (\mathcal{E}, \mathcal{M})$ of 7.2 satisfy the hypotheses of 6.25, and the latter theorem entails that the idempotent \mathcal{V} -core $\widetilde{\mathbb{T}}$ of \mathbb{T} can be formed as follows: For each object $V \in \mathcal{V}$, $\iota_V : \widetilde{TV} \hookrightarrow TV$ is the \mathbb{T} -closure of the image of $\eta_V : V \rightarrow TV$. But by 7.2, this is simply the usual closure of the image of the canonical map $V \rightarrow V^{**}$; i.e., \widetilde{TV} is the usual completion of V .

8. Example: Sheafification, closure, and density

Let j be a Lawvere-Tierney topology on an (elementary) topos \mathcal{X} , let Ω_j be the associated retract of the subobject classifier Ω , let \mathbb{T} be the double-dualization \mathcal{X} -monad on \mathcal{X} determined by Ω_j (4.18), and let $(\mathcal{E}, \mathcal{M}) = (\text{Epi } \mathcal{X}, \text{Mono } \mathcal{X})$. We now show by means of 6.25 that the idempotent \mathcal{X} -core of \mathbb{T} exists and is given by j -sheafification, and that moreover, the notions of \mathbb{T} -density, \mathbb{T} -closure, \mathbb{T} -separated object, and \mathbb{T} -complete object coincide with the familiar notions of j -density, j -closure, j -separated object, and j -sheaf.

8.1. The *universal closure operator* $\overline{(-)}$ determined by j is, in particular, a weakly hereditary idempotent closure operator on \mathcal{M} in \mathcal{X} ; indeed, see [16, A4.3.2, A4.3.3(ii)]. This closure operator determines an associated factorization system $(j\text{-Dense}, j\text{-CIEmb})$ (2.4.2). The elements of $j\text{-Dense}$ are called *j -dense morphisms* and those of $j\text{-CIEmb}$ *j -closed monomorphisms*. An object $Y \in \mathcal{X}$ is said to be *j -separated* (resp., a *j -sheaf*) if for every j -dense monomorphism $d : D \hookrightarrow X$, the mapping $\mathcal{X}(d, Y) : \mathcal{X}(X, Y) \rightarrow \mathcal{X}(D, Y)$ is a injective (resp. bijective). Hence, letting $j\text{-DenseEmb}$ be the class of all j -dense monomorphisms, the class of all j -sheaves is $\text{Shv}(\mathcal{X}, j) = (j\text{-DenseEmb})^\perp$. Also, by 2.4.4, $j\text{-CIEmb} = (j\text{-DenseEmb})^\perp \cap \mathcal{M}$.

8.2. LEMMA. *An object $X \in \mathcal{X}$ is j -separated (resp. a j -sheaf) if and only if there exists a monomorphism (resp. a j -closed mono) $X \rightarrow \underline{\mathcal{X}}(Y, \Omega_j)$ for some $Y \in \mathcal{X}$.*

PROOF. If an object $X \in \mathcal{X}$ is j -separated, then by [22, V.3.4] there is a monomorphism $X \rightarrow \underline{\mathcal{X}}(X, \Omega_j)$. If X is moreover a j -sheaf, then since $\underline{\mathcal{X}}(X, \Omega_j)$ is a j -sheaf by [15, 3.27, 3.24] we deduce by [15, 3.26] that the above monomorphism $X \rightarrow \underline{\mathcal{X}}(X, \Omega_j)$ is j -closed. Conversely, if there is a monomorphism $m : X \rightarrow \underline{\mathcal{X}}(Y, \Omega_j)$ for some $Y \in \mathcal{X}$, then by [15, 3.27, 3.24, 3.26], X is j -separated. If the given monomorphism m is j -closed, then we deduce by [15, 3.27, 3.24, 3.26] that X is a j -sheaf. ■

8.3. LEMMA. *The class j -DenseEmb of all j -dense monomorphisms is closed under tensors in $\underline{\mathcal{X}}$.*

PROOF. Let $d : D \rightarrow X$ be a j -dense monomorphism, and let $Y \in \mathcal{X}$. Then $Y \times d : Y \times D \rightarrow Y \times X$ is a pullback of d (along the projection $Y \times X \rightarrow X$), and $(-)$ commutes (up to isomorphism) with pullback (e.g., by [16, 4.3.2]), so the object $\overline{Y \times d} : \overline{Y \times D} \rightarrow Y \times X$ of the category $\mathbf{Sub}_{\mathcal{M}}(Y \times X)$ is isomorphic to the pullback $Y \times \overline{d} : Y \times \overline{D} \rightarrow Y \times X$ of $\overline{d} : \overline{D} \rightarrow X$. But $\overline{d} \cong 1_X$ in $\mathbf{Sub}_{\mathcal{M}}(X)$ and hence $\overline{Y \times d} \cong Y \times \overline{d} \cong Y \times 1_X = 1_{Y \times X}$ in $\mathbf{Sub}_{\mathcal{M}}(Y \times X)$, so $Y \times d$ is a j -dense monomorphism. ■

8.4. COROLLARY. $\mathbf{Shv}(\mathcal{X}, j) = (j\text{-DenseEmb})^{\perp x}$, and $j\text{-CIEmb} = (j\text{-DenseEmb})^{\downarrow x} \cap \mathcal{M}$.

PROOF. It was remarked in 8.1 that

$$\mathbf{Shv}(\mathcal{X}, j) = (j\text{-DenseEmb})^{\perp}, \quad j\text{-CIEmb} = (j\text{-DenseEmb})^{\downarrow} \cap \mathcal{M},$$

so in view of 8.3 the needed equations follow from 3.6 and [21, 5.4]. ■

8.5. LEMMA. *$(j\text{-Dense}, j\text{-CIEmb})$ is an \mathcal{X} -factorization-system on $\underline{\mathcal{X}}$.*

PROOF. By [21, 2.4], monomorphisms in \mathcal{X} are the same as \mathcal{X} -enriched monomorphisms in $\underline{\mathcal{X}}$, so $\mathcal{M} = \mathbf{Mono}_{\mathcal{X}} \underline{\mathcal{X}}$ and hence by [21, 2.11], \mathcal{M} is closed under cotensors in $\underline{\mathcal{X}}$. Since $(j\text{-DenseEmb}^{\downarrow x \uparrow x}, j\text{-DenseEmb}^{\downarrow x})$ is an \mathcal{X} -prefactorization-system on $\underline{\mathcal{X}}$, $j\text{-DenseEmb}^{\downarrow x}$ is also closed under cotensors in $\underline{\mathcal{X}}$ by 2.3.1. Hence, since $j\text{-CIEmb}$ is the intersection of the classes $(j\text{-DenseEmb})^{\downarrow x}$ and \mathcal{M} by 8.4, $j\text{-CIEmb}$ is closed under cotensors in $\underline{\mathcal{X}}$, so by [21, 5.7], the factorization system $(j\text{-Dense}, j\text{-CIEmb})$ is an \mathcal{X} -factorization-system on $\underline{\mathcal{X}}$. ■

8.6. LEMMA. $j\text{-DenseEmb} \subseteq \Sigma_T$, where Σ_T is the class of all morphisms inverted by the endofunctor T .

PROOF. Ω_j is a j -sheaf (e.g., by [15, 3.27]), so by 8.4, $\Omega_j \in (j\text{-DenseEmb})^{\perp x}$. Hence each morphism $d \in j\text{-DenseEmb}$ is \mathcal{X} -orthogonal to Ω_j and therefore is inverted by the dualization functor $\underline{\mathcal{X}}(-, \Omega_j) : \mathcal{X} \rightarrow \mathcal{X}^{\text{op}}$ and hence by the double-dualization functor T for Ω_j . ■

8.7. LEMMA. $\Sigma_T \subseteq j\text{-Dense}$.

PROOF. Suppose $h : X_1 \rightarrow X_2$ lies in Σ_T . Since the monad \mathbb{T} is induced by the dualization adjunction $\underline{\mathcal{X}}(-, \Omega_j) \dashv \underline{\mathcal{X}}(-, \Omega_j) : \mathcal{X}^{\text{op}} \rightarrow \mathcal{X}$ (4.18), we know by 4.13 that $\Sigma_T = \Sigma_{\underline{\mathcal{X}}(-, \Omega_j)}$, so $\underline{\mathcal{X}}(h, \Omega_j) : \underline{\mathcal{X}}(X_2, \Omega_j) \rightarrow \underline{\mathcal{X}}(X_1, \Omega_j)$ is an isomorphism. But Ω_j classifies j -closed subobjects ([22, V.2.2]), so $h^{-1} : \text{Sub}_{\mathcal{M}}(X_2) \rightarrow \text{Sub}_{\mathcal{M}}(X_1)$ induces a bijection between the j -closed subobjects of X_2 and X_1 , respectively. Letting $h(X_1)$ denote the image of h , considered as a subobject of X_2 , we observe that its j -closure $\overline{h(X_1)}$ is a j -closed subobject of X_2 and has inverse image $h^{-1}(\overline{h(X_1)}) = X_1$. But X_2 itself is also a j -closed subobject of X_2 with the same inverse image, $h^{-1}(X_2) = X_1$, so $\overline{h(X_1)} = X_2$ (as subobjects of X_2), showing that h is j -dense. ■

8.8. THEOREM. *Given a Lawvere-Tierney topology j on a topos \mathcal{X} , if we take \mathbb{T} to be the double-dualization \mathcal{X} -monad for Ω_j and let $(\mathcal{E}, \mathcal{M})$ be the epi-mono factorization system on \mathcal{X} , then the following hold:*

1. *A morphism is \mathbb{T} -dense if and only if it is j -dense.*
2. *A monomorphism is \mathbb{T} -closed if and only if it is j -closed.*
3. *Every morphism in \mathcal{X} factors as a \mathbb{T} -dense morphism followed by a \mathbb{T} -closed monomorphism.*
4. *The \mathbb{T} -closure of a subobject is the same as its j -closure.*
5. *An object of \mathcal{X} is \mathbb{T} -complete if and only if it is a j -sheaf.*
6. *An object of \mathcal{X} is \mathbb{T} -separated if and only if it is j -separated.*
7. *The idempotent \mathcal{X} -core $\widetilde{\mathbb{T}}$ of \mathbb{T} exists, and its associated \mathcal{X} -reflective-subcategory of $\underline{\mathcal{X}}$ consists of the j -sheaves.*

PROOF. To prove 2, observe first that by 8.6 and 8.4,

$$\mathbb{T}\text{-CIEmb} = \Sigma_T^{\downarrow \mathcal{X}} \cap \mathcal{M} \subseteq (j\text{-DenseEmb})^{\downarrow \mathcal{X}} \cap \mathcal{M} = j\text{-CIEmb}.$$

Also, by 8.5 and 8.7,

$$j\text{-CIEmb} = (j\text{-Dense})^{\downarrow \mathcal{X}} \subseteq \Sigma_T^{\downarrow \mathcal{X}},$$

so $\mathbb{T}\text{-CIEmb} = j\text{-CIEmb}$ and 2 is proved. Now 1, 3, and 4 follow, since by 8.5 and 6.3 we find that

$$j\text{-Dense} = (j\text{-CIEmb})^{\uparrow \mathcal{X}} = (\mathbb{T}\text{-CIEmb})^{\uparrow \mathcal{X}} = \mathbb{T}\text{-Dense}.$$

It now follows by 8.2 that an object $X \in \mathcal{X}$ is j -separated (resp. a j -sheaf) if and only if there is an \mathcal{M} -embedding (resp. a \mathbb{T} -closed \mathcal{M} -embedding) $X \hookrightarrow \underline{\mathcal{X}}(Y, \Omega_j)$ for some $Y \in \mathcal{X}$. Hence, since \mathbb{T} is induced by the \mathcal{X} -adjunction $F \dashv G : \mathcal{X}^{\text{op}} \rightarrow \underline{\mathcal{X}}$ with $F = G = \underline{\mathcal{X}}(-, \Omega_j)$ (4.18), we deduce 5 and 6 by 6.13, using the fact that by 6.25, every \mathbb{T} -complete object is \mathbb{T} -separated.

In view of the above, 7 now follows from 6.25. ■

8.9. Let \mathcal{Y} be a *Grothendieck quasitopos*; equivalently, let \mathcal{Y} be the category of K -separated J -sheaves on a small *bisite* (\mathcal{C}, J, K) ([16, C2.2.13]), so that \mathcal{C} is a small category and J, K are coverages on \mathcal{C} with $J \subseteq K$. Let $\mathcal{X} := [\mathcal{C}^{\text{op}}, \text{Set}]$ be the presheaf topos, and let j, k be the Lawvere-Tierney topologies associated to J, K , respectively. For each $i := j, k$, let \mathbb{T}_i be the double-dualization \mathcal{X} -monad on $\underline{\mathcal{X}}$ for Ω_i , and let $\Sigma_i := \Sigma_{T_i}$ be the class of all morphisms inverted by T_i . By 8.8, \mathcal{Y} consists of the \mathbb{T}_k -separated \mathbb{T}_j -complete (equivalently, Σ_j -complete) objects of \mathcal{X} , i.e. $\mathcal{Y} = \underline{\mathcal{X}}_{(\mathbb{T}_k, \Sigma_j)}$. By 4.14, $(\Sigma_i, \Sigma_i^{\perp \mathcal{X}})$ is an \mathcal{X} -orthogonal-pair for each $i := j, k$, and by 8.8 5,

$$\Sigma_k^{\perp \mathcal{X}} = \text{Shv}(\mathcal{C}, K) \subseteq \text{Shv}(\mathcal{C}, J) = \Sigma_j^{\perp \mathcal{X}},$$

so we deduce that $\Sigma_j \subseteq \Sigma_k$. Applying 8.8 3 to j , we find that \mathbb{T}_k and Σ_j satisfy the hypotheses of 6.20, and we deduce that $\mathcal{Y} = \underline{\mathcal{X}}_{(\mathbb{T}_k, \Sigma_j)}$ is an \mathcal{X} -enriched reflective subcategory of $\underline{\mathcal{X}}$.

References

- [1] H. Applegate and M. Tierney, *Iterated cotriples*, Lecture Notes in Math. **137** (1970), 56–99.
- [2] F. Borceux and B. Day, *Universal algebra in a closed category*, J. Pure Appl. Algebra **16** (1980), no. 2, 133–147.
- [3] C. Casacuberta and A. Frei, *Localizations as idempotent approximations to completions*, J. Pure Appl. Algebra **142** (1999), no. 1, 25–33.
- [4] C. Casacuberta, G. Peschke, and M. Pfenniger, *On orthogonal pairs in categories and localisation*, London Math. Soc. Lecture Note Ser., vol. 175, Cambridge Univ. Press, 1992, pp. 211–223.
- [5] C. Cassidy, M. Hébert, and G. M. Kelly, *Reflective subcategories, localizations and factorization systems*, J. Austral. Math. Soc. Ser. A **38** (1985), no. 3, 287–329.
- [6] B. Day, *On adjoint-functor factorisation*, Lecture Notes in Math. **420** (1974), 1–19.
- [7] D. Dikranjan and E. Giuli, *Closure operators. I*, Topology Appl. **27** (1987), no. 2, 129–143.
- [8] D. Dikranjan and W. Tholen, *Categorical structure of closure operators*, Kluwer Academic Publishers Group, Dordrecht, 1995.
- [9] E. J. Dubuc, *Kan extensions in enriched category theory*, Lecture Notes in Mathematics, Vol. 145, Springer-Verlag, 1970.
- [10] S. Eilenberg and G. M. Kelly, *Closed categories*, Proc. Conf. Categorical Algebra (La Jolla, Calif., 1965), Springer, 1966, pp. 421–562.

- [11] S. Fakir, *Monade idempotente associée à une monade*, C. R. Acad. Sci. Paris Sér. A-B **270** (1970), A99–A101.
- [12] P. J. Freyd and G. M. Kelly, *Categories of continuous functors. I*, J. Pure Appl. Algebra **2** (1972), 169–191.
- [13] J. W. Gray, *Formal category theory: adjointness for 2-categories*, Lecture Notes in Mathematics, Vol. 391, Springer-Verlag, 1974.
- [14] M. Hébert, *Weak reflections and weak factorization systems*, Appl. Categ. Structures **19** (2011), no. 1, 9–38.
- [15] P. T. Johnstone, *Topos theory*, Academic Press, London-New York, 1977.
- [16] ———, *Sketches of an elephant: a topos theory compendium. Vols. 1,2*, Oxford University Press, 2002.
- [17] G. M. Kelly, *Basic concepts of enriched category theory*, Repr. Theory Appl. Categ. (2005), no. 10, Reprint of the 1982 original [Cambridge Univ. Press].
- [18] G. M. Kelly and R. Street, *Review of the elements of 2-categories*, Lecture Notes in Math. **420** (1974), 75–103.
- [19] J. Lambek and B. A. Rattray, *Localization and sheaf reflectors*, Trans. Amer. Math. Soc. **210** (1975), 279–293.
- [20] R. B. B. Lucyshyn-Wright, *Riesz-Schwartz extensive quantities and vector-valued integration in closed categories*, Ph.D. thesis, York University, 2013, arXiv:1307.8088.
- [21] ———, *Enriched factorization systems*, Theory Appl. Categ. **29** (2014), no. 18, 475–495.
- [22] S. Mac Lane and I. Moerdijk, *Sheaves in geometry and logic*, Springer-Verlag, 1994.
- [23] D. Pumplün, *Universelle und spezielle Probleme*, Math. Ann. **198** (1972), 131–146.
- [24] H. H. Schaefer and M. P. Wolff, *Topological vector spaces*, second ed., Springer-Verlag, 1999.
- [25] W. Tholen, *Closure operators and their middle-interchange law*, Topology Appl. **158** (2011), no. 17, 2437–2441.

Department of Pure Mathematics and Mathematical Statistics
University of Cambridge
Wilberforce Road, Cambridge, UK
Email: rbb12@dpms.cam.ac.uk

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is $\text{T}_{\text{E}}\text{X}$, and $\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}2\epsilon$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

$\text{T}_{\text{E}}\text{X}$ TECHNICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT $\text{T}_{\text{E}}\text{X}$ EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Ieke Moerdijk, Radboud University Nijmegen: i.moerdijk@math.ru.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Edinburgh: Alex.Simpson@ed.ac.uk

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: rfcwalters@gmail.com

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca