

A GALOIS THEORY FOR MONOIDS

Dedicated to Manuela Sobral on the occasion of her seventieth birthday

ANDREA MONTOLI, DIANA RODELO AND TIM VAN DER LINDEN

ABSTRACT. We show that the adjunction between monoids and groups obtained via the Grothendieck group construction is admissible, relatively to surjective homomorphisms, in the sense of categorical Galois theory. The central extensions with respect to this Galois structure turn out to be the so-called *special homogeneous surjections*.

Introduction

An ACTION of a monoid B on a monoid X can be defined as a monoid homomorphism $B \rightarrow \text{End}(X)$, where $\text{End}(X)$ is the monoid of endomorphisms of X . These actions were studied in [15], where it is shown that they are equivalent to a certain class of split epimorphisms, called *Schreier split epimorphisms* in the recent paper [13]. Some properties of Schreier split epimorphisms, as well as the closely related notions of *special Schreier surjection* and *Schreier reflexive relation*, were then studied in [2] and [3], where the foundations for a cohomology theory of monoids are laid. Many typical properties of the category of groups, such as the *Split Short Five Lemma* or the fact that any internal reflexive relation is transitive, remain valid in the category of monoids when, in the spirit of relative homological algebra, those properties are restricted to Schreier split epimorphisms and Schreier reflexive relations. When an action $B \rightarrow \text{End}(X)$ factors through the group $\text{Aut}(X)$ of automorphisms of X , the corresponding split epimorphism is called *homogeneous* [2]. Some properties of homogeneous split epimorphisms and of the related notions of *special homogeneous surjection* and *homogeneous reflexive relation* were also studied in [2] and [3].

The first and the second author were supported by the Centro de Matemática da Universidade de Coimbra (CMUC), funded by the European Regional Development Fund through the program COMPETE and by the Portuguese Government through the FCT–Fundação para a Ciência e a Tecnologia under the project PEst-C/MAT/UI0324/2013 and grants number PTDC/MAT/120222/2010 and SFRH/BPD/69661/2010. The third author is a Research Associate of the Fonds de la Recherche Scientifique–FNRS. He would like to thank CMUC for its kind hospitality during his stays in Coimbra.

Received by the editors 2014-02-05 and, in revised form, 2014-04-25.

Transmitted by Walter Tholen. Published on 2014-05-01.

2010 Mathematics Subject Classification: 20M32, 20M50, 11R32, 19C09, 18F30.

Key words and phrases: categorical Galois theory; homogeneous split epimorphism; special homogeneous surjection; central extension; group completion; Grothendieck group.

© Andrea Montoli, Diana Rodelo and Tim Van der Linden, 2014. Permission to copy for private use granted.

The aim of the present paper is to approach the concept of homogeneous split epimorphism from the point of view of categorical Galois theory [6, 7]. Recall that the classical *Grothendieck group* or *group completion* construction [10, 11, 12] gives an adjunction between the categories **Mon** of monoids and **Gp** of groups, which is relevant for instance in *K*-theory, where it is used in the definition of K_0 . We prove that this adjunction is admissible in the sense of categorical Galois theory, when it is considered with respect to the class of surjective homomorphisms both in **Mon** and in **Gp**. We further show that the central extensions with respect to this adjunction are the special homogeneous surjections. This gives a positive answer to the question whether homogeneous split epimorphisms can be characterised in a way which does not refer to the underlying split epimorphism of sets.

The paper is organised as follows. In Section 1 we recall some basic notions of categorical Galois theory. In Section 2 we prove that the Grothendieck group adjunction is part of an admissible Galois structure (Theorem 2.2). In Section 3 we recall the definitions of Schreier split epimorphism and homogeneous split epimorphism, special Schreier surjection and special homogeneous surjection together with some of their properties. In Section 4 we show that the central extensions with respect to the Galois structure under consideration are exactly the special homogeneous surjections (Theorem 4.3).

1. Galois structures

We recall the definition of *Galois structure* and the concepts of *trivial*, *normal* and *central extension* arising from it, as introduced in [6, 7, 8]. For the sake of simplicity we restrict ourselves to the context of Barr-exact categories [1].

1.1. DEFINITION. A GALOIS STRUCTURE $\Gamma = (\mathcal{C}, \mathcal{X}, H, I, \eta, \epsilon, \mathcal{E}, \mathcal{F})$ consists of an adjunction

$$\mathcal{C} \begin{array}{c} \xrightarrow{I} \\ \perp \\ \xleftarrow{H} \end{array} \mathcal{X}$$

with unit $\eta: 1_{\mathcal{C}} \Rightarrow HI$ and counit $\epsilon: IH \Rightarrow 1_{\mathcal{X}}$ between Barr-exact categories \mathcal{C} and \mathcal{X} , as well as classes of morphisms \mathcal{E} in \mathcal{C} and \mathcal{F} in \mathcal{X} such that:

- (1) \mathcal{E} and \mathcal{F} contain all isomorphisms;
- (2) \mathcal{E} and \mathcal{F} are pullback-stable;
- (3) \mathcal{E} and \mathcal{F} are closed under composition;
- (4) $H(\mathcal{F}) \subseteq \mathcal{E}$;
- (5) $I(\mathcal{E}) \subseteq \mathcal{F}$.

We will follow [7] and call the morphisms in \mathcal{E} and \mathcal{F} FIBRATIONS.

1.2. DEFINITION. A TRIVIAL EXTENSION is a fibration $f: A \rightarrow B$ in \mathcal{C} such that the square

$$\begin{array}{ccc} A & \xrightarrow{\eta_A} & HI(A) \\ f \downarrow & \lrcorner & \downarrow HI(f) \\ B & \xrightarrow{\eta_B} & HI(B) \end{array}$$

is a pullback. A CENTRAL EXTENSION is a fibration f whose pullback $p^*(f)$ along *some* fibration p is a trivial extension. A NORMAL EXTENSION is a fibration such that its kernel pair projections are trivial extensions.

It is well known and easy to see that trivial extensions are always central extensions and that any normal extension is automatically central.

Given an object B in \mathcal{C} we consider the induced adjunction

$$(\mathcal{E} \downarrow B) \begin{array}{c} \xrightarrow{I^B} \\ \xleftarrow{H^B} \end{array} (\mathcal{F} \downarrow I(B)),$$

where we write $(\mathcal{E} \downarrow B)$ for the full subcategory of the slice category $(\mathcal{C} \downarrow B)$ determined by morphisms in \mathcal{E} ; similarly for $(\mathcal{F} \downarrow I(B))$. Here I^B is the restriction of I , and H^B sends a fibration $g: X \rightarrow I(B)$ to the pullback

$$\begin{array}{ccc} A & \longrightarrow & H(X) \\ H^B(g) \downarrow & \lrcorner & \downarrow H(g) \\ B & \xrightarrow{\eta_B} & HI(B) \end{array}$$

of $H(g)$ along η_B .

1.3. DEFINITION. A Galois structure $\Gamma = (\mathcal{C}, \mathcal{X}, H, I, \eta, \epsilon, \mathcal{E}, \mathcal{F})$ is said to be ADMISSIBLE when all functors H^B are full and faithful.

1.4. PROPOSITION. [9, Proposition 2.4] *If Γ is admissible, then $I: \mathcal{C} \rightarrow \mathcal{X}$ preserves pullbacks along trivial extensions. In particular, the trivial extensions are pullback-stable, so that every trivial extension is a normal extension.*

2. The Grothendieck group of a monoid

The GROTHENDIECK GROUP (or GROUP COMPLETION) of a monoid $(M, \cdot, 1)$ is given by a group $\text{Gp}(M)$ together with a monoid homomorphism $M \rightarrow \text{Gp}(M)$ which is universal with respect to monoid homomorphisms from M to groups [10, 11, 12]. More precisely, we have

$$\text{Gp}(M) = \frac{\text{GpF}(M)}{\text{N}(M)},$$

where $\text{GpF}(M)$ denotes the free group on M and $N(M)$ is the normal subgroup generated by elements of the form $[m_1][m_2][m_1 \cdot m_2]^{-1}$ (from now on, we simply write $m_1 m_2$ instead of $m_1 \cdot m_2$). This gives us an equivalence relation \equiv on $\text{GpF}(M)$ generated by $[m_1][m_2] \equiv [m_1 m_2]$ with equivalence classes $\overline{[m_1][m_2]} = \overline{[m_1 m_2]}$. Thus, an arbitrary element in $\text{Gp}(M)$ —an equivalence class of words—can be represented by a word of the form

$$[m_1][m_2]^{-1}[m_3][m_4]^{-1} \cdots [m_n]^{\iota(n)} \quad \text{or} \quad [m_1]^{-1}[m_2][m_3]^{-1}[m_4] \cdots [m_n]^{\iota(n)},$$

where $\iota(n) = \pm 1$, $n \in \mathbb{N}$, $m_1, \dots, m_n \in M$ and no further cancellation is possible.

Let Mon and Gp represent the categories of monoids and of groups, respectively. The group completion of a monoid determines an adjunction

$$\text{Mon} \begin{array}{c} \xrightarrow{\text{Gp}} \\ \perp \\ \xleftarrow{\text{Mon}} \end{array} \text{Gp}, \tag{A}$$

where Mon is the forgetful functor. To simplify notation, we write $\text{Gp}(M)$ instead of $\text{MonGp}(M)$ when referring to the monoid structure of $\text{Gp}(M)$. The counit is $\epsilon = 1_{\text{Gp}}$ and the unit is defined, for any monoid M , by

$$\eta_M: M \rightarrow \text{Gp}(M): m \mapsto \overline{[m]}.$$

2.1. REMARK. It is well known that in general η_M is neither surjective nor injective. For example:

- The additive monoid of natural numbers is such that $\eta_{\mathbb{N}}: \mathbb{N} \rightarrow \mathbb{Z}$ is an injection. In fact, η_M is injective whenever M is a monoid with cancellation.
- The monoid $M = (\{0, 1\}, \cdot, 1)$ has a trivial Grothendieck group and therefore η_M is surjective.
- The product $\mathbb{N} \times M$, for M as above, is such that $\text{Gp}(\mathbb{N} \times M) = \mathbb{Z}$ (in fact, it is not difficult to see that the group completion functor preserves products) and $\eta_{\mathbb{N} \times M}: \mathbb{N} \times M \rightarrow \mathbb{Z}$ is neither surjective nor injective.

By choosing the classes of morphisms \mathcal{E} and \mathcal{F} to be the surjections in Mon and Gp , respectively, we obtain a Galois structure

$$\Gamma_{\text{Mon}} = (\text{Mon}, \text{Gp}, \text{Mon}, \text{Gp}, \eta, \epsilon, \mathcal{E}, \mathcal{F}).$$

Since this is the only Galois structure we shall consider in detail, without further mention we take all normal, central and trivial extensions in this paper with respect to Γ_{Mon} .

2.2. THEOREM. *The Galois structure Γ_{Mon} is admissible.*

PROOF. For any monoid M , we must prove that the functor

$$\text{Mon}^M : (\mathcal{F} \downarrow \text{Gp}(M)) \rightarrow (\mathcal{E} \downarrow M)$$

is fully faithful. Given a morphism $\alpha : (A, f) \rightarrow (B, g)$ in $(\mathcal{F} \downarrow \text{Gp}(M))$, its image through Mon^M is defined by the universal property of the front pullback below:

$$\begin{array}{ccccc}
 M \times_{\text{Gp}(M)} A & \xrightarrow{\pi_A} & A & & \\
 \downarrow \text{Mon}^M(f) & \searrow \text{Mon}^M(\alpha) & \downarrow & \searrow \alpha & \\
 & M \times_{\text{Gp}(M)} B & \xrightarrow{\pi_B} & B & \\
 & \downarrow & \downarrow f & \searrow g & \\
 & & M & \xrightarrow{\eta_M} & \text{Gp}(M)
 \end{array} \tag{B}$$

First we prove that Mon^M is faithful. Consider morphisms $\alpha, \beta : (A, f) \rightarrow (B, g)$ such that $\text{Mon}^M(\alpha) = \text{Mon}^M(\beta)$. For any $a \in A$, we prove that $\alpha(a) = \beta(a)$ by induction on the length n (supposing that no cancellations are possible) of the word that represents the class $f(a)$.

If $f(a) = \overline{[m]}$, then $(m, a) \in M \times_{\text{Gp}(M)} A$ and

$$\text{Mon}^M(\alpha)(m, a) = \text{Mon}^M(\beta)(m, a)$$

implies that $\alpha(a) = \beta(a)$. If $f(a) = \overline{[m]^{-1}}$, then $f(a^{-1}) = \overline{[m]}$ and we find $\alpha(a^{-1}) = \beta(a^{-1})$ as in the previous case; hence $\alpha(a) = \beta(a)$.

Suppose that $\alpha(a') = \beta(a')$ for those $a' \in A$ which have $f(a')$ represented by a word of length $n - 1$ or smaller. Suppose that $f(a)$ is represented by a word of length n , $n \geq 2$. It can be written as the product (= concatenation) of a word of length one and a word of length $n - 1$. By the surjectivity of f , their corresponding classes can be written as $f(a_1)$ and $f(a_1^{-1}a)$, for some $a_1 \in A$. Then

$$\alpha(a) = \alpha(a_1)\alpha(a_1^{-1}a) = \beta(a_1)\beta(a_1^{-1}a) = \beta(a)$$

by the induction hypothesis.

Now we have to show that Mon^M is full. The proof goes in two steps: first a proof by induction in the case when M is a free monoid (Lemma 2.3 below), then an extension from the free case to the general case (the subsequent Lemma 2.4). ■

2.3. LEMMA. *The functor Mon^M is full for all free monoids M .*

PROOF. Let M be a free monoid. To simplify notation, we identify the classes in $\text{Gp}(M)$ with their representatives. Consider group surjections f and g as in Diagram (B) and a monoid homomorphism

$$\gamma: (M \times_{\text{Gp}(M)} A, \text{Mon}^M(f)) \rightarrow (M \times_{\text{Gp}(M)} B, \text{Mon}^M(g)).$$

We define a group homomorphism $\alpha: (A, f) \rightarrow (B, g)$ as follows. For any $a \in A$, we define $\alpha(a)$ by decomposing a into a product of elements in the image of π_A . The main difficulty lies in proving that the result is independent of the chosen decomposition.

If $f(a) = [m]$ for some $m \in M$, then $(m, a) \in M \times_{\text{Gp}(M)} A$ and we define

$$\alpha(a) := \pi_B(\gamma(m, a)).$$

If $f(a) = [m]^{-1}$, then $f(a^{-1}) = [m]$ and we define

$$\alpha(a) := \pi_B(\gamma(m, a^{-1}))^{-1}.$$

Suppose that $a = a_1 a_2^{-1} \cdots a_n^{\iota(n)}$ such that $f(a_i) = [m_i]$, with $m_i \in M$, and n is the smallest number for which such a decomposition in $\text{Gp}(M)$ exists. Then we must put

$$\alpha(a) = \pi_B(\gamma(m_1, a_1)) \pi_B(\gamma(m_2, a_2))^{-1} \cdots \pi_B(\gamma(m_n, a_n))^{\iota(n)}; \quad (\text{C})$$

the case $a = a_1^{-1} a_2 \cdots a_n^{\iota(n)}$ can be treated similarly.

To prove that α is a homomorphism, it now suffices to show that it is well defined. That is to say, if $a = x_1 x_2^{-1} \cdots x_k^{\iota(k)}$ such that $f(x_i) = [l_i]$, with $l_i \in M$, then (C) must agree with

$$\pi_B(\gamma(l_1, x_1)) \pi_B(\gamma(l_2, x_2))^{-1} \cdots \pi_B(\gamma(l_k, x_k))^{\iota(k)}.$$

Since M is free, and hence the group $\text{Gp}(M)$ is free, if the words

$$[m_1][m_2]^{-1} \cdots [m_n]^{\iota(n)} \quad \text{and} \quad [l_1][l_2]^{-1} \cdots [l_k]^{\iota(k)}$$

are both of minimal length, then $k = n$ and $l_i = m_i$. Thus we only have to prove the result for decompositions of equal length mapping down to the same word in $\text{Gp}(M)$. We do this by induction on n .

Case $n = 1$. Suppose that $a_1 = a = x_1$ and $f(a_1) = [m_1] = f(x_1)$ for some $m_1 \in M$. Then obviously $\alpha(a_1) = \alpha(x_1)$. The same happens if $a_1 = a = x_1$ and $f(a_1) = [m_1]^{-1} = f(x_1)$ for some $m_1 \in M$.

More generally, let $a, x \in A$ be such that $f(a) = [m] = f(x)$ for some $m \in M$. Then $f(x^{-1}a) = [1]$, so

$$\alpha(a) = \pi_B(\gamma(m, a)) = \pi_B(\gamma(m, x)) \pi_B(\gamma(1, x^{-1}a)) = \alpha(x) \alpha(x^{-1}a)$$

which implies that

$$\alpha(x^{-1}a) = \alpha(x)^{-1} \alpha(a).$$

This formula will be useful in the sequel of the proof.

Case $n = 2$. Now consider $a \in A$ such that $a_1 a_2^{-1} = a = x_1 x_2^{-1}$ and $f(a_i) = [m_i] = f(x_i)$ with $m_i \in M$. Then $\alpha(x_i^{-1} a_i) = \alpha(x_i)^{-1} \alpha(a_i)$ by the formula above. Hence $x_1^{-1} a_1 = x_2^{-1} a_2$ implies $\alpha(x_1)^{-1} \alpha(a_1) = \alpha(x_2)^{-1} \alpha(a_2)$, so that

$$\alpha(a_1) \alpha(a_2)^{-1} = \alpha(x_1) \alpha(x_2)^{-1}.$$

The case in which $a_1^{-1} a_2 = a = x_1^{-1} x_2$ and $f(a_i) = [m_i] = f(x_i)$ is similar.

Case $n = 3$. Suppose $a_1 a_2^{-1} a_3 = a = x_1 x_2^{-1} x_3$ such that $f(a_i) = [m_i] = f(x_i)$, with $m_i \in M$. Then

$$x_1^{-1} a_1 a_2^{-1} a_3 = x_2^{-1} x_3$$

gives

$$\alpha(a_2 a_1^{-1} x_1)^{-1} \alpha(a_3) = \alpha(x_2)^{-1} \alpha(x_3)$$

because they both map to the same word $[m_2]^{-1} [m_3]$. Similarly,

$$a_1 a_2^{-1} = x_1 x_2^{-1} x_3 a_3^{-1}$$

gives

$$\alpha(a_1) \alpha(a_2)^{-1} = \alpha(x_1) \alpha(a_3 x_3^{-1} x_2)^{-1}$$

because they both map to $[m_1] [m_2]^{-1}$. As a consequence, the equality

$$a_3 x_3^{-1} x_2 = a_2 a_1^{-1} x_1$$

above the word $[m_2]$ of length one gives

$$\alpha(a_3 x_3^{-1} x_2) = \alpha(a_2 a_1^{-1} x_1)$$

so that $\alpha(x_1)^{-1} \alpha(a_1) \alpha(a_2)^{-1} = \alpha(x_2)^{-1} \alpha(x_3) \alpha(a_3)^{-1}$ and thus

$$\alpha(a_1) \alpha(a_2)^{-1} \alpha(a_3) = \alpha(x_1) \alpha(x_2)^{-1} \alpha(x_3).$$

Again, the case $a_1^{-1} a_2 a_3^{-1} = a = x_1^{-1} x_2 x_3^{-1}$ can be treated analogously.

Case $n \geq 4$. Suppose that the result holds for all decompositions which map down to words of minimal length $n - 1$ or shorter in $\text{Gp}(M)$. Suppose that $a_1 a_2^{-1} a_3 \cdots a_n^{\iota(n)} = a = x_1 x_2^{-1} x_3 \cdots x_n^{\iota(n)}$ such that $f(a_i) = [m_i] = f(x_i)$, with $m_i \in M$. Then

$$(x_1^{-1} a_1 a_2^{-1}) a_3 \cdots a_n^{\iota(n)} = x_2^{-1} x_3 \cdots x_n^{\iota(n)}$$

both map to $[m_2]^{-1} \cdots [m_n]^{\iota(n)}$, so by the induction hypothesis we find

$$\alpha(a_2 a_1^{-1} x_1)^{-1} \alpha(a_3) \cdots \alpha(a_n)^{\iota(n)} = \alpha(x_2)^{-1} \alpha(x_3) \cdots \alpha(x_n)^{\iota(n)}.$$

Furthermore, $\alpha(a_2 a_1^{-1} x_1) = \alpha(a_2) \alpha(a_1)^{-1} \alpha(x_1)$ as shown above (case $n = 3$) so that

$$\alpha(a_1) \alpha(a_2)^{-1} \alpha(a_3) \cdots \alpha(a_n)^{\iota(n)} = \alpha(x_1) \alpha(x_2)^{-1} \alpha(x_3) \cdots \alpha(x_n)^{\iota(n)}.$$

The case $a_1^{-1} a_2 a_3^{-1} \cdots a_n^{\iota(n)} = a = x_1^{-1} x_2 x_3^{-1} \cdots x_n^{\iota(n)}$ being similar, this concludes the proof. ■

2.4. LEMMA. *The functor Mon^M is full for all monoids M .*

PROOF. As in the previous lemma, we simplify notation by identifying the classes in $\text{Gp}(M)$ with their representatives.

Consider group surjections f and g as in Diagram **(B)** as well as a group homomorphism

$$\gamma: (M \times_{\text{Gp}(M)} A, \text{Mon}^M(f)) \rightarrow (M \times_{\text{Gp}(M)} B, \text{Mon}^M(g)).$$

We cover the monoid M with the free monoid $F(M)$ on M , then apply the Grothendieck group functor to obtain the following commutative diagram with exact columns:

$$\begin{array}{ccc} \text{Ker}(r_M) & \longrightarrow & \text{N}(M) \\ \downarrow & & \downarrow \\ F(M) & \xrightarrow{\eta_{F(M)}} & \text{Gp}F(M) \\ r_M \downarrow & & \downarrow q_M \\ M & \xrightarrow{\eta_M} & \text{Gp}(M). \end{array}$$

We pull back $\text{Mon}^M(f)$, $\text{Mon}^M(g)$ and the morphism γ between them along the surjection r_M . We thus obtain a diagram

$$\begin{array}{ccccc} F(M) \times_{\text{Gp}(M)} A & \xrightarrow{r_M \times_{\text{Gp}(M)} 1_A} & M \times_{\text{Gp}(M)} A & \xrightarrow{\pi_A} & A \\ \downarrow r_M^*(\text{Mon}^M(f)) & \searrow r_M^*(\gamma) & \downarrow & \searrow \gamma & \downarrow f \\ F(M) \times_{\text{Gp}(M)} B & \xrightarrow{r_M \times_{\text{Gp}(M)} 1_B} & M \times_{\text{Gp}(M)} B & \xrightarrow{\pi_B} & B \\ \downarrow r_M^*(\text{Mon}^M(f)) & \searrow r_M^*(\text{Mon}^M(g)) & \downarrow \text{Mon}^M(f) & \searrow \text{Mon}^M(g) & \downarrow f \\ F(M) & \xrightarrow{r_M} & M & \xrightarrow{\eta_M} & \text{Gp}(M). \end{array}$$

Since $\eta_M r_M = q_M \eta_{F(M)}$, the left hand side triangle of this diagram can also be obtained by taking the pullbacks of f and g along $q_M \eta_{F(M)}$:

$$\begin{array}{ccccc} F(M) \times_{\text{Gp}(M)} A & \xrightarrow{\eta_{F(M)} \times_{\text{Gp}(M)} 1_A} & \text{Gp}F(M) \times_{\text{Gp}(M)} A & \xrightarrow{p_A} & A \\ \downarrow r_M^*(\text{Mon}^M(f)) & \searrow r_M^*(\gamma) & \downarrow & \searrow \beta & \downarrow f \\ F(M) \times_{\text{Gp}(M)} B & \xrightarrow{\eta_{F(M)} \times_{\text{Gp}(M)} 1_B} & \text{Gp}F(M) \times_{\text{Gp}(M)} B & \xrightarrow{p_B} & B \\ \downarrow r_M^*(\text{Mon}^M(f)) & \searrow r_M^*(\text{Mon}^M(g)) & \downarrow q_M^*(f) & \searrow q_M^*(g) & \downarrow f \\ F(M) & \xrightarrow{\eta_{F(M)}} & \text{Gp}F(M) & \xrightarrow{q_M} & \text{Gp}(M). \end{array}$$

Since the functor $\text{Mon}^{F(M)}$ is full by Lemma 2.4, we find a morphism β given by the dotted arrow above. It suffices to show that β keeps the elements of the kernel $\text{N}(M)$ (of q_M ,

thus also) of p_A and p_B fixed, because then it induces the needed $\alpha: (A, f) \rightarrow (B, g)$ by the universal property of p_A as a cokernel of its kernel.

The group $N(M)$ is generated by words $[m_1][m_2][m_1m_2]^{-1}$ as a normal subgroup of $\text{GpF}(M)$. Hence it suffices to prove for elements of the type

$$([m_1][m_2][m_1m_2]^{-1}, 1)$$

in $\text{GpF}(M) \times_{\text{Gp}(M)} A$ that

$$\beta([m_1][m_2][m_1m_2]^{-1}, 1) = ([m_1][m_2][m_1m_2]^{-1}, 1) \in \text{GpF}(M) \times_{\text{Gp}(M)} B.$$

Since f is a surjection, there exists an element $a \in A$ such that

$$(r_M \times_{\text{Gp}(M)} 1_A)([m_1][m_2], a) = (m_1m_2, a) = (r_M \times_{\text{Gp}(M)} 1_A)([m_1m_2], a).$$

For some $b \in B$ we have $\gamma(m_1m_2, a) = (m_1m_2, b)$, so using the commutativity of the second diagram, we see that

$$r_M^*(\gamma)([m_1][m_2], a) = ([m_1][m_2], b)$$

and

$$r_M^*(\gamma)([m_1m_2], a) = ([m_1m_2], b).$$

On the other hand, using the commutativity of the third diagram we find

$$\begin{aligned} \beta([m_1][m_2], a) &= \beta(\eta_{\text{F}(M)} \times_{\text{Gp}(M)} 1_A)([m_1][m_2], a) \\ &= (\eta_{\text{F}(M)} \times_{\text{Gp}(M)} 1_B)(r_M^*(\gamma)([m_1][m_2], a)) \\ &= ([m_1][m_2], b) \end{aligned}$$

and, similarly, $\beta([m_1m_2], a) = ([m_1m_2], b)$, for some $b \in B$ as above. Since β is a group homomorphism, we obtain

$$\begin{aligned} \beta([m_1][m_2][m_1m_2]^{-1}, 1) &= \beta([m_1][m_2], a)\beta([m_1m_2], a)^{-1} \\ &= ([m_1][m_2], b)([m_1m_2]^{-1}, b^{-1}) \\ &= ([m_1][m_2][m_1m_2]^{-1}, 1) \end{aligned}$$

which concludes the proof. ■

2.5. REMARK. We can restrict the group completion to commutative monoids: it is easily seen that then Γ_{Mon} restricts to an admissible Galois structure

$$\Gamma_{\text{CMon}} = (\text{CMon}, \text{Ab}, \text{CMon}, \text{Gp}|_{\text{CMon}}, \eta', \epsilon', \mathcal{E}', \mathcal{F}')$$

induced by the (co)restriction

$$\text{CMon} \begin{array}{c} \xrightarrow{\text{Gp}|_{\text{CMon}}} \\ \perp \\ \xleftarrow{\text{CMon}} \end{array} \text{Ab},$$

of the adjunction **(A)** to commutative monoids and abelian groups.

We end this section with an example showing that the adjunction **(A)** is not SEMI-LEFT-EXACT [5]: it is not admissible with respect to all morphisms, instead of just the surjections [4].

2.6. EXAMPLE. Consider $\eta_{\mathbb{N}^2}: \mathbb{N}^2 \rightarrow \mathbb{Z}^2$ with morphisms f and g as in Diagram **(B)**, where A is the subgroup of \mathbb{Z}^2 generated by $(1, -1)$, f is determined by $f(1, -1) = (1, -1)$ and

$$g: \mathbb{Z}^3 \rightarrow \mathbb{Z}^2: (k, l, m) \mapsto (k, l).$$

Then $\mathbb{N}^2 \times_{\mathbb{Z}^2} A = 0$ while $\mathbb{N}^2 \times_{\mathbb{Z}^2} \mathbb{Z}^3 = \mathbb{N}^2 \times \mathbb{Z}$, so that the functor Mon^M is not faithful: it maps, for instance, both $A \rightarrow B: (1, -1) \mapsto (1, -1, 0)$ and $(1, -1) \mapsto (1, -1, 1)$ to the zero morphism $0 \rightarrow \mathbb{N}^2 \times \mathbb{Z}$.

3. Schreier split epimorphisms and homogeneous split epimorphisms

In this section we recall some definitions and results from [2] and [3]. We work in the category Mon of monoids.

3.1. DEFINITION. Consider a split epimorphism (f, s) with its kernel:

$$N \triangleright \xrightarrow[k]{} X \begin{matrix} \xleftarrow{f} \\ \xrightarrow{s} \end{matrix} \gg Y. \tag{D}$$

It is called a SCHREIER SPLIT EPIMORPHISM when, for any $x \in X$, there exists a unique $n \in N$ such that $x = n s f(x)$.

Note that when we say “split epimorphism” we consider the chosen splitting as part of the structure; and for the sake of simplicity, we take canonical kernels—so N is a subset of X .

3.2. DEFINITION. The split epimorphism **(D)** is said to be RIGHT HOMOGENEOUS when, for every element $y \in Y$, the function $\mu_y: N \rightarrow f^{-1}(y)$ defined through multiplication on the right by $s(y)$, so $\mu_y(n) = n s(y)$, is bijective. Similarly, by duality, we can define a LEFT HOMOGENEOUS split epimorphism: now the function $N \rightarrow f^{-1}(y): n \mapsto s(y)n$ must be a bijection for all $y \in Y$. A split epimorphism is said to be HOMOGENEOUS when it is both right and left homogeneous.

3.3. PROPOSITION. [2, Propositions 2.3 and 2.4] Consider a split epimorphism (f, s) as in **(D)**. The following statements are equivalent:

- (i) (f, s) is a Schreier split epimorphism;
- (ii) there exists a unique function $q: X \rightarrow N$ such that $q(x)s f(x) = x$, for all $x \in X$;
- (iii) there exists a function $q: X \rightarrow N$ such that $q(x)s f(x) = x$ and $q(n s(y)) = n$, for all $n \in N$, $x \in X$ and $y \in Y$;
- (iv) (f, s) is right homogeneous.

3.4. DEFINITION. Given monoids Y and N , an ACTION of Y on N is a monoid homomorphism $\varphi: Y \rightarrow \text{End}(N)$, where $\text{End}(N)$ is the monoid of endomorphisms of N .

Actions correspond to Schreier split epimorphisms via a semidirect product construction:

3.5. SEMIDIRECT PRODUCTS. It is shown in [13] that any Schreier split epimorphism **(D)** corresponds to an action φ of Y on N defined by

$$\varphi(y)(n) = {}^y n = q(s(y) n)$$

for $y \in Y$ and $n \in N$. Thus (f, s) is isomorphic, as a split epimorphism, to

$$N \xrightarrow{\langle 1, 0 \rangle} N \rtimes_{\varphi} Y \xleftarrow[\langle 0, 1 \rangle]{\pi_Y} Y,$$

where $N \rtimes_{\varphi} Y$ is the semidirect product of N and Y with respect to φ : the cartesian product of sets $N \times Y$ equipped with the operation

$$(x_1, y_1) \cdot (x_2, y_2) = (x_1 \varphi_{y_1}(x_2), y_1 y_2),$$

where $\varphi_{y_1} = \varphi(y_1) \in \text{Aut}(N)$. See [13], [2] or Chapter 5 in [3] for more details.

3.6. PROPOSITION. [2, Proposition 3.8] *A Schreier split epimorphism **(D)** is homogeneous if and only if the corresponding action $\varphi: Y \rightarrow \text{End}(N)$ factors through the group $\text{Aut}(N)$ of automorphisms of N .*

3.7. LEMMA. [2, Lemma 4.1] *Consider the morphism of Schreier split epimorphisms*

$$\begin{array}{ccccc} N & \begin{array}{c} \xleftarrow{q} \\ \xrightarrow{k} \end{array} & X & \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{s} \end{array} & Y \\ \tilde{u} \downarrow & & u \downarrow & & \downarrow v \\ N' & \begin{array}{c} \xleftarrow{q'} \\ \xrightarrow{k'} \end{array} & X' & \begin{array}{c} \xleftarrow{f'} \\ \xrightarrow{s'} \end{array} & Y' \end{array}$$

and their kernels, and the restriction \tilde{u} of u to N . Then the left hand side square consisting of the functions q and q' also commutes: $q'u = \tilde{u}q$.

This lemma has the following useful consequence.

3.8. COROLLARY. *Given a morphism of Schreier split epimorphisms as in Lemma 3.7, the homomorphism \tilde{u} preserves the action of the object Y on N : for all $y \in Y$ and $n \in N$,*

$$\tilde{u}({}^y n) = {}^{v(y)} \tilde{u}(n).$$

PROOF. $\tilde{u}({}^y n) = \tilde{u}q(s(y) n) = q'u(s(y) n) = q'(us(y) u(n)) = q'(s'v(y) \tilde{u}(n)) = {}^{v(y)} \tilde{u}(n) \blacksquare$

We now extend these concepts to surjections which are not necessarily split.

3.9. DEFINITION. Given a surjective homomorphism g of monoids and its kernel pair

$$\text{Eq}(g) \begin{array}{c} \xrightarrow{\pi_1} \\ \xleftarrow{\Delta} \\ \xrightarrow{\pi_2} \end{array} X \xrightarrow{g} Y, \tag{E}$$

g is called a SPECIAL SCHREIER SURJECTION when (π_1, Δ) is a Schreier split epimorphism. It is called a SPECIAL HOMOGENEOUS SURJECTION when (π_1, Δ) is a homogeneous split epimorphism.

As a consequence of Theorem 5.5 in [2], if g is a special Schreier surjection, then its kernel is necessarily a group.

3.10. REMARK. The name *Schreier extension* was used in [16, 14] to describe a different, but closely related concept.

3.11. REMARK. A special Schreier (resp. homogeneous) surjection which is a split epimorphism is always a Schreier (resp. homogeneous) split epimorphism. Yet a Schreier (resp. homogeneous) split epimorphism is not necessarily a special Schreier (resp. homogeneous) surjection. Indeed, according to Proposition 3.1.12 in [3], a Schreier (resp. homogeneous) split epimorphism is a special Schreier (resp. homogeneous) surjection if and only if its kernel is a group. In fact, by Proposition 2.3.4 in [3], taking the kernel pair of a Schreier split epimorphism (f, s) as in (D), we do obtain a Schreier split epimorphism $(\pi_1, \langle sf, 1_X \rangle)$. Nevertheless, the split epimorphism (π_1, Δ) need not be Schreier.

As a consequence of Theorem 5.5 in [2] and of the remark above we have:

3.12. COROLLARY. *A surjective homomorphism g as in (E) is a special Schreier (resp. special homogeneous) surjection if and only if the kernel pair projection π_1 is a special Schreier (resp. special homogeneous) surjection.*

3.13. PROPOSITION. [3, Proposition 7.1.4] *Special Schreier and special homogeneous surjections are stable under products and pullbacks.*

3.14. PROPOSITION. [3, Proposition 7.1.5] *Given any pullback*

$$\begin{array}{ccc} X & \xrightarrow{g} & X' \\ f \downarrow & \lrcorner & \downarrow f' \\ Y & \xrightarrow{h} & Y' \end{array}$$

with g and h surjective homomorphisms, if f is a special Schreier (resp. special homogeneous) surjection, then so is f' .

3.15. PROPOSITION. [2, Proposition 3.4] *Any split epimorphism (D) such that Y is a group is a homogeneous split epimorphism.*

3.16. REMARK. According to the proposition above and to Remark 3.11, a split epimorphism (\mathbf{D}) such that Y is a group is a special homogeneous surjection if and only if its kernel N is a group. Furthermore, in that case also X is a group [3, Corollary 2.2.5]. Conversely, every surjective homomorphism between groups is a special homogeneous surjection.

4. Normal extensions and central extensions

In this section we characterise the trivial split extensions, the central and the normal extensions in the Galois structure Γ_{Mon} . The central extensions turn out to be precisely the special homogeneous surjections, while a split epimorphism of monoids is a trivial extension if and only if it is a special homogeneous surjection. This gives a characterisation which does not refer to the underlying split epimorphism of sets: Definition 3.2 in terms of elements, Proposition 3.3 where the splitting q is a *function* rather than a morphism of monoids.

4.1. LEMMA. *Any morphism of homogeneous split epimorphisms and their kernels*

$$\begin{array}{ccccc}
 N & \xrightarrow{k} & X & \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{s} \end{array} & Y \\
 \tilde{u} \downarrow & & u \downarrow & & \downarrow \eta_Y \\
 N' & \xrightarrow{k'} & X' & \begin{array}{c} \xleftarrow{f'} \\ \xrightarrow{s'} \end{array} & \text{Gp}(Y)
 \end{array}$$

factors into the composite

$$\begin{array}{ccccc}
 N & \xrightarrow{k} & X & \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{s} \end{array} & Y \\
 \parallel & & \downarrow \overline{\eta_Y} & & \downarrow \eta_Y \\
 N & \xrightarrow{k''} & X'' & \begin{array}{c} \xleftarrow{f''} \\ \xrightarrow{s''} \end{array} & \text{Gp}(Y) \\
 \tilde{u} \downarrow & & \downarrow \bar{u} & & \parallel \\
 N' & \xrightarrow{k'} & X' & \begin{array}{c} \xleftarrow{f'} \\ \xrightarrow{s'} \end{array} & \text{Gp}(Y)
 \end{array}$$

of morphisms of homogeneous split epimorphisms and their kernels, where φ is as in Proposition 3.6 and $X'' = N \rtimes_{\overline{\varphi}} \text{Gp}(Y)$ for $\overline{\varphi}: \text{Gp}(Y) \rightarrow \text{Aut}(N)$, the unique group homomorphism satisfying $\varphi = \overline{\varphi}\eta_Y$.

PROOF. As mentioned above, we have $X \cong N \rtimes_{\varphi} Y$ for $\varphi: Y \rightarrow \text{Aut}(N)$. By adjointness, this monoid morphism φ gives rise to a unique group homomorphism $\overline{\varphi}: \text{Gp}(Y) \rightarrow \text{Aut}(N)$ for which $\overline{\varphi}\eta_Y = \varphi$. Note that $\overline{\varphi}$ is necessarily given by

$$\overline{\varphi}(\overline{[y_1][y_2]^{-1} \cdots [y_n]^{\iota(n)}}) = \varphi_{y_1} \varphi_{y_2}^{-1} \cdots \varphi_{y_n}^{\iota(n)} \in \text{Aut}(N) \tag{F}$$

and

$$\overline{\varphi}(\overline{[y_1]^{-1}[y_2] \cdots [y_n]^{\iota(n)}}) = \varphi_{y_1}^{-1} \varphi_{y_2} \cdots \varphi_{y_n}^{\iota(n)} \in \text{Aut}(N). \tag{G}$$

Via the functoriality of the semidirect product construction this already yields the upper part of the diagram, where $\overline{\eta}_Y = 1_N \rtimes \eta_Y$. This leaves us with finding $\overline{u}: X'' \rightarrow X'$.

The needed morphism $\overline{u}: N \rtimes_{\overline{\varphi}} \text{Gp}(Y) \rightarrow N' \rtimes_{\psi} \text{Gp}(Y)$, where ψ is the action for which $X' \cong N' \rtimes_{\psi} \text{Gp}(Y)$, is induced once we prove that \tilde{u} is a morphism of $\text{Gp}(Y)$ -actions. More precisely, we have to show that

$$\tilde{u}(\overline{\varphi}_z(n)) = \psi_z(\tilde{u}(n))$$

for all $z \in \text{Gp}(Y)$ and $n \in N$. Corollary 3.8 and the fact that $\varphi = \overline{\varphi}\eta_Y$ tell us precisely that this equality holds for generators $z = \eta_Y(y)$ of $\text{Gp}(Y)$, so it suffices to check that it extends to all elements of $\text{Gp}(Y)$. This needs a straightforward verification based on **(F)** and **(G)**. ■

4.2. PROPOSITION. *Consider a split epimorphism (f, s) as in **(D)**. The following statements are equivalent:*

- (i) *f is a trivial extension;*
- (ii) *f is a special homogeneous surjection.*

PROOF. (i) \Rightarrow (ii) If f is a trivial extension, then by definition the diagram

$$\begin{array}{ccc} X & \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{s} \end{array} & Y \\ \eta_X \downarrow & & \downarrow \eta_Y \\ \text{Gp}(X) & \begin{array}{c} \xleftarrow{\text{Gp}(f)} \\ \xrightarrow{\text{Gp}(s)} \end{array} & \text{Gp}(Y) \end{array} \tag{H}$$

is a pullback. By Remark 3.16, the group homomorphism $\text{Gp}(f)$ is a special homogeneous surjection; hence so is f by Proposition 3.13.

(ii) \Rightarrow (i) Given a split epimorphism (f, s) which is a special homogeneous surjection, we have to show that the square **(H)** is a pullback. Taking kernels we obtain the morphism of special homogeneous surjections and their kernels

$$\begin{array}{ccccc} N & \xrightarrow{k} & X & \begin{array}{c} \xleftarrow{f} \\ \xrightarrow{s} \end{array} & Y \\ \tilde{\eta}_X \downarrow & & \eta_X \downarrow & & \downarrow \eta_Y \\ K & \xrightarrow{k'} & \text{Gp}(X) & \begin{array}{c} \xleftarrow{\text{Gp}(f)} \\ \xrightarrow{\text{Gp}(s)} \end{array} & \text{Gp}(Y) \end{array}$$

where, in particular, the kernel N of f is a group by Remark 3.11. By Theorem 2.3.7 in [3], the square **(H)** is a pullback precisely when $\tilde{\eta}_X$ is an isomorphism.

Lemma 4.1 gives us the diagram of solid arrows

$$\begin{array}{ccccc}
 N & \xrightarrow{k} & X & \xleftarrow[f]{s} & Y \\
 \parallel & & \downarrow \overline{\eta_Y} & & \downarrow \eta_Y \\
 N & \xrightarrow{k''} & X'' & \xleftarrow[f'']{s''} & \text{Gp}(Y) \\
 \downarrow \widetilde{\eta_X} & & \downarrow \overline{\eta_X} & \uparrow g & \parallel \\
 K & \xrightarrow{k'} & \text{Gp}(X) & \xleftarrow[\text{Gp}(s)]{\text{Gp}(f)} & \text{Gp}(Y).
 \end{array}$$

On the other hand, since X'' is a group (thanks to Remark 3.16), the universal property of $\text{Gp}(X)$ makes $\overline{\eta_Y}$ induce a unique group homomorphism $g: \text{Gp}(X) \rightarrow X''$ such that $g\eta_X = \overline{\eta_Y}$. Note that this g is actually a morphism of split epimorphisms:

$$f''g\eta_X = f''\overline{\eta_Y} = \eta_Y f = \text{Gp}(f)\eta_X$$

so that $f''g = \text{Gp}(f)$ by the universal property of η_X , while

$$g\text{Gp}(s)\eta_Y = g\eta_X s = \overline{\eta_Y} s = s''\eta_Y$$

and thus $g\text{Gp}(s) = s''$.

Finally, we have $\overline{\eta_X}g = 1_{\text{Gp}(X)}$ since $\overline{\eta_X}g\eta_X = \overline{\eta_X}\overline{\eta_Y} = \eta_X$. On the other hand, using Lemma 2.1.6 in [3]—which says that Schreier split epimorphisms are strongly split epimorphisms, that is, the kernel and the section are jointly strongly epimorphic—from

$$g\overline{\eta_X}k'' = g\overline{\eta_X}\overline{\eta_Y}k = g\eta_X k = \overline{\eta_Y}k = k'' \quad \text{and} \quad g\overline{\eta_X}s'' = g\text{Gp}(s) = s''$$

we conclude that $g\overline{\eta_X} = 1_{X''}$. In particular, the arrow $\widetilde{\eta_X}$ is an isomorphism, hence the square **(H)** is a pullback. \blacksquare

4.3. THEOREM. *For a surjective homomorphism of monoids g , the following statements are equivalent:*

- (i) g is a central extension;
- (ii) g is a normal extension;
- (iii) g is a special homogeneous surjection.

PROOF. Consider a surjective homomorphism and its kernel pair **(E)**. Then g is a normal extension

$$\begin{aligned}
 & \stackrel{(1.2)}{\Leftrightarrow} \pi_1 \text{ is a trivial extension} \\
 & \stackrel{(4.2)}{\Leftrightarrow} \pi_1 \text{ is a special homogeneous surjection} \\
 & \stackrel{(3.12)}{\Leftrightarrow} g \text{ is a special homogeneous surjection.}
 \end{aligned}$$

A normal extension is always a central extension by definition. To prove that (i) implies (iii), let us suppose that g is a central extension. Then there exists a fibration p such that $p^*(g)$ is a trivial extension, which makes it a normal extension by Proposition 1.4, hence a special homogeneous surjection by (ii) \Rightarrow (iii). Since p is a surjective homomorphism, we can apply Proposition 3.14 to conclude that g is a special homogeneous surjection, too. ■

4.4. WHAT ABOUT SPECIAL SCHREIER SURJECTIONS? A natural question that arises is, whether the special Schreier surjections admit a similar characterisation. More precisely, does the reflection **(A)** factorise in such a way that the special Schreier surjections become the central extensions with respect to this new adjunction? As explained in the proof of Proposition 4.2, any split epimorphism of groups is necessarily special homogeneous, which implies that so are the central extensions. Hence we would need a reflective subcategory \mathcal{X} of **Mon** in which all split epimorphisms are Schreier split epimorphisms. By Corollary 3.1.7 in [3] though, this would mean that \mathcal{X} is contained in the category of groups, which defeats the purpose.

References

- [1] M. Barr, *Exact categories*, Exact categories and categories of sheaves, Lecture Notes in Math., vol. 236, Springer, 1971, pp. 1–120.
- [2] D. Bourn, N. Martins-Ferreira, A. Montoli, and M. Sobral, *Schreier split epimorphisms between monoids*, Semigroup Forum, published online, DOI 10.1007/s00233-014-9571-6, 2014.
- [3] D. Bourn, N. Martins-Ferreira, A. Montoli, and M. Sobral, *Schreier split epimorphisms in monoids and in semirings*, Textos de Matemática (Série B), vol. 45, Departamento de Matemática da Universidade de Coimbra, 2014.
- [4] A. Carboni, G. Janelidze, G. M. Kelly, and R. Paré, *On localization and stabilization for factorization systems*, Appl. Categ. Struct. **5** (1997), 1–58.
- [5] C. Cassidy, M. Hébert, and G. M. Kelly, *Reflective subcategories, localizations and factorizations systems*, J. Austral. Math. Soc. **38** (1985), 287–329.
- [6] G. Janelidze, *Pure Galois theory in categories*, J. Algebra **132** (1990), no. 2, 270–286.
- [7] G. Janelidze, *Categorical Galois theory: revision and some recent developments*, Galois connections and applications, Math. Appl., vol. 565, Kluwer Acad. Publ., 2004, pp. 139–171.
- [8] G. Janelidze and G. M. Kelly, *Galois theory and a general notion of central extension*, J. Pure Appl. Algebra **97** (1994), no. 2, 135–161.
- [9] G. Janelidze and G. M. Kelly, *The reflectiveness of covering morphisms in algebra and geometry*, Theory Appl. Categ. **3** (1997), no. 6, 132–159.

- [10] A. I. Mal'cev, *On the immersion of an algebraic ring into a field*, Math. Ann. **113** (1937), 686–691.
- [11] A. I. Mal'cev, *On the immersion of associative systems into groups, I*, Mat. Sbornik N. S. **6** (1939), 331–336.
- [12] A. I. Mal'cev, *On the immersion of associative systems into groups, II*, Mat. Sbornik N. S. **8** (1940), 241–264.
- [13] N. Martins-Ferreira, A. Montoli, and M. Sobral, *Semidirect products and crossed modules in monoids with operations*, J. Pure Appl. Algebra **217** (2013), 334–347.
- [14] A. Patchkoria, *On extensions of semimodules*, Bull. Acad. Sci. Georgian SSR **84** (1976), no. 3, 545–548.
- [15] A. Patchkoria, *Crossed semimodules and Schreier internal categories in the category of monoids*, Georgian Math. J. **5** (1998), no. 6, 575–581.
- [16] L. Rédei, *Die Verallgemeinerung der Schreierschen Erweiterungstheorie*, Acta Sci. Math. (Szeged) **14** (1952), 252–273.

CMUC, Universidade de Coimbra, 3001–501 Coimbra, Portugal

Departamento de Matemática, Faculdade de Ciências e Tecnologia, Universidade do Algarve, Campus de Gambelas, 8005–139 Faro, Portugal

Institut de Recherche en Mathématique et Physique, Université catholique de Louvain, chemin du cyclotron 2 bte L7.01.02, B–1348 Louvain-la-Neuve, Belgium

Email: `montoli@mat.uc.pt`

`drodelo@ualg.pt`

`tim.vanderlinden@uclouvain.be`

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES (ISSN 1201-561X) will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods.

Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

Full text of the journal is freely available in .dvi, Postscript and PDF from the journal's server at <http://www.tac.mta.ca/tac/> and by ftp. It is archived electronically and in printed paper format.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. For institutional subscription, send enquiries to the Managing Editor, Robert Rosebrugh, rrosebrugh@mta.ca.

INFORMATION FOR AUTHORS The typesetting language of the journal is \TeX , and $\text{\LaTeX}2\epsilon$ strongly encouraged. Articles should be submitted by e-mail directly to a Transmitting Editor. Please obtain detailed information on submission format and style files at <http://www.tac.mta.ca/tac/>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

\TeX NICAL EDITOR. Michael Barr, McGill University: barr@math.mcgill.ca

ASSISTANT \TeX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Lawrence Breen, Université de Paris 13: breen@math.univ-paris13.fr

Ronald Brown, University of North Wales: [ronnie.profbrown\(at\)btinternet.com](mailto:ronnie.profbrown(at)btinternet.com)

Valeria de Paiva: valeria.depaiiva@gmail.com

Ezra Getzler, Northwestern University: [getzler\(at\)northwestern\(dot\)edu](mailto:getzler(at)northwestern(dot)edu)

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Martin Hyland, University of Cambridge: M.Hyland@dpmms.cam.ac.uk

Anders Kock, University of Aarhus: kock@imf.au.dk

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Ieke Moerdijk, Radboud University Nijmegen: i.moerdijk@math.ru.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Edinburgh: Alex.Simpson@ed.ac.uk

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: street@math.mq.edu.au

Walter Tholen, York University: tholen@mathstat.yorku.ca

Myles Tierney, Rutgers University: tierney@math.rutgers.edu

Robert F. C. Walters, University of Insubria: robert.walters@uninsubria.it

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca