

HOMOTOPY THEORY WITH *-CATEGORIES

ULRICH BUNKE

ABSTRACT. We construct model category structures on various types of (marked) *-categories. These structures are used to present the infinity categories of (marked) *-categories obtained by inverting (marked) unitary equivalences. We use this presentation to explicitly calculate the ∞ -categorical G -fixed points and G -orbits for G -equivariant (marked) *-categories.

Contents

1	Introduction	781
2	(Marked) *-categories and linear versions	787
3	Adjunctions	791
4	Classifier categories	795
5	Unitary equivalences and ∞ -categories of *-categories	798
6	The tensor and power structure over groupoids	802
7	The resolution	810
8	Completeness, cocompleteness, and local presentability	812
9	Verification of the model category axioms	822
10	The simplicial axioms	829
11	Cofibrant generation	832
12	The construction $\mathbf{A} \mapsto \hat{\mathbf{A}}^G$	837
13	Infinity-categorical G -fixed points	839
14	Infinity-categorical G -orbits	845

1. Introduction

1.1. MODEL CATEGORIES. If \mathcal{C} is a category and W is a set of morphisms in \mathcal{C} , then one can consider the ∞ -category $\mathcal{C}[W^{-1}]$. If the relative category (\mathcal{C}, W) extends to a

Acknowledgement: The author was supported by the SFB 1085 *Higher Invariants* funded by the Deutsche Forschungsgemeinschaft DFG. He thanks Markus Land for valuable discussions and pointing out the reference [Del10]. He profited from discussions with Denis-Charles Cisinski about injective model category structures. He furthermore thanks Dmitri Pavlov for convincing him that Proposition 8.7 should be true, and Daniel Kasprowski and Christoph Winges for discussing applications of the results of the present paper to coarse homology theories.

Received by the editors 2017-12-19 and, in final form, 2019-09-13.

Transmitted by Ezra Getzler. Published on 2019-09-16.

2010 Mathematics Subject Classification: 18C35, 55U35.

Key words and phrases: *-categories, model categories, ∞ -categories, limits and colimits.

© Ulrich Bunke, 2019. Permission to copy for private use granted.

simplicial model category in which all objects are cofibrant, then we have an equivalence between $\mathcal{C}[W^{-1}]$ and the nerve $N(\mathcal{C}^{\text{cf}})$ of the simplicial category of cofibrant/fibrant objects of \mathcal{C} . This explicit description of $\mathcal{C}[W^{-1}]$ is sometimes very helpful in order to calculate mapping spaces in $\mathcal{C}[W^{-1}]$, or in order to identify limits or colimits of diagrams in $\mathcal{C}[W^{-1}]$.

In this note \mathcal{C} is a category of $*$ -categories¹. A $*$ -category is a category with an involution $*$ fixing the objects. In such a category one can talk about unitary morphisms. Furthermore, one can talk about unitary transformations between functors between $*$ -categories and therefore about unitary equivalences between $*$ -categories. One natural choice for W is the set the unitary equivalences.

There are cases where one is interested in $*$ -categories with a distinguished subset of the unitary morphisms called marked morphisms. We call such a $*$ -category a marked $*$ -category. We can then consider the category \mathcal{C}^+ of such marked $*$ -categories with functors preserving the marked morphisms. Moreover we can talk about marked isomorphisms between functors between marked $*$ -categories. In this case we let W be the subset of morphisms which are invertible up to marked isomorphism.

In the present paper we consider the following categories \mathcal{C} of $*$ -categories and their marked versions \mathcal{C}^+ .

1. $*$ -categories ${}^*\mathbf{Cat}_1$: categories \mathbf{A} with an involution $*$: $\mathbf{A} \rightarrow \mathbf{A}^{\text{op}}$.
2. \mathbb{C} -linear $*$ -categories ${}^*\mathbb{C}\mathbf{Cat}_1$: $*$ -categories enriched over \mathbb{C} -vector spaces with an anti-linear involution.
3. pre- C^* -categories $C_{\text{pre}}^*\mathbf{Cat}_1$: \mathbb{C} -linear $*$ -categories which admit a maximal C^* -completion.
4. C^* -categories $C^*\mathbf{Cat}_1$: pre- C^* -categories whose Hom-vector spaces are complete in the maximal norm.

If \mathbf{A} belongs to one of these examples, then a unitary morphism in \mathbf{A} is a morphism u whose inverse is given by u^* . A marking on \mathbf{A} is a choice of a subset of unitary morphisms containing all identities which is closed under composition and the $*$ -operation. A morphism between marked categories must send marked morphisms to marked morphisms. We write ${}^*\mathbf{Cat}_1^+$, ${}^*\mathbb{C}\mathbf{Cat}_1^+$, $C_{\text{pre}}^*\mathbf{Cat}_1^+$ and $C^*\mathbf{Cat}_1^+$ for the categories of marked objects in these examples. The subscript 1 indicates that we consider them as 1-categories.

The case of C^* -categories has been considered previously in [Del10]. Many arguments in the present paper are modifications of the arguments given in [Del10] in order to be applicable in the other cases.

¹In order to fix size issues we use three Grothendieck universes $\mathcal{U} \subseteq \mathcal{V} \subseteq \mathcal{W}$. The objects of \mathcal{C} will be categories in \mathcal{V} which are locally \mathcal{U} -small. The category \mathcal{C} itself belongs to \mathcal{W} and is locally \mathcal{V} -small.

1.2. **REMARK.** We consider \mathbb{C} -linear $*$ -categories since this case fits with the C^* -examples. The assertions about the model category on ${}^*\mathbb{C}\mathbf{Cat}_1$ and the version of Theorem 13.7 extends to the case where \mathbb{C} is replaced by an arbitrary ring with involution.

An analogous theory for marked preadditive and additive categories appears in [BEKW].

We now state the main result in detail. Let \mathcal{C} belong to the list of categories

$$\{ {}^*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \} \quad (1)$$

1.3. **DEFINITION.**

1. A weak equivalence in \mathcal{C} is a (marked) unitary equivalence.
2. A cofibration is a morphism in \mathcal{C} which is injective on objects.
3. A fibration is a morphism in \mathcal{C} which has the right-lifting property with respect to trivial cofibrations.

In condition 1 the word *marked* only applies to the four marked versions. For the simplicial structure we refer to Definition 6.19 below since its introduction needs more notation. For the definition of the notion of a cofibrantly generated model category we refer to [Hov99, Def. 2.1.17].

1.4. **THEOREM.** *The structures described in Definition 1.3 and Definition 6.19 equip \mathcal{C} with a simplicial model category structure.*

If \mathcal{C} belongs to the list

$$\{ {}^*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C^* \mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \},$$

then the model category structure is cofibrantly generated and the underlying category is locally presentable.

1.5. **REMARK.** In the case of $\mathcal{C} = C^* \mathbf{Cat}_1$ a proof of this theorem (except the local presentability) was given in [Del10].

1.6. **REMARK.** A cofibrantly generated simplicial model category which is in addition locally presentable is called combinatorial [Dug01], [Lur09, A.2.6.1]. Hence ${}^*\mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1^+, C^* \mathbf{Cat}_1$ and $C^* \mathbf{Cat}_1^+$ have combinatorial simplicial model category structures.

At the moment we do not know whether $C_{\text{pre}}^* \mathbf{Cat}_1$ or $C_{\text{pre}}^* \mathbf{Cat}_1^+$ are cofibrantly generated or locally presentable.

1.7. **REMARK.** The existence and combinatoriality of this model category structure on $^*\mathbf{Cat}_1$ has been previously asserted by Joyal in [Joy10].²

All categories \mathcal{C} in the list (1) have a notion of (marked) unitary equivalences. Inverting the (marked) unitary equivalences $W_{\mathcal{C}}$ in the realm of $(\infty, 1)$ (short ∞)-categories we obtain the list

$$\{^*\mathbf{Cat}, {}^*_C\mathbf{Cat}, C_{\text{pre}}^*\mathbf{Cat}, C^*\mathbf{Cat}, {}^*\mathbf{Cat}^+, {}^*_C\mathbf{Cat}^+, C_{\text{pre}}^*\mathbf{Cat}^+, C^*\mathbf{Cat}^+\}$$

of ∞ -categories $\mathcal{C}_{\infty} := \mathcal{C}[W_{\mathcal{C}}^{-1}]$.

More precisely, we model ∞ -categories as quasi-categories. Our basic references are [Lur09] and [Cis19]. We identify categories with ∞ -categories using the nerve functor. In this case we will omit the nerve from the notation. If (\mathcal{C}, W) is a relative category, then there exists a localization functor ³

$$\ell : \mathcal{C} \rightarrow \mathcal{C}_{\infty} := \mathcal{C}[W^{-1}] , \tag{2}$$

see [Lur17, Def. 1.3.4.1], [Cis19, 7.1.2]. It is characterized essentially uniquely by the universal property that

$$\ell^* : \mathbf{Fun}(\mathcal{C}_{\infty}, \mathcal{D}) \rightarrow \mathbf{Fun}_W(\mathcal{C}, \mathcal{D})$$

is an equivalence for every ∞ -category \mathcal{D} , where \mathbf{Fun}_W denotes the full subcategory of functors sending the morphisms in W to equivalences.

1.8. **REMARK.** The model category structure on \mathcal{C} asserted in Theorem 1.4 provides a model for \mathcal{C}_{∞} .

In general, let \mathcal{C} be a simplicial model category with weak equivalences W and set $\mathcal{C}_{\infty} := \mathcal{C}[W^{-1}]$. We consider the full subcategory \mathcal{C}^{cf} of \mathcal{C} of cofibrant/fibrant objects which is enriched in Kan complexes. If either all objects of \mathcal{C} are cofibrant, or \mathcal{C} admits functorial factorizations (e.g., if \mathcal{C} is combinatorial), then by [DK80a], or [Lur17, Def. 1.3.4.15, Thm. 1.3.4.20] (and in addition [Lur17, Rem. 1.3.4.16] in the second case) we have an equivalence

$$\mathcal{C}_{\infty} \simeq \mathbf{N}(\mathcal{C}^{\text{cf}}) ,$$

where $\mathbf{N}(\mathcal{C}^{\text{cf}})$ is the nerve [Lur09, Def. 1.1.5.5] of the fibrant simplicial category \mathcal{C}^{cf} . In particular, for A, B in \mathcal{C}^{cf} we have an equivalence of spaces

$$\text{Map}_{\mathcal{C}_{\infty}}(\ell(A), \ell(B)) \simeq \ell_{\mathbf{sSet}}(\text{Map}_{\mathcal{C}}(A, B)) , \tag{3}$$

where $\text{Map}_{\mathcal{C}}(A, B)$ is the simplicial mapping set and $\ell_{\mathbf{sSet}} : \mathbf{sSet} \rightarrow \mathbf{sSet}[W^{-1}] \simeq \mathbf{Spc}$ is the usual localization of the category of simplicial sets at the weak homotopy equivalences.

²I thank Philip Hackney for pointing this out.

³In [Hin16] this localization is called the Dwyer-Kan localization, since it has been first considered by [DK80b] in the context of simplicial categories, and in order to distinguish it from the localizations considered in the book [Lur09] which are versions of Bousfield localizations.

⁴a more precise notation would be $\mathbf{N}(\mathcal{C}) \rightarrow \mathbf{N}(\mathcal{C})[W^{-1}]$

In order to see this we could use [DK80a, 1.1.(iv)] in order to relate $\ell_{\mathbf{sSet}}(\mathrm{Map}_{\mathcal{C}}(A, B))$ with $\ell_{\mathbf{sSet}}(\mathrm{Map}_{L^H(\mathcal{C}, W)}(A, B))$, where L^H denotes the hammock localization, and [Hin16, Prop. 1.2.1] in order to relate $L^H(\mathcal{C}, W)$ with the ∞ -categorical localization \mathcal{C}_∞ .

Note that for the equivalence (3) it actually suffices to assume that A is cofibrant and B is fibrant.

By [Lur09, A.3.7.6] the ∞ -category, associated as described in Remark 1.8, to a simplicial and combinatorial model category is presentable. Consequently Theorem 1.4 implies:

1.9. COROLLARY. *The ∞ -categories $^*\mathbf{Cat}$, $^*\mathbf{Cat}^+$, $^*\mathbb{C}\mathbf{Cat}$, $^*\mathbb{C}\mathbf{Cat}^+$, $C^*\mathbf{Cat}$ and $C^*\mathbf{Cat}^+$ are presentable.*

1.10. HOMOTOPY FIXED POINTS AND ORBITS. Let G be a group. The category of G -objects in a category \mathcal{C} is defined as the functor category $\mathbf{Fun}(BG, \mathcal{C})$. Here BG is the category with one object pt and $\mathrm{Hom}_{BG}(pt, pt) = G$ such that the composition is given by the multiplication in G .

We now assume that the category \mathcal{C} belongs to the list

$$\{^*\mathbf{Cat}_1, ^*\mathbb{C}\mathbf{Cat}_1, C_{\mathrm{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, ^*\mathbf{Cat}_1^+, ^*\mathbb{C}\mathbf{Cat}_1^+, C_{\mathrm{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

By $\ell : \mathcal{C} \rightarrow \mathcal{C}_\infty$ we denote the localization (2) which inverts the (marked) unitary equivalences. Furthermore, we let

$$\ell_{BG} : \mathbf{Fun}(BG, \mathcal{C}) \rightarrow \mathbf{Fun}(BG, \mathcal{C}_\infty) \tag{4}$$

denote the functor given by post-composition with ℓ . We consider an object \mathbf{A} of \mathcal{C} with an action of G , i.e, an object of $\mathbf{Fun}(BG, \mathcal{C})$. One of the purposes of the present paper is to calculate the object

$$\lim_{BG} \ell_{BG}(\mathbf{A}) .$$

Calculation of this limit amounts more precisely to provide an object \mathbf{B} of \mathcal{C} and an equivalence

$$\lim_{BG} \ell_{BG}(\mathbf{A}) \simeq \ell(\mathbf{B}) .$$

Such an object \mathbf{B} will be defined in Definition 12.1 where it is denoted by $\hat{\mathbf{A}}^G$. The construction of $\hat{\mathbf{A}}^G$ as such is not very surprising and reflects the construction of a two-categorical limit. In Theorem 13.7 we verify that it indeed represents the ∞ -categorical limit, i.e., that we have an equivalence

$$\lim_{BG} \ell_{BG}(\mathbf{A}) \simeq \ell(\hat{\mathbf{A}}^G) .$$

In order to approach the task of the calculation of the ∞ -categorical limit of the G -object $\ell_{BG}(\mathbf{A})$, using Theorem 1.4, we present the ∞ -category $\mathbf{Fun}(BG, \mathcal{C}_\infty)$ in terms of an injective model category structure on $\mathbf{Fun}(BG, \mathcal{C})$, see Remark 1.8. We then observe that

$$\lim_{BG} R(\mathbf{A}) \cong \hat{\mathbf{A}}^G ,$$

where $R : \mathbf{Fun}(BG, \mathcal{C}) \rightarrow \mathbf{Fun}(BG, \mathcal{C})$ is an explicitly given fibrant replacement functor. In model categorical language one would say that $\hat{\mathbf{A}}^G$ represents the homotopy G -invariants in \mathbf{A} . We then use general results from ∞ -category theory in order to justify that these homotopy invariants indeed represent the limit in the ∞ -categorical sense.

We now turn to G -orbits. We assume that \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

If G is a group and \mathbf{A} is an object of \mathcal{C} , then by $\underline{\mathbf{A}}$ we denote the object of $\mathbf{Fun}(BG, \mathcal{C})$ given by \mathbf{A} with the trivial action of G . We are interested in the calculation of the colimit

$$\operatorname{colim}_{BG} \ell_{BG}(\underline{\mathbf{A}})$$

in \mathcal{C}_{∞} . This again amounts to provide an object \mathbf{B} of \mathcal{C} and an equivalence

$$\operatorname{colim}_{BG} \ell_{BG}(\underline{\mathbf{A}}) \simeq \ell(\mathbf{B}) .$$

In Section 6 we construct a bifunctor

$$\mathcal{C} \times \mathbf{Grpd}_1 \rightarrow \mathcal{C} , \quad (\mathbf{A}, \mathcal{G}) \mapsto \mathbf{A}\sharp\mathcal{G} .$$

Our main result is Theorem 14.6 which asserts that

$$\operatorname{colim}_{BG} \ell_{BG}(\underline{\mathbf{A}}) \simeq \ell(\mathbf{A}\sharp BG) .$$

The main point here is again that we calculate a colimit in the infinity-categorical sense. To this end, using Theorem 1.4, we present the ∞ -category $\mathbf{Fun}(BG, \mathcal{C}_{\infty})$ in terms of a projective model category structure on $\mathbf{Fun}(BG, \mathcal{C})$. Then we show that

$$\operatorname{colim}_{BG} L(\underline{\mathbf{A}}) \cong \mathbf{A}\sharp BG ,$$

where L is an explicit cofibrant replacement functor. In model categorical language one would say that $\mathbf{A}\sharp BG$ represents the homotopy G -orbits of $\underline{\mathbf{A}}$. We then again use general results from ∞ -category theory in order to justify that these homotopy orbits represent the colimit in the ∞ -categorical sense.

The calculation of G -orbits will be applied in order to identify the values of an induction functor (Definition 14.10)

$$J^G : \mathbf{C} \xrightarrow{(-)} \mathbf{Fun}(BG, \mathcal{C}) \xrightarrow{\ell_{BG}} \mathbf{Fun}(BG, \mathcal{C}_{\infty}) \xrightarrow{\operatorname{LKan}} \mathbf{Fun}(\mathbf{Orb}(G), \mathcal{C}_{\infty}) ,$$

where LKan is the left-Kan extension functor associated to the canonical inclusion $BG \rightarrow \mathbf{Orb}(G)$. By Proposition 14.11, for a subgroup H of G , we get an equivalence

$$J^G(\mathbf{C})(H \backslash G) \simeq \ell(\mathbf{C}\sharp BH) .$$

This result will be applied in order to identify the coefficients of certain equivariant homology theories.

2. (Marked) *-categories and linear versions

In this section we introduce the notion of a *-category and various \mathbb{C} -linear versions.

2.1. DEFINITION. A *-category $(\mathbf{A}, *)$ is a small category \mathbf{A} with an involution $*$: $\mathbf{A} \rightarrow \mathbf{A}^{\text{op}}$ which fixes the objects. A morphism between *-categories is a functor between the underlying categories which preserves the involutions.

We let $*\mathbf{Cat}_1$ denote the category of *-categories and morphisms between *-categories. Usually we will just use the notation \mathbf{A} instead of $(\mathbf{A}, *)$.

If f is a morphism in a *-category, then we will write f^* for the image of f under the involution $*$.

2.2. EXAMPLE. Let G be a group. Then the category BG with morphisms $\text{Hom}_{BG}(pt, pt) = G$ can be turned into a *-category by setting

$$g^* := g^{-1} .$$

More generally, if \mathbf{G} is any groupoid, then we can consider \mathbf{G} as a *-category with the *-operation given by $g^* := g^{-1}$.

2.3. DEFINITION. A \mathbb{C} -linear *-category is a *-category \mathbf{A} which is in addition enriched over \mathbb{C} -vector spaces such that and for all objects a, a' of \mathbf{A} the map

$$* : \text{Hom}_{\mathbf{A}}(a, a') \rightarrow \text{Hom}_{\mathbf{A}}(a', a)$$

is anti-linear. A morphism between \mathbb{C} -linear *-categories is a morphism between *-categories which is also a functor between \mathbb{C} -vector space enriched categories.

We let $*_{\mathbb{C}}\mathbf{Cat}_1$ denote the category of \mathbb{C} -linear *-categories and morphisms between \mathbb{C} -linear *-categories.

2.4. REMARK. Note that in a \mathbb{C} -linear *-category \mathbf{A} for any two objects a, a' in \mathbf{A} the morphism space $\text{Hom}_{\mathbf{A}}(a, a')$ is not empty. But it may be the zero vector space. This in particular applies to the endomorphisms $\text{End}_{\mathbf{A}}(a)$. If this is the zero vector space, then we have $\text{id}_a = 0$. In this case the object a will be called a zero object. The morphism spaces from and to zero objects are zero vector spaces.

2.5. EXAMPLE. An algebra A over \mathbb{C} with an anti-linear involution can be considered as a *-category \mathbf{A} with one object pt and the \mathbb{C} -vector space of endomorphisms $\text{Hom}_{\mathbf{A}}(pt, pt) := A$. The composition is given by the algebra multiplication, and the involution on A induces the *-functor.

An important example is the underlying algebra with involution of a C^* -algebra.

We will also encounter the \mathbb{C} -linear *-category $\Delta_{*\mathbb{C}\mathbf{Cat}_1}^0$ associated to the algebra \mathbb{C} . The notation indicates that this \mathbb{C} -linear *-category is the object classifier (see Definition 4.2) in $*_{\mathbb{C}}\mathbf{Cat}_1$.

A particular example is the zero algebra 0 and the associated \mathbb{C} -linear *-category $\mathbf{0}$. The unique object in this \mathbb{C} -linear *-category is a zero object.

In the following we consider the zero algebra as a C^* -algebra. Let \mathbf{A} be a \mathbb{C} -linear $*$ -category.

2.6. DEFINITION. A representation of \mathbf{A} in a C^* -algebra B is a map $\rho : \text{Mor}(\mathbf{A}) \rightarrow B$ with the following properties:

1. For every two objects a, a' of \mathbf{A} the restriction of ρ to the subset $\text{Hom}_{\mathbf{A}}(a, a')$ of $\text{Mor}(\mathbf{A})$ is \mathbb{C} -linear.
2. $\rho(f \circ g) = \rho(f)\rho(g)$ for all pairs of composable morphisms f, g in $\text{Mor}(\mathbf{A})$
3. $\rho(f^*) = \rho(f)^*$ for all f in $\text{Mor}(\mathbf{A})$

2.7. REMARK. One could say that a representation ρ as in Definition 2.6 is a possibly non-unital morphism of \mathbb{C} -linear $*$ -categories $\mathbf{A} \rightarrow \mathbf{B}$, where \mathbf{B} is the (possibly non-unital) \mathbb{C} -linear $*$ -category associated to the C^* -algebra B as in Example 2.5. Since we do not want to talk about non-unital morphisms we avoid to use this interpretation.

2.8. EXAMPLE. We consider the commutative \mathbb{C} -algebra

$$A := \mathbb{C}[x](((x - r)^{-1})_{r \in \mathbb{R}})$$

with $*$ -operation given by $x^* = x$ as a \mathbb{C} -linear $*$ -category \mathbf{A} . Assume that $\rho : \mathbf{A} \rightarrow B$ is a representation in a non-zero C^* -algebra. Then $\rho(1)$ is a non-zero selfadjoint idempotent which commutes with $\rho(a)$ for all elements a of A . We can form the non-zero C^* -algebra $B' := \rho(1)B\rho(1)$ with unit $1' := \rho(1)$ and obtain a unital representation $\rho' : \mathbf{A} \rightarrow B'$ given by $\rho'(a) := \rho(1)\rho(a)\rho(1)$. Then $\rho'(x)$ would be a selfadjoint element in B' with empty spectrum. This is impossible. Therefore \mathbf{A} does not admit any representation in a non-zero C^* -algebra.

Let \mathbf{A} be a \mathbb{C} -linear $*$ -category and f be a morphism in \mathbf{A} .

2.9. DEFINITION. We define the maximal norm of the morphism f by

$$\|f\|_{\max} := \sup_{\rho} \|\rho(f)\|_B ,$$

where the supremum is taken over all representations of \mathbf{A} in C^* -algebras B and $\| - \|_B$ denotes the norm of B .

A priori we have $\|f\|_{\max} \in [-\infty, \infty]$. We note the following facts about the maximal norm on a \mathbb{C} -linear $*$ -category \mathbf{A} :

1. For every morphism f of \mathbf{A} we have $\|f\|_{\max} \geq 0$ since we always have the representation into the zero algebra.
2. For every two composable morphisms f and g of \mathbf{A} we have the inequality $\|g \circ f\|_{\max} \leq \|g\|_{\max} \|f\|_{\max}$.

- 3. For every morphism f of \mathbf{A} we have the C^* -equality $\|f\|_{\max}^2 = \|f^*f\|_{\max}$.
- 4. For every pair f, g of parallel morphisms in \mathbf{A} we have the C^* -inequality $\|f\|_{\max}^2 \leq \|f^* \circ f + g^* \circ g\|_{\max}$.

The last three properties hold true since they are satisfied in every representation of \mathbf{A} .

The maximal norm is preserved by unitary equivalences between \mathbb{C} -linear $*$ -categories, see Lemma 5.5 below.

2.10. DEFINITION. A *pre- C^* -category* is a \mathbb{C} -linear $*$ -category in which every morphism has a finite norm.

We let $C_{\text{pre}}^* \mathbf{Cat}_1$ denote the full subcategory of ${}^*_{\mathbb{C}} \mathbf{Cat}_1$ of pre- C^* -categories.

2.11. EXAMPLE. If \mathbf{A} comes from a C^* -algebra A with $\|-\|_A$ as in Example 2.5, then the representations of \mathbf{A} in C^* -algebras B correspond to C^* -algebra homomorphisms $A \rightarrow B$. Since morphisms between C^* -algebras are norm-bounded by 1 we have the equality

$$\|-\|_{\max} = \|-\|_A .$$

Therefore \mathbf{A} is a pre- C^* -category.

2.12. EXAMPLE. Let $A := \mathbb{C}[x]$ be the polynomial ring with the $*$ -operation given by complex conjugation. We consider A as a \mathbb{C} -linear $*$ -category \mathbf{A} . Every real number μ provides a $*$ -homomorphism $\rho_\mu : A \rightarrow \mathbb{C}$ given by evaluation of the polynomials at μ . We have $\|\rho_\mu(x)\| = |\mu|$. Hence $\|x\|_{\max} = \infty$. Consequently, the \mathbb{C} -linear $*$ -category \mathbf{A} is not a pre- C^* -category.

If K is a compact subset of \mathbb{R} , then we define for f in $\mathbb{C}[x]$

$$\|f\|_K := \sup_{k \in K} \rho_k(f) .$$

If we take the closure of $\mathbb{C}[x]$ with respect to this norm, then we get the C^* -algebra of continuous functions $C(K)$ on K . This shows that a \mathbb{C} -linear $*$ -category which is not a pre- C^* -category still may have many non-trivial C^* -closures.

2.13. EXAMPLE. We consider the \mathbb{C} -linear $*$ -category \mathbf{A} from Example 2.8. Since it has no non-trivial $*$ -representations the maximal norm on it is trivial. Consequently it is a pre- C^* -category.

2.14. DEFINITION. A *C^* -category* is a pre- C^* -category in which for every pair of objects the morphism vector space is a Banach space with respect to the norm $\|-\|_{\max}$.

We let $C^* \mathbf{Cat}_1$ denote the full subcategory of $C_{\text{pre}}^* \mathbf{Cat}_1$ of C^* -categories.

2.15. **REMARK.** According to the usual definition a C^* -category \mathbf{C} is a \mathbb{C} -linear $*$ -category in which the morphism spaces are equipped with norms such that:

1. The morphism spaces are complete.
2. For composable morphisms we have $\|f \circ g\| \leq \|g\| \|f\|$.
3. The C^* -identity $\|f^* \circ f\| = \|f\|^2$ holds true for every morphism f .
4. For every pair f, g of parallel morphisms the C^* -inequality $\|f\|^2 \leq \|f^* f + g^* g\|$ holds true.

We claim that the maximal norm on such a category considered as a \mathbb{C} -linear $*$ -category coincides with the given norm. Since every representation of C^* -categories (in the usual definition) is norm-decreasing we conclude that the maximal norm on \mathbf{C} is smaller than the given norm.

In order to show equality we form the C^* -algebra

$$A(\mathbf{C}) := \bigoplus_{c, c' \in \mathbf{C}} \text{Hom}_{\mathbf{C}}(c, c')$$

(the sum is taken in the category of Banach spaces and involves completion) with the composition given by matrix multiplication and the obvious $*$ -operation. Let $\rho : \mathbf{C} \rightarrow A(\mathbf{C})$ denote the canonical representation. Then $\|f\| = \|\rho(f)\|_{A(\mathbf{C})}$.

This implies that our definition of a C^* -category coincides with the classical one. Furthermore, being a C^* -category is just a property of a \mathbb{C} -linear $*$ -category and not an additional structure.

We have a chain of functors

$$C^* \mathbf{Cat}_1 \subseteq C^*_{\text{pre}} \mathbf{Cat}_1 \subseteq {}^*_{\mathbb{C}} \mathbf{Cat}_1 \rightarrow {}^* \mathbf{Cat}_1 \rightarrow \mathbf{Cat}_1, \tag{5}$$

where the first two are fully faithful.

2.16. **REMARK.** The categories $C^* \mathbf{Cat}_1$, ${}^*_{\mathbb{C}} \mathbf{Cat}_1$, ${}^* \mathbf{Cat}_1$ and \mathbf{Cat}_1 are closed under taking full subcategories. For $C^* \mathbf{Cat}_1$ this follows from Remark 2.15, and for the other three examples this is clear.

For $C^*_{\text{pre}} \mathbf{Cat}_1$, going to a full subcategory may increase the maximal norm since there might be functors from the subcategory to C^* -algebras which do not extend to the whole category. We can not exclude that it becomes infinite.

Let \mathcal{C} be a member of the list

$$\{ {}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, {}^*_{\mathbb{C}} \mathbf{Cat}_1, C^*_{\text{pre}} \mathbf{Cat}_1 \} \tag{6}$$

and \mathbf{A} be an object of \mathcal{C} .

Let a, a' be objects of \mathbf{A} , and u be a morphism in $\text{Hom}_{\mathbf{A}}(a, a')$.

2.17. DEFINITION. *The morphism u is called unitary if $u^*u = \text{id}_a$ and $uu^* = \text{id}_{a'}$.*

Note that the zero morphism between two zero objects is unitary.

2.18. DEFINITION. *A marking on \mathbf{A} is a choice of a subset of the unitary morphisms containing all identities which is preserved by the involution $*$ and closed under composition.*

We can talk about marked objects in \mathcal{C} . A marked object in \mathcal{C} has an underlying object in \mathcal{C} obtained by forgetting the marking.

2.19. DEFINITION. *A morphism between two marked objects in \mathcal{C} is a morphism between the underlying objects in \mathcal{C} which sends marked morphisms to marked morphisms.*

In this way we obtain categories $\mathcal{C}^*\mathbf{Cat}_1^+$, ${}^*\mathcal{C}\mathbf{Cat}_1^+$, ${}^*\mathbf{Cat}_1^+$, or \mathbf{Cat}_1^+ of marked objects and morphisms between marked objects.

2.20. REMARK. Let \mathcal{C} be in the list $\{\mathcal{C}^*\mathbf{Cat}_1^+, {}^*\mathcal{C}\mathbf{Cat}_1^+, {}^*\mathbf{Cat}_1^+, \mathbf{Cat}_1^+\}$ and \mathbf{A} be an object of \mathcal{C} . Then we can consider the subcategory \mathbf{A}^+ of \mathbf{A} with the same objects as \mathbf{A} and marked morphisms. Note that \mathbf{A}^+ is a groupoid. A morphism $f : \mathbf{A} \rightarrow \mathbf{B}$ in \mathcal{C} induces a morphism of categories $f^+ : \mathbf{A}^+ \rightarrow \mathbf{B}^+$.

2.21. EXAMPLE. Let \mathcal{C} be in the list $\{{}^*\mathbf{Cat}_1, \mathcal{C}^*\mathbf{Cat}_1, {}^*\mathcal{C}\mathbf{Cat}_1, \mathcal{C}^*\mathbf{Cat}_1\}$ and \mathbf{A} be an object of \mathcal{C} . Then we can consider the marked category $\text{mi}(\mathbf{A})$ in \mathcal{C}^+ whose marked morphisms are exactly the identities. We can also consider the marked category $\text{ma}(\mathbf{A})$ in \mathcal{C}^+ whose marked morphisms are all unitary morphisms.

3. Adjunctions

The inclusions in the chain (5) are right or left adjoints of adjunctions which we will now describe. The presence of these adjunctions turns out to be useful at various places. They will be lifted to infinity-categorical versions in Section 5.

As a convention we will denote forgetful functors by the symbol \mathcal{F} with a subscript indicating which structure or property is forgotten. Most of the functors below come in two versions, one for the unmarked, and one for the marked case. We will use the same notation for both.

Given a (marked) category \mathbf{A} we can form the free (marked) $*$ -category $\text{Free}_*(\mathbf{A})$ on \mathbf{A} . We have adjunctions

$$\text{Free}_* : \mathbf{Cat}_1 \rightleftarrows {}^*\mathbf{Cat}_1 : \mathcal{F}_* , \quad \text{Free}_* : \mathbf{Cat}_1^+ \rightleftarrows {}^*\mathbf{Cat}_1^+ : \mathcal{F}_* , \quad (7)$$

where \mathcal{F}_* denotes the functor which forgets the $*$ -operation.

3.1. REMARK. If \mathbf{A} is a category, then $\text{Free}_*(\mathbf{A})$ is obtained from \mathbf{A} by adding a morphism $f^* : a' \rightarrow a$ for every morphism $f : a \rightarrow a'$ in \mathbf{A} with the only relation that $(f \circ g)^* = g^* \circ f^*$.

A marked category (i.e., an object of \mathbf{Cat}_1^+) is a category with a distinguished set of isomorphisms. For a marked category \mathbf{A} , in the definition of $\text{Free}_*(\mathbf{A})$, we adopt the

additional relation $f^* = f^{-1}$ for all marked morphisms f of \mathbf{A} (which are isomorphisms by definition).

We furthermore have adjunctions

$$\text{Lin}_{\mathbb{C}} : {}^* \mathbf{Cat}_1 \rightleftarrows {}^*_{\mathbb{C}} \mathbf{Cat}_1 : \mathcal{F}_{\mathbb{C}} , \quad \text{Lin}_{\mathbb{C}} : {}^* \mathbf{Cat}_1^+ \rightleftarrows {}^*_{\mathbb{C}} \mathbf{Cat}_1^+ : \mathcal{F}_{\mathbb{C}} , \tag{8}$$

where $\text{Lin}_{\mathbb{C}}$ is the linearization functor, and $\mathcal{F}_{\mathbb{C}}$ denotes the functor which forgets the \mathbb{C} -linear structure.

3.2. REMARK. Here is the explicit description of $\text{Lin}_{\mathbb{C}}$. If \mathbf{A} is a $*$ -category, then $\text{Lin}_{\mathbb{C}}(\mathbf{A})$ has the same objects as \mathbf{A} , but its \mathbb{C} -vector space of morphisms is given by

$$\text{Hom}_{\text{Lin}_{\mathbb{C}}(\mathbf{A})}(a, a') := \mathbb{C}[\text{Hom}_{\mathbf{A}}(a, a')] ,$$

and the composition is defined in the canonical way.

The functor $*$: $\text{Lin}_{\mathbb{C}}(\mathbf{A}) \rightarrow \text{Lin}_{\mathbb{C}}(\mathbf{A})^{\text{op}}$ is defined as the anti-linear extension of $*$ on \mathbf{A} .

For a set X and an element x of X we consider x as an element of the complex vector space $\mathbb{C}[X]$ generated by X in the canonical way. This gives a canonical map of sets $X \rightarrow \mathbb{C}[X]$.

In the marked case the set of marked morphisms in $\text{Hom}_{\text{Lin}_{\mathbb{C}}(\mathbf{A})}(a, a')$ is defined to be the image of the set of marked morphisms in \mathbf{A} under the canonical map

$$\text{Hom}_{\mathbf{A}}(a, a') \rightarrow \mathbb{C}[\text{Hom}_{\mathbf{A}}(a, a')] = \text{Hom}_{\text{Lin}_{\mathbb{C}}(\mathbf{A})}(a, a') .$$

For a \mathbb{C} -linear $*$ -category \mathbf{B} we check the natural bijection

$$\text{Hom}_{{}^*_{\mathbb{C}} \mathbf{Cat}_1}(\text{Lin}_{\mathbb{C}}(\mathbf{A}), \mathbf{B}) \cong \text{Hom}_{{}^* \mathbf{Cat}_1}(\mathbf{A}, \mathcal{F}_{\mathbb{C}}(\mathbf{B})) . \tag{9}$$

It identifies a morphism $\Phi : \mathbf{A} \rightarrow \mathcal{F}_{\mathbb{C}}(\mathbf{B})$ with a morphism $\Psi : \text{Lin}_{\mathbb{C}}(\mathbf{A}) \rightarrow \mathbf{B}$. The functors coincide on objects. Given Φ and a morphism f in $\text{Hom}_{\text{Lin}_{\mathbb{C}}(\mathbf{A})}(a, a')$ we define

$$\Psi(f) := \sum_{\phi \in \text{Hom}_{\mathbf{A}}(a, a')} \lambda_{\phi} \Phi(\phi)$$

in $\text{Hom}_{\mathbf{B}}(\Phi(a), \Phi(a'))$, where the equality $f = \sum_{\phi \in \text{Hom}_{\mathbf{A}}(a, a')} \lambda_{\phi} \phi$ uniquely determines the collection of complex numbers $(\lambda_{\phi})_{\phi \in \text{Hom}_{\mathbf{A}}(a, a')}$. Vice versa, if Ψ is given, then for f in $\text{Hom}_{\mathbf{A}}(a, a')$ we define $\Phi(f) := \Psi(f)$ in $\text{Hom}_{\mathbf{B}}(\Psi(a), \Psi(a'))$.

In the marked case, by an inspection of these explicit formulas, one checks that the bijection (9) restricts to a bijection between the sets of morphisms between marked objects.

3.3. REMARK. If \mathbf{B} contains zero objects, then $\text{Hom}_{{}^*_{\mathbb{C}} \mathbf{Cat}_1}(\mathbf{B}, \text{Lin}_{\mathbb{C}}(\mathbf{A})) = \emptyset$ for every $*$ -category \mathbf{A} .

We have adjunctions

$$\text{Compl} : C_{\text{pre}}^* \mathbf{Cat}_1 \rightleftarrows C^* \mathbf{Cat}_1 : \mathcal{F}_- , \quad \text{Compl} : C_{\text{pre}}^* \mathbf{Cat}_1^+ \rightleftarrows C^* \mathbf{Cat}_1^+ : \mathcal{F}_- \tag{10}$$

where \mathcal{F}_- forgets the completeness condition and Compl is the completion functor.

3.4. REMARK. In the following we give an explicit description of the completion functor.

Let \mathbf{A} be a pre- C^* -category. The completion functor is the identity on objects. Furthermore, it completes the morphism spaces in the norm $\| - \|_{\max}$. Note that this completion involves factoring out the subspace of elements of zero norm. In this completion process the objects a of \mathbf{A} with $\|id_a\| = 0$ become zero objects.

In the marked case the set of marked morphisms in $\text{Hom}_{\text{Compl}(\mathbf{A})}(a, a')$ is defined to be the image of the set of marked morphisms in $\text{Hom}_{\mathbf{A}}(a, a')$ under the natural map $\text{Hom}_{\mathbf{A}}(a, a') \rightarrow \text{Hom}_{\text{Compl}(\mathbf{A})}(a, a')$. This is well-defined since this map preserves unitaries.

Let us check the bijection

$$\text{Hom}_{C^*\text{Cat}_1}(\text{Compl}(\mathbf{A}), \mathbf{B}) \cong \text{Hom}_{C^*\text{Cat}_1}(\mathbf{A}, \mathcal{F}_-(\mathbf{B})) \tag{11}$$

for a C^* -category \mathbf{B} . This bijection sends a morphism $\Phi : \mathbf{A} \rightarrow \mathcal{F}_-(\mathbf{B})$ to a morphism $\Psi : \text{Compl}(\mathbf{A}) \rightarrow \mathbf{B}$ and vice versa. These functors coincide on objects. Given Φ we can define Ψ by extension by continuity (using that the morphism spaces in \mathbf{B} are complete). Note that Φ necessarily annihilates all morphisms of zero norm and therefore factorizes over the quotients taken in the process of completion.

Vice versa, given Ψ we define Φ by restriction along the natural maps

$$\text{Hom}_{\mathbf{A}}(a, a') \rightarrow \text{Hom}_{\text{Compl}(\mathbf{A})}(a, a')$$

for all pairs of objects a, a' of \mathbf{A} .

One easily checks that these processes are inverse to each other.

In the marked case we observe by an inspection that the bijection (11) identifies morphisms between marked objects.

3.5. EXAMPLE. We continue with Example 2.2. The linearization

$$\text{Lin}_{\mathbb{C}}(BG) \cong \mathbb{C}[G]$$

of G with its usual involution is a \mathbb{C} -linear $*$ -category. It is actually a pre- C^* -category. In order to see this note that the elements of the group go to partial isometries in every representation. Furthermore,

$$\text{End}_{\text{Compl}(\text{Lin}_{\mathbb{C}}(BG))}(pt) =: C^*_{\max}(G)$$

is the maximal group- C^* -algebra.

3.6. EXAMPLE. We consider the \mathbb{C} -linear $*$ -category \mathbf{A} from Example 2.8. We get $\text{Compl}(\mathbf{A}) \cong \mathbf{0}$.

The relation between \mathbb{C} -linear $*$ -categories and pre- C^* -categories is more complicated. Let \mathbf{A} be a \mathbb{C} -linear $*$ -category. Then we can define a subcategory $\text{Bd}(\mathbf{A})$ of \mathbf{A} which has the same objects as \mathbf{A} , and whose morphisms are those morphisms of \mathbf{A} with finite maximal norm. Note that $\text{Bd}(\mathbf{A})$ is closed under composition and contains the identities of the objects since they are sent to selfadjoint projections in any representation of \mathbf{A} in a

C^* -algebra and therefore have maximal norm bounded above by 1. Hence $\text{Bd}(\mathbf{A})$ is indeed a category. It is furthermore clear that $\text{Bd}(\mathbf{A})$ is a \mathbb{C} -linear $*$ -category with enrichment and $*$ -operation induced from \mathbf{A} .

Since $\text{Bd}(\mathbf{A})$ may have representations to C^* -algebras which do not extend to \mathbf{A} we can not expect that $\text{Bd}(\mathbf{A})$ is a pre- C^* -category. But we can iterate the construction and consider

$$\text{Bd}^\infty(\mathbf{A}) := \bigcap_{n \in \mathbb{N}} \text{Bd}^n(\mathbf{A}) .$$

This is a \mathbb{C} -linear $*$ -subcategory of \mathbf{A} .

Assume now that \mathbf{A} is a marked \mathbb{C} -linear $*$ -category. Then all marked morphisms of \mathbf{A} also belong to $\text{Bd}(\mathbf{A})$ since unitary morphisms in \mathbf{A} have norm bounded by one. So $\text{Bd}(\mathbf{A})$ and hence $\text{Bd}^\infty(\mathbf{A})$ has a canonical marking consisting of all marked morphisms in \mathbf{A} .

3.7. EXAMPLE. We continue with Example 2.12. The bounded elements in $\mathbb{C}[x]$ are the constant polynomials. Hence $\text{Bd}(\mathbb{C}[x]) \cong \Delta_{\mathbb{C}\text{Cat}_1}^0$.

Let \mathbf{A} be a (marked) \mathbb{C} -linear $*$ -category.

3.8. LEMMA.

1. $\text{Bd}^\infty(\mathbf{A})$ is a (marked) pre- C^* -category.
2. Any functor $\mathbf{C} \rightarrow \mathbf{A}$ of (marked) \mathbb{C} -linear $*$ -categories where \mathbf{C} is a (marked) pre- C^* -category factorizes uniquely over $\text{Bd}^\infty(\mathbf{A})$.
3. There are adjunctions

$$\mathcal{F}_{\text{pre}} : C_{\text{pre}}^* \text{Cat}_1 \rightleftarrows {}^*_{\mathbb{C}} \text{Cat}_1 : \text{Bd}^\infty , \quad \mathcal{F}_{\text{pre}} : C_{\text{pre}}^* \text{Cat}_1^+ \rightleftarrows {}^*_{\mathbb{C}} \text{Cat}_1^+ : \text{Bd}^\infty , \quad (12)$$

where \mathcal{F}_{pre} denotes the inclusion.

PROOF. We can identify $\text{Bd}^\infty(\mathbf{A}) \cong \text{Bd}(\text{Bd}^\infty(\mathbf{A}))$. Let f be a morphism in $\text{Bd}^\infty(\mathbf{A})$. Then we can consider f as a morphism in $\text{Bd}(\text{Bd}^\infty(\mathbf{A}))$ which implies that f has a finite maximal norm. This proves 1.

Let $\phi : \mathbf{C} \rightarrow \mathbf{A}$ be a morphism of \mathbb{C} -linear $*$ -categories where \mathbf{C} is a pre- C^* -category. We show 2 by contradiction. Assume that there exists a natural number n such that $\phi(\mathbf{C}) \subseteq \text{Bd}^n(\mathbf{A})$, but $\phi(\mathbf{C}) \not\subseteq \text{Bd}^{n+1}(\mathbf{A})$. Then there exists a morphism f in \mathbf{C} and a family of representations $(\rho_k)_{k \in \mathbb{N}}$ of $\text{Bd}^n(\mathbf{A})$ in a family of C^* -algebras $(B_k)_{k \in \mathbb{N}}$ such that $\|\rho_k(\phi(f))\|_{B_k} \geq k$ for every natural number k . Since the composition $\rho_k \circ \phi$ is a representation of \mathbf{C} we see that $k \leq \|f\|_{\text{max}}$ for every natural number k . This contradicts the assumption that \mathbf{C} is a pre- C^* -category.

In the marked case we observe that ϕ sends marked morphisms in \mathbf{C} to $\text{Bd}^\infty(\mathbf{A})$ since marked morphisms are unitary and ϕ preserves unitaries.

The Assertion 3 now follows from Assertion 2. ■

Let \mathcal{C} be a member of the list

$$\{\mathbf{Cat}_1, {}^*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1\}$$

and \mathcal{C}^+ denote the corresponding marked version. We have a canonical functor $\mathcal{F}_+ : \mathcal{C}^+ \rightarrow \mathcal{C}$ which forgets the marking. This functor fits into adjunctions

$$\text{mi} : \mathcal{C} \rightleftharpoons \mathcal{C}^+ : \mathcal{F}_+ , \quad \mathcal{F}_+ : \mathcal{C}^+ \rightleftharpoons \mathcal{C} : \text{ma} \tag{13}$$

The left-adjoint mi of \mathcal{F}_+ marks the identities, and the right-adjoint ma marks all unitaries (or invertibles in the case of \mathbf{Cat}_1 , respectively).

4. Classifier categories

In this section we discuss the representability of the functors which take the sets objects, (bounded) morphisms, unitary morphisms (or marked morphisms) of a (marked) *-category in the respective cases. The role of this section is an illustration. We use the opportunity to explain the categorical meaning of the examples which will be used later.

Let \mathcal{C} be in the list

$$\{\mathbf{Cat}_1, {}^*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, \mathbf{Cat}_1^+, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+\} .$$

4.1. LEMMA. *The functor $\mathcal{C} \rightarrow \mathbf{Set}$ which sends a category in \mathcal{C} to its set of objects is representable.*

4.2. DEFINITION. *The object $\Delta_{\mathcal{C}}^0$ which represents this functor will be called the object classifier.*

PROOF. This is a case-by-case discussion. In \mathbf{Cat}_1 we can set

$$\Delta_{\mathbf{Cat}_1}^0 := pt ,$$

the category with one object pt and one morphism id_{pt} . Then in view of the adjunction (7) we have

$$\Delta_{*\mathbf{Cat}_1}^0 \cong \text{Free}_*(\Delta_{\mathbf{Cat}_1}^0) .$$

Its underlying category is again $\Delta_{\mathbf{Cat}_1}^0$.

In view of the adjunction (8) we have

$$\Delta_{*\mathbb{C}\mathbf{Cat}_1}^0 \cong \text{Lin}_{\mathbb{C}}(\Delta_{*\mathbf{Cat}_1}^0) .$$

This is the \mathbb{C} -linear *-category associated to the C^* -algebra \mathbb{C} and hence is a C^* -category. We conclude that

$$\Delta_{C^* \mathbf{Cat}_1}^0 \cong \Delta_{C_{\text{pre}}^* \mathbf{Cat}_1}^0 \cong \Delta_{*\mathbb{C}\mathbf{Cat}_1}^0$$

(as \mathbb{C} -linear *-categories). For \mathcal{C} in the list

$$\{\mathbf{Cat}_1, {}^*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1\}$$

the marked version of the object classifier is characterized by

$$\Delta_{\mathcal{C}^+}^0 \cong \text{mi}(\Delta_{\mathcal{C}}^0) ,$$

see (13) for notation. ■

Usually we will omit the subscript \mathcal{C} when the context is clear and just write Δ^0 for the object classifier. A similar conventions applies to the other classifier objects below.

4.3. LEMMA. For \mathcal{C} in $\{\mathbf{Cat}_1, {}^*\mathbf{Cat}_1, {}^*_\mathbb{C}\mathbf{Cat}_1, \mathbf{Cat}_1^+, {}^*\mathbf{Cat}_1^+, {}^*_\mathbb{C}\mathbf{Cat}_1^+\}$ the functor $\mathcal{C} \rightarrow \mathbf{Set}$ which sends a category in \mathcal{C} to its set of morphisms is representable.

For \mathcal{C} in $\{C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+\}$ this functor is not representable.

4.4. DEFINITION. The object $\Delta_{\mathcal{C}}^1$ which represents this functor will be called the morphism classifier.

PROOF. This is again a case-by-case discussion. The morphism classifiers have two objects 0 and 1 corresponding to the source and target of the morphism.

In \mathbf{Cat}_1 we let $\Delta_{\mathbf{Cat}_1}^1$ be the category with one non-trivial morphism $a : 0 \rightarrow 1$. Then in view of the adjunction (7) we have

$$\Delta_{*\mathbf{Cat}_1}^1 \cong \text{Free}_*(\Delta_{\mathbf{Cat}_1}^1) .$$

For example, a morphism $0 \rightarrow 1$ in this category is a word $aa^*aa^* \dots a^*a$. In view of the adjunction (8) we have

$$\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1 \cong \text{Lin}_{\mathbb{C}}(\Delta_{*\mathbf{Cat}_1}^1) .$$

In the marked cases, for \mathcal{C} in the list $\{\mathbf{Cat}_1, {}^*\mathbf{Cat}_1, {}^*_\mathbb{C}\mathbf{Cat}_1\}$, the morphism classifiers are characterized by

$$\Delta_{\mathcal{C}^+}^1 \cong \text{mi}_{\mathcal{C}}(\Delta_{\mathcal{C}}^1) .$$

We now come to the non-existence assertion. Assume that the pre- C^* -category \mathbf{C} represents the morphism-set functor in $C_{\text{pre}}^* \mathbf{Cat}_1$. Let $a : 0 \rightarrow 1$ be the universal morphism. As in any non-trivial pre- C^* -category there exists morphisms of arbitrary large maximal norm (just scale) we have $\|a\|_{\max} = \infty$ contradicting the assumption that \mathbf{C} is a pre- C^* -category. The same reasoning applies to $C^* \mathbf{Cat}_1$ and the marked versions. ■

The following replaces the morphisms classifier in the C^* -category cases. Let \mathcal{C} be a member of the list $\{C^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1^+\}$. The following result is [Del10, Ex. 3.8].

4.5. LEMMA. The functor $\mathcal{C} \rightarrow \mathbf{Set}$ which sends a category in \mathcal{C} to its set of morphisms with maximal norm bounded by 1 is representable.

4.6. DEFINITION. The object $\Delta_{\mathcal{C}}^{1,\text{bd}}$ which represents this functor will be called the bounded morphism classifier.

PROOF. In order to construct the bounded morphism classifier for $C^* \mathbf{Cat}_1$ we start with the \mathbb{C} -linear $*$ -category

$$\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1 \cong \text{Lin}_{\mathbb{C}}(\text{Free}_*(\Delta_{\mathbf{Cat}_1}^1)) .$$

The universal morphism a in $\Delta_{\mathbf{Cat}_1}^1$ can be considered as a morphism of $\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1$ in the natural way. We add formal inverses of $\lambda \text{id}_0 - a^*a$ in $\text{End}_{\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1}(0)$ and $\lambda \text{id}_1 - aa^*$ in $\text{End}_{\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1}(1)$ for all λ in \mathbb{C} with $|\lambda| > 1$ and obtain a new \mathbb{C} -linear $*$ -category $\widetilde{\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1}$.

Spectral theory implies that $\|a\|_{\max} = \|a^*\|_{\max} \leq 1$. Hence $\widetilde{\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1}$ is a pre- C^* -category. One easily checks that

$$\Delta_{C^*\mathbf{Cat}_1}^{1,\text{bd}} := \text{Compl}(\widetilde{\Delta_{*\mathbb{C}\mathbf{Cat}_1}^1})$$

has the required universal properties. In the marked case we have

$$\Delta_{C^*\mathbf{Cat}_1^+}^{1,\text{bd}} \cong \text{mi}(\Delta_{C^*\mathbf{Cat}_1^+}^{1,\text{bd}}).$$

■

4.7. REMARK. We do not know whether $C_{\text{pre}}^*\mathbf{Cat}_1^{(+)}$ has a bounded morphism classifier or an appropriate replacement. This is the reason that we can not show that the model category structure on $C_{\text{pre}}^*\mathbf{Cat}_1^{(+)}$ is cofibrantly generated.

We now discuss unitaries.

4.8. REMARK. If u is a unitary morphism in a \mathbb{C} -linear $*$ -category, then any representation sends u to a partial isometry. Hence $\|u\|_{\max} \leq 1$. But it may happen that $\|u\| = 0$. This is e.g. the case if u is the identity of the unique object of the pre- C^* -category considered in Example 2.8.

For a $*$ -category \mathbf{A} the counit $\mathbf{A} \rightarrow \text{Lin}_{\mathbb{C}}(\mathbf{A})$ preserves unitaries. Similarly, for a pre- C^* -category \mathbf{A} the natural morphism $\mathbf{A} \rightarrow \text{Compl}(\mathbf{A})$ preserves unitaries.

For a \mathbb{C} -linear $*$ -category \mathbf{B} the counit $\text{Bd}^\infty(\mathbf{B}) \rightarrow \mathbf{B}$ is bijective on unitaries.

Let \mathcal{C} be in the list

$$\{*\mathbf{Cat}_1, *\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *\mathbb{C}\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}.$$

4.9. LEMMA. *The functor $\mathcal{C} \rightarrow \mathbf{Set}$ which sends a category in \mathcal{C} to its set of unitary morphisms is representable.*

4.10. DEFINITION. *The object $\mathbf{1}_{\mathcal{C}}$ which represents this functor will be called the unitary morphism classifier.*

PROOF. We perform a case-by-case discussion. In $*\mathbf{Cat}_1$ we define $\mathbf{1}_{*\mathbf{Cat}_1}$ to be the category with objects 0 and 1 and non-trivial morphisms $u : 0 \rightarrow 1$ and $u^* = u^{-1} : 1 \rightarrow 0$. In view of the adjunction (8) we have

$$\mathbf{1}_{*\mathbb{C}\mathbf{Cat}_1} \cong \text{Lin}_{\mathbb{C}}(\mathbf{1}_{*\mathbf{Cat}_1}).$$

Since the generator u is sent to a unitary in any representation it is clear that $\mathbf{1}_{*\mathbb{C}\mathbf{Cat}_1}$ is a pre- C^* -category. Since $\|u\|_{\max} = 1$ it is actually a C^* -category.

Hence we have isomorphisms

$$\mathbf{1}_{C_{\text{pre}}^*\mathbf{Cat}_1} \cong \mathbf{1}_{*\mathbb{C}\mathbf{Cat}_1} \cong \mathbf{1}_{C^*\mathbf{Cat}_1}$$

(as \mathbb{C} -linear $*$ -categories). In the marked cases, for \mathcal{C} in the list $\{*\mathbf{Cat}_1, *\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1\}$ we have $\mathbf{1}_{\mathcal{C}^+} \cong \text{mi}_{\mathcal{C}}(\mathbf{1}_{\mathcal{C}})$, i.e., the universal unitary in $\mathbf{1}_{\mathcal{C}^+}$ is not marked. ■

Let \mathcal{C} be a member of the list

$$\{*\mathbf{Cat}_1, {}_{\mathbb{C}}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1\}$$

4.11. LEMMA. *The functor $\mathcal{C}^+ \rightarrow \mathbf{Set}$ which sends a category in \mathcal{C}^+ to its set of marked morphisms is representable.*

4.12. DEFINITION. *The object $\mathbf{1}_{\mathcal{C}}^+$ which represents this functor will be called the marked morphism classifier.*


PROOF. We have $\mathbf{1}_{\mathcal{C}}^+ \cong \text{ma}(\mathbf{1}_{\mathcal{C}})$, i.e., the universal unitary is now marked. ■

We now consider just categories.

4.13. LEMMA. *The functor which sends a category to its set of invertible morphisms is representable.*

4.14. DEFINITION. *We call a category \mathbb{I} which represents this functor the classifier of invertible morphisms.*

PROOF. The groupoid \mathbb{I} of the shape



has the desired properties. The morphism $0 \rightarrow 1$ is the universal invertible morphism. ■

4.15. REMARK. The groupoid \mathbb{I} is also the morphism classifier in \mathbf{Grpd}_1 .

5. Unitary equivalences and ∞ -categories of $*$ -categories

In this section we introduce the ∞ -categories of $*$ -categories, \mathbb{C} -linear $*$ -categories, pre- C^* -categories, C^* -categories and their marked versions by inverting unitary (or marked, respectively) equivalences.

Let \mathcal{C} belong to the list

$$\{*\mathbf{Cat}_1, {}_{\mathbb{C}}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, \mathbf{Cat}_1^+, *\mathbf{Cat}_1^+, {}_{\mathbb{C}}\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

Furthermore, let $f, g : \mathbf{A} \rightarrow \mathbf{B}$ be a parallel pair of morphisms in \mathcal{C} .

5.1. DEFINITION. *We say that f and g are (marked) unitary equivalent, if there exists a natural isomorphism of functors $u : f \rightarrow g$ such that $u(a)$ is a (marked) unitary morphism in $\text{Hom}_{\mathbf{B}}(f(a), g(a))$ for every object a of \mathbf{A} .*

Here the word *marked* applies in the marked cases. In these cases the word unitary can be omitted since marked morphisms are unitary by definition.

Let $f : \mathbf{A} \rightarrow \mathbf{B}$ be a morphism in \mathcal{C} .

5.2. DEFINITION. *The morphism f is a (marked) unitary equivalence if there exists a morphism $g : \mathbf{B} \rightarrow \mathbf{A}$ in \mathcal{C} such that $f \circ g$ is unitary (marked) isomorphic to $\text{id}_{\mathbf{B}}$ and $g \circ f$ is (marked) unitary isomorphic to $\text{id}_{\mathbf{A}}$.*

The following characterization of unitary or marked equivalences will be useful later. We let

$$\mathcal{F}_{\text{all}} : \mathcal{C} \rightarrow \mathbf{Cat} \tag{14}$$

be the functor which takes the underlying category (i.e., forgets all additional structures and properties). Furthermore, in the marked cases, we consider the functor

$$(-)^+ : \mathcal{C} \rightarrow \mathbf{Cat}$$

which takes the subcategory of marked morphisms, see Remark 2.20. Finally recall the functor ma from the unmarked to the marked versions which marks all unitaries, see (13).

Let $f : \mathbf{A} \rightarrow \mathbf{B}$ be a morphism in \mathcal{C} .

5.3. LEMMA.

1. *in the marked cases: The morphism f is a marked equivalence if and only if $\mathcal{F}_{\text{all}}(f)$ and f^+ are equivalences of categories.*
2. *in the unmarked cases: The morphism f is a unitary equivalence if and only if $\mathcal{F}_{\text{all}}(f)$ and $\text{ma}(f)^+$ are equivalences of categories.*

PROOF. We start with 1. If f is a marked equivalence, then by Definition 5.2 there is an inverse morphism $g : \mathbf{B} \rightarrow \mathbf{A}$ up to marked isomorphism. Then $\mathcal{F}_{\text{all}}(g)$ and g^+ are inverse equivalences of $\mathcal{F}_{\text{all}}(f)$ and f^+ , respectively.

We now assume that $\mathcal{F}_{\text{all}}(f)$ and f^+ are equivalences of categories. Then there exists a functor $g^+ : \mathbf{B}^+ \rightarrow \mathbf{A}^+$ and isomorphisms of functors

$$u : \text{id}_{\mathbf{B}^+} \rightarrow f^+ \circ g^+ , \quad v : \text{id}_{\mathbf{A}^+} \rightarrow g^+ \circ f^+ .$$

We define a morphism $g : \mathbf{B} \rightarrow \mathbf{A}$ in \mathcal{C} as follows:

1. on objects: For an object b of \mathbf{B} we define $g(b) := g^+(b)$.
2. on morphisms: For objects b, b' of B we define $g : \text{Hom}_{\mathbf{B}}(b, b') \rightarrow \text{Hom}_{\mathbf{A}}(g(b), g(b'))$ as the composition

$$\text{Hom}_{\mathbf{B}}(b, b') \xrightarrow{\cong, !} \text{Hom}_{\mathbf{B}}(f(g(b)), f(g(b'))) \xleftarrow{\cong, \mathcal{F}_{\text{all}}(f)} \text{Hom}_{\mathbf{A}}(g(b), g(b')) ,$$

where the isomorphism marked by $!$ is given by

$$\phi \mapsto u_{b'} \circ \phi \circ u_b^{-1}$$

and we use that $\mathcal{F}_{\text{all}}(f)$ is an equivalence of categories for the second isomorphism.

Note that g preserves marked morphisms since u_b and $u_{b'}$ are marked and $\mathcal{F}_{\text{all}}(f)$ induces a bijection between the subsets of marked morphisms (since f^+ is assumed to be an equivalence). Furthermore, since u_b and $u_{b'}$ are unitary (since marked morphisms must be unitary), g is a morphism of $*$ -categories. Finally, in the \mathbb{C} -enriched cases, g is compatible with the enrichments.

The morphism g is the required inverse to f up to marked isomorphism. The transformations u and v can be interpreted as marked isomorphisms

$$u : \text{id}_{\mathbf{B}} \rightarrow f \circ g, \quad v : \text{id}_{\mathbf{A}} \rightarrow g \circ f.$$

We now show 2. If f is a unitary equivalence, then there is an inverse morphism $g : \mathbf{B} \rightarrow \mathbf{A}$ up to unitary isomorphism. Then $\mathcal{F}_{\text{all}}(g)$ and $\text{ma}(g)^+$ are inverse equivalences of $\mathcal{F}_{\text{all}}(f)$ and $\text{ma}(f)^+$, respectively.

We now assume that $\mathcal{F}_{\text{all}}(f)$ and $\text{ma}(f)^+$ are equivalences of categories. Then by the first case 1 we know that $\text{ma}(f) : \text{ma}(\mathbf{A}) \rightarrow \text{ma}(\mathbf{B})$ is a marked equivalence. Let $g : \text{ma}(\mathbf{B}) \rightarrow \text{ma}(\mathbf{A})$ be an inverse of $\text{ma}(f)$ up to marked isomorphism. Then $\mathcal{F}_+(g) : \mathbf{B} \rightarrow \mathbf{A}$ (\mathcal{F}_+ forgets the marking, see (13)) is an inverse of f up to unitary isomorphism. ■

5.4. REMARK. In the case $\mathcal{C} = C^*\mathbf{Cat}_1$ it was shown in [Del10, Lemma 4.6] that a morphism $f : \mathbf{A} \rightarrow \mathbf{B}$ is a unitary equivalence if and only if $\mathcal{F}_{\text{all}}(f)$ is an equivalence of categories, i.e., that the second condition in Lemma 5.3.2 involving $\text{ma}(f)^+$ is redundant. The argument uses a special property of C^* -categories, namely the existence of polar decompositions of morphisms [Del10, Prop. 2.6].

The following lemma about maximal norms morally belongs to Section 2 but can only be stated at this place since it involves the notion of unitary equivalences.

5.5. LEMMA. *If $\Phi : \mathbf{A} \rightarrow \mathbf{B}$ is a unitary equivalence between \mathbb{C} -linear $*$ -categories, then for every morphism f in \mathbf{A} we have $\|\Phi(f)\|_{\max} = \|f\|_{\max}$.*

PROOF. By precomposition with Φ every representation of \mathbf{B} in a C^* -algebra yields a representation of \mathbf{A} in the same C^* -algebra. This immediately implies the inequality

$$\|\Phi(f)\|_{\max} \leq \|f\|.$$

Let now $\Psi : \mathbf{B} \rightarrow \mathbf{A}$ be an inverse equivalence. Then there exists a unitary morphism u in \mathbf{A} such that $u \circ \Psi(\Phi(f)) \circ u^* = f$. This gives (using $\|u\|_{\max} \leq 1$, see Remark 4.8)

$$\|f\|_{\max} = \|u \circ \Psi(\Phi(f)) \circ u^*\|_{\max} \leq \|\Psi(\Phi(f))\|_{\max} \leq \|\Phi(f)\|_{\max}.$$

■

5.6. **REMARK.** We will use the following fact about adjunctions. We consider two relative categories $(\mathcal{C}, W_{\mathcal{C}})$ and $(\mathcal{D}, W_{\mathcal{D}})$ and a pair of adjoint functors

$$L : \mathcal{C} \rightleftarrows \mathcal{D} : R . \quad (15)$$

By

$$\ell_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}[W_{\mathcal{C}}^{-1}] , \quad \ell_{\mathcal{D}} : \mathcal{D} \rightarrow \mathcal{D}[W_{\mathcal{D}}^{-1}]$$

we denote the corresponding localization functors, see (2).

We now assume that L and R are compatible with the sets $W_{\mathcal{C}}$ and $W_{\mathcal{D}}$ in the sense that $\ell_{\mathcal{D}} \circ L$ sends the morphisms in $W_{\mathcal{C}}$ to equivalences in $\mathcal{D}[W_{\mathcal{D}}^{-1}]$, and that $\ell_{\mathcal{C}} \circ R$ sends the morphisms in $W_{\mathcal{D}}$ to equivalences in $\mathcal{C}[W_{\mathcal{C}}^{-1}]$. Then the functors L and R descend essentially uniquely to functors

$$\bar{L} : \mathcal{C}[W_{\mathcal{C}}^{-1}] \rightleftarrows \mathcal{D}[W_{\mathcal{D}}^{-1}] : \bar{R} .$$

In this case the adjunction $L \dashv R$ naturally induces an adjunction $\bar{L} \dashv \bar{R}$. A reference for these facts is [Cis19, Prop. 7.1.14].⁵

If L comes from a left Quillen functor between combinatorial model categories, then we could also proceed as sketched in [Lur17, Rem. 1.3.4.27].

Let \mathcal{C} be in the list

$$\{ * \mathbf{Cat}_1, {}^*_{\mathbb{C}} \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, {}^* \mathbf{Cat}_1^+, {}^*_{\mathbb{C}} \mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \} ,$$

and let $W_{\mathcal{C}}$ denote the (marked) unitary equivalences in \mathcal{C} as defined in Definition 5.2.

5.7. **DEFINITION.** We define the ∞ -categories

$$\begin{aligned} * \mathbf{Cat} &:= * \mathbf{Cat}_1[W_{* \mathbf{Cat}_1}^{-1}] , & * \mathbf{Cat}^+ &:= * \mathbf{Cat}_1^+[W_{* \mathbf{Cat}_1^+}^{-1}] , \\ {}^*_{\mathbb{C}} \mathbf{Cat} &:= {}^*_{\mathbb{C}} \mathbf{Cat}_1[W_{{}^*_{\mathbb{C}} \mathbf{Cat}_1}^{-1}] , & {}^*_{\mathbb{C}} \mathbf{Cat}^+ &:= {}^*_{\mathbb{C}} \mathbf{Cat}_1^+[W_{{}^*_{\mathbb{C}} \mathbf{Cat}_1^+}^{-1}] , \\ C_{\text{pre}}^* \mathbf{Cat} &:= C_{\text{pre}}^* \mathbf{Cat}_1[W_{C_{\text{pre}}^* \mathbf{Cat}_1}^{-1}] , & C_{\text{pre}}^* \mathbf{Cat}^+ &:= C_{\text{pre}}^* \mathbf{Cat}_1^+[W_{C_{\text{pre}}^* \mathbf{Cat}_1^+}^{-1}] , \\ C^* \mathbf{Cat} &:= C^* \mathbf{Cat}_1[W_{C^* \mathbf{Cat}_1}^{-1}] , & C^* \mathbf{Cat}^+ &:= C^* \mathbf{Cat}_1^+[W_{C^* \mathbf{Cat}_1^+}^{-1}] . \end{aligned}$$

⁵Alternatively, one can use [Lur09, Prop. 5.2.2.8] as follows. Let $u : \text{id}_{\mathcal{C}} \rightarrow R \circ L$ be the unit and $v : L \circ R \rightarrow \text{id}_{\mathcal{D}}$ be the counit of the adjunction (15). Then u induces a transformation

$$\bar{u} : \text{id}_{\mathcal{C}[W_{\mathcal{C}}^{-1}]} \rightarrow \bar{R} \circ \bar{L} .$$

We must show that \bar{u} is a unit transformation in the sense of [Lur09, Def. 5.2.2.7], i.e., that for any two objects C in $\mathcal{C}[W_{\mathcal{C}}^{-1}]$ and D in $\mathcal{D}[W_{\mathcal{D}}^{-1}]$ the induced morphism

$$\text{Map}_{\mathcal{D}[W_{\mathcal{D}}^{-1}]}(\bar{L}(C), D) \xrightarrow{\bar{R}} \text{Map}_{\mathcal{C}[W_{\mathcal{C}}^{-1}]}(\bar{R}(\bar{L}(C)), \bar{R}(D)) \xrightarrow{\bar{u}(C)} \text{Map}_{\mathcal{C}[W_{\mathcal{C}}^{-1}]}(C, \bar{R}(D))$$

is an equivalence of spaces. Using the fact that \bar{u} and $\bar{v} : \bar{L} \circ \bar{R} \rightarrow \text{id}_{\mathcal{D}[W_{\mathcal{D}}^{-1}]}$ induced by v satisfy the triangle identities up to equivalence we see that the desired inverse equivalence is given by

$$\text{Map}_{\mathcal{C}[W_{\mathcal{C}}^{-1}]}(C, \bar{R}(D)) \xrightarrow{\bar{L}} \text{Map}_{\mathcal{D}[W_{\mathcal{D}}^{-1}]}(\bar{L}(C), \bar{L}(\bar{R}(D))) \xrightarrow{\bar{v}(D)} \text{Map}_{\mathcal{D}[W_{\mathcal{D}}^{-1}]}(\bar{L}(C), D) .$$

5.8. LEMMA. *The adjunctions (8) induce adjunctions*

$$\text{Lin}_{\mathbb{C}} : {}^*\mathbf{Cat} \rightleftarrows {}^*\mathbb{C}\mathbf{Cat} : \mathcal{F}_{\mathbb{C}} , \quad \text{Lin}_{\mathbb{C}} : {}^*\mathbf{Cat}^+ \rightleftarrows {}^*\mathbb{C}\mathbf{Cat}^+ : \mathcal{F}_{\mathbb{C}} \quad (16)$$

PROOF. We observe that the forgetful functor $\mathcal{F}_{\mathbb{C}}$ and the linearization functor $\text{Lin}_{\mathbb{C}}$ preserve (marked) unitary equivalences. Hence they descend naturally to the ∞ -categories. By Remark 5.6 we obtain an adjunction between these descended functors. ■

5.9. LEMMA. *The adjunctions (10) induce adjunctions*

$$\text{Compl} : C_{\text{pre}}^* \mathbf{Cat} \rightleftarrows C^* \mathbf{Cat} : \mathcal{F}_- , \quad \text{Compl} : C_{\text{pre}}^* \mathbf{Cat}^+ \rightleftarrows C^* \mathbf{Cat}^+ : \mathcal{F}_- \quad (17)$$

PROOF. We first observe that the forgetful functor \mathcal{F}_- and the completion functor Compl preserve (marked) unitary equivalences. Hence they descend naturally to the ∞ -categories. By Remark 5.6 we obtain an adjunction between these descended functors. ■

5.10. LEMMA. *The adjunctions (12) induce adjunctions*

$$\mathcal{F}_{\text{pre}} : C_{\text{pre}}^* \mathbf{Cat} \rightleftarrows {}^*\mathbb{C}\mathbf{Cat} : \text{Bd}^{\infty} , \quad \mathcal{F}_{\text{pre}} : C_{\text{pre}}^* \mathbf{Cat}^+ \rightleftarrows {}^*\mathbb{C}\mathbf{Cat}^+ : \text{Bd}^{\infty} . \quad (18)$$

PROOF. The forgetful functor \mathcal{F}_{pre} preserves (marked) unitary equivalences. The operation Bd^{∞} also preserves (marked) unitary equivalences since $\text{Bd}^{\infty}(\mathbf{A})$ contains all unitary (marked, resp.) morphisms of \mathbf{A} . Hence both functors descend naturally to the ∞ -categories. By Remark 5.6 we obtain an adjunction between these descended functors. ■

5.11. CONVENTION. We use the same notation ℓ for all the localization functors: For \mathcal{C} in the list

$$\{ {}^*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \}$$

we write

$$\ell : \mathcal{C} \rightarrow \mathcal{C}_{\infty} \quad (19)$$

for the corresponding localization.

6. The tensor and power structure over groupoids

In this section we let \mathbf{G} be a category. Later we will assume that it is a groupoid.

For a category \mathbf{A} we consider the functor category $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ whose objects are functors from \mathbf{G} to \mathbf{A} , and whose morphisms are natural transformations between functors. If \mathbf{A} is a $*$ -category, then we define an involution

$$* : \mathcal{F}un(\mathbf{G}, \mathbf{A}) \rightarrow \mathcal{F}un(\mathbf{G}, \mathbf{A})$$

such that it sends a morphism $f = (f_g)_{g \in \mathbf{G}} : a \rightarrow a'$ in $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ with $f_g : a(g) \rightarrow a'(g)$ to the morphism $f^* := (f_g^*)_{g \in \mathbf{G}} : a' \rightarrow a$.

Assume furthermore that \mathbf{A} is a \mathbb{C} -linear $*$ -category. Then the enrichment of \mathbf{A} over complex vector spaces naturally induces an enrichment of $\mathcal{F}un(\mathbf{G}, \mathbf{A})$. In this case $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ has the structure of a \mathbb{C} -linear $*$ -category. If \mathbf{A} is marked, then $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ is a marked $*$ -category or marked \mathbb{C} -linear $*$ -category whose marked morphisms are those transformations $(f_g)_{g \in G}$ where f_g is marked for all g in G .

For \mathcal{C} in the list

$$\{*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathbb{C}}\mathbf{Cat}_1, *_{\mathbb{C}}\mathbf{Cat}_1^+\}$$

we therefore get a functor

$$\mathcal{F}un(-, -) : \mathbf{Cat}_1^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C} .$$

Let \mathbf{A} be a (marked) $*$ -category.

6.1. DEFINITION. We call a functor a in $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ (marked) unitary if $a(\phi)$ is unitary (marked) for all morphisms ϕ in $\text{Hom}_{\mathbf{G}}(g, h)$.

We let $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ denote the full subcategory of $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ of unitary functors. It is a $*$ -category by Remark 2.16. Similarly, if \mathbf{A} is a \mathbb{C} -linear $*$ -category, then so is $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$.

If \mathbf{A} is marked, then we let $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ denote the full subcategory of $\mathcal{F}un(\mathbf{G}, \mathbf{A})$ of marked functors. It is again a marked $*$ -category. If \mathbf{A} is a marked \mathbb{C} -linear $*$ -category, then so is $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$.

Note that a functor in $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ sends all morphisms of \mathbf{G} to invertible morphisms in \mathbf{A} . This observation will be used e.g. in Remark 6.8 below.

For \mathcal{C} in the list

$$\{*\mathbf{Cat}_1, *_{\mathbb{C}}\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathbb{C}}\mathbf{Cat}_1^+\}$$

we have a functor

$$\mathcal{F}un^u(-, -) : \mathbf{Cat}_1^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C} .$$

6.2. REMARK. In the marked case the notation $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ is actually an abuse of notation since this could also be interpreted as the category of unitary functors between \mathbf{G} and \mathbf{A} after forgetting the marking. But we prefer to use this notation with the interpretation as above in order to state formulas below in a form which applies to the unmarked as well as to the marked cases.

If \mathbf{A} is a (marked) pre- C^* -category or a (marked) C^* -category, then as a convention, in order to form the (marked) \mathbb{C} -linear $*$ -category $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ we will consider \mathbf{A} as a (marked) \mathbb{C} -linear category and interpret $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ as a (marked) \mathbb{C} -linear $*$ -category.

6.3. REMARK. If \mathbf{C} is a C^* -category, then we can not expect that $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ is a C^* -category again. For the simplest counter example let \mathbf{G} be an infinite set and \mathbf{A} be the category associated to a C^* -algebra A . Then $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ is the \mathbb{C} -linear $*$ -category with one object and with morphisms $\prod_{\mathbf{G}} A$. But this is not even a pre- C^* -category. In order to

get a C^* -category, for the morphisms we should take the uniformly bounded sequences $\prod_{g \in \mathbf{G}}^{\text{bd}} A$. So we must define a uniformly bounded subfunctor

$$\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A}) \subseteq \mathcal{F}un^u(\mathbf{G}, \mathbf{A}) .$$

6.4. DEFINITION. For a (marked) \mathbb{C} -linear $*$ -category \mathbf{A} we define the uniformly bounded subfunctor by

$$\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A}) := \text{Bd}^\infty(\mathcal{F}un^u(\mathbf{G}, \mathbf{A})) .$$

By definition $\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})$ is a (marked) pre- C^* -category. In particular, for \mathcal{C} in the list $\{C_{\text{pre}}^* \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1^+\}$ we have defined a functor

$$\mathcal{F}un^{\text{bd}}(-, -) : \mathbf{Cat}_1^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C} .$$

6.5. EXAMPLE. We have $\mathcal{F}un^{\text{bd}}(\Delta_{\mathbf{Cat}_1}^0, \mathbf{A}) \cong \text{Bd}^\infty(\mathbf{A})$.

6.6. EXAMPLE. Assume that \mathbf{G} is a set and that \mathbf{A} is associated to a C^* -algebra A . Then $\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})$ can be identified with the category with one object and the morphisms given by the C^* -algebra $\prod_{g \in \mathbf{G}}^{\text{bd}} A$.

6.7. DEFINITION. If \mathbf{A} is a (marked) \mathbb{C} -linear $*$ -category, then we define the (marked) C^* -category

$$\mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A}) := \text{Compl}(\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})) .$$

In particular, for \mathcal{C} in the list $\{C^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1^+\}$ we have defined a functor

$$\mathcal{F}un^{C^*}(-, -) : \mathbf{Cat}_1^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C} .$$

6.8. REMARK. If \mathbf{A} is a (marked) C^* -category, then in Definition 6.7 one can actually omit the application of the completion functor. Let $\mathbf{G}[\mathbf{G}^{-1}]$ denote the groupoid obtained from \mathbf{G} by universally inverting all morphisms. Then the natural functor $\mathbf{G} \rightarrow \mathbf{G}[\mathbf{G}^{-1}]$ induces an isomorphism

$$\mathcal{F}un^u(\mathbf{G}[\mathbf{G}^{-1}], \mathbf{A}) \xrightarrow{\cong} \mathcal{F}un^u(\mathbf{G}, \mathbf{A})$$

for any (marked) $*$ -category \mathbf{A} . If \mathbf{A} is \mathbb{C} -linear, then we get an isomorphism

$$\mathcal{F}un^{\text{bd}}(\mathbf{G}[\mathbf{G}^{-1}], \mathbf{A}) \xrightarrow{\cong} \mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A}) .$$

If \mathbf{A} is a (marked) C^* -category, then in Corollary 6.17 we will see that $\mathcal{F}un^{\text{bd}}(\mathbf{G}[\mathbf{G}^{-1}], \mathbf{A})$ is a (marked) C^* -category. Consequently, $\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})$ is already a (marked) C^* -category.

6.9. REMARK. If \mathbf{A} is a C^* -category, then by Remark 6.8 $\mathcal{F}un^{bd}(\mathbf{G}, \mathbf{A})$ is again a C^* -category.

We have an obvious alternative candidate $\mathcal{F}un^r(\mathbf{G}, \mathbf{A})$ for the functor C^* -category which is also defined as a subcategory of $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ as follows. The objects of $\mathcal{F}un^r(\mathbf{G}, \mathbf{A})$ are the objects of $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$. Recall that given two functors a, a' in $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$ we have an inclusion

$$\text{Hom}_{\mathcal{F}un^u(\mathbf{G}, \mathbf{A})}(a, a') \subseteq \prod_{g \in \mathbf{G}} \text{Hom}_{\mathbf{A}}(a(g), a'(g)) ,$$

where a family $(f_g)_{g \in \mathbf{G}}$ belongs to this subspace if for every morphism ϕ in $\text{Hom}_{\mathbf{G}}(g, h)$

$$f_h \circ a(\phi) = a'(\phi) \circ f_g . \tag{20}$$

We now define the morphisms of $\mathcal{F}un^r(\mathbf{G}, \mathbf{A})$ by

$$\text{Hom}_{\mathcal{F}un^r(\mathbf{G}, \mathbf{A})}(a, a') := \text{Hom}_{\mathcal{F}un^u(\mathbf{G}, \mathbf{A})}(a, a') \cap \prod_{g \in \mathbf{G}}^{bd} \text{Hom}_{\mathbf{A}}(a(g), a'(g)) .$$

Since the relations (20) are linear and continuous the morphism space $\text{Hom}_{\mathcal{F}un^r(\mathbf{G}, \mathbf{A})}(a, a')$ is a closed linear subspace of the Banach space $\prod_{g \in \mathbf{G}}^{bd} \text{Hom}_{\mathbf{A}}(a(g), a'(g))$ with the norm

$$\|(f_g)_{g \in \mathbf{G}}\| := \sup_{g \in \mathbf{G}} \|f_g\|_{\max} .$$

So $\text{Hom}_{\mathcal{F}un^r(\mathbf{G}, \mathbf{A})}(a, a')$ inherits a Banach space structure. Let now $f \in \text{Hom}_{\mathcal{F}un^r(\mathbf{G}, \mathbf{A})}(a, a')$. Then we have

$$\|f^* \circ f\| = \sup_{g \in \mathbf{G}} \|f_g^* \circ f_g\| = \sup_{g \in \mathbf{G}} \|f_g\| = \|f\| ,$$

i.e., the C^* -identity is satisfied. Similarly we conclude the C^* -inequality 4. It follows from the universal property of Bd^∞ that we have a morphism of C^* -categories

$$\mathcal{F}un^r(\mathbf{G}, \mathbf{A}) \rightarrow \mathcal{F}un^{bd}(\mathbf{G}, \mathbf{A}) .$$

In general we do not know whether this is an isomorphism. But it is an isomorphism if \mathbf{G} is a set and \mathbf{A} is associated to a C^* -algebra, see Example 6.6.

Let \mathbf{G} be a groupoid and \mathbf{A} be a $*$ -category. We consider \mathbf{G} as a $*$ -category in the canonical manner (see Example 2.2) and form the $*$ -category $\mathbf{A} \times \mathbf{G}$ (the existence of the product is ensured by Theorem 8.1). Explicitly, the $*$ -category $\mathbf{A} \times \mathbf{G}$ is the product category with the $*$ -operation

$$(f, \phi)^* := (f^*, \phi^{-1}) . \tag{21}$$

Since this definition involves the inverse of the morphism ϕ of \mathbf{G} it is important to assume that \mathbf{G} is a groupoid.

If f is a unitary morphism in \mathbf{A} and ϕ is any morphism in \mathbf{G} , then (f, ϕ) is a unitary morphism in $\mathbf{A} \times \mathbf{G}$. If \mathbf{A} is a marked $*$ -category, then in the $*$ -category $\mathbf{A} \times \mathbf{G}$ we mark all morphisms of the form (f, ϕ) with f marked in \mathbf{A} and ϕ in \mathbf{G} arbitrary.

For \mathcal{C} in the list $\{*\mathbf{Cat}_1, *\mathbf{Cat}_1^+\}$ we thus have defined a functor

$$- \times - : \mathcal{C} \times \mathbf{Grpd}_1 \rightarrow \mathcal{C} .$$

In the \mathbb{C} -linear case we must modify this construction. For a \mathbb{C} -linear $*$ -category \mathbf{A} and a groupoid \mathbf{G} we define $\mathbf{A} \otimes \mathbf{G}$ to be the category with the objects of $\mathbf{A} \times \mathbf{G}$, and whose morphisms are given by the complex vector spaces

$$\mathrm{Hom}_{\mathbf{A} \otimes \mathbf{G}}((a, g), (a', g')) := \bigoplus_{\phi \in \mathrm{Hom}_{\mathbf{G}}(g, g')} \mathrm{Hom}_{\mathbf{A}}(a, a')$$

with the obvious $*$ -operation and composition. Note that the sum over an empty index set is the zero vector space.

We note that $\mathbf{A} \times \mathbf{G}$ is a wide subcategory of $\mathbf{A} \otimes \mathbf{G}$ in a natural way. If \mathbf{A} is marked, then in $\mathbf{A} \otimes \mathbf{G}$ we again mark all morphisms of the form (f, ϕ) for f a marked morphism in \mathbf{A} and an arbitrary morphism ϕ of \mathbf{G} .

For \mathcal{C} in the list $\{*\mathbb{C}\mathbf{Cat}_1, *\mathbb{C}\mathbf{Cat}_1^+\}$ we thus have defined a functor

$$- \times - : \mathcal{C} \times \mathbf{Grpd}_1 \rightarrow \mathcal{C} .$$

6.10. EXAMPLE. If \mathbf{A} is a (marked) $*$ -category, then we have an isomorphism

$$\mathrm{Lin}_{\mathbb{C}}(\mathbf{A}) \otimes \mathbf{G} \cong \mathrm{Lin}_{\mathbb{C}}(\mathbf{A} \times \mathbf{G}) .$$

6.11. EXAMPLE. If $\mathbf{G} = BH$ for some group H , then the \mathbb{C} -linear $*$ -category $\Delta_{*\mathbb{C}\mathbf{Cat}_1}^0 \otimes BH$ is isomorphic to the \mathbb{C} -linear $*$ -category associated to the group ring $\mathbb{C}[H]$ with its usual involution.

Let \mathbf{G} be a groupoid.

6.12. LEMMA. *If \mathbf{A} is a (marked) pre- C^* -category, then so is $\mathbf{A} \otimes \mathbf{G}$.*

PROOF. It suffices to show that

$$\mathbf{A} \otimes \mathbf{G} = \mathrm{Bd}^\infty(\mathbf{A} \otimes \mathbf{G}) . \tag{22}$$

Every morphism in $\mathbf{A} \otimes \mathbf{G}$ is a finite linear combination of morphisms of the form (f, ϕ) of $\mathbf{A} \times \mathbf{G}$ with $f : a \rightarrow a'$ and $\phi : g \rightarrow g'$. We can decompose $(f, \phi) = (\mathrm{id}_{a'}, \phi) \circ (f, \mathrm{id}_g)$.

Let $\rho : \mathbf{A} \otimes \mathbf{G} \rightarrow B$ be a representation in a C^* -algebra B . Then we can restrict ρ to a representation of $\mathbf{A} \cong \mathbf{A} \times \{g\} \subseteq \mathbf{A} \otimes \mathbf{G}$. We conclude that $\|\rho(f, \mathrm{id}_g)\|_B \leq \|f\|_{\max}$, where $\|-\|_{\max}$ denotes the maximal norm on \mathbf{A} . Furthermore, because of (21) we know that $\rho(\mathrm{id}_{a'}, g)$ is a partial isometry in B . Therefore $\|\rho(f, \phi)\|_B \leq \|f\|_{\max}$. This shows that $(f, g) \in \mathrm{Bd}^\infty(\mathbf{A} \otimes \mathbf{G})$. Hence also all finite linear combinations of such elements belong to $\mathrm{Bd}^\infty(\mathbf{A} \otimes \mathbf{G})$. This shows the desired equality (22). ■

For \mathcal{C} in the list $\{C_{\text{pre}}^* \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1^+\}$ we thus have defined a functor

$$- \otimes - : \mathcal{C} \times \mathbf{Grpd}_1 \rightarrow \mathcal{C} .$$

For a (marked) \mathbb{C} -linear *-category \mathbf{A} and a groupoid \mathbf{G} we define

$$\mathbf{A} \otimes_{\max} \mathbf{G} := \text{Compl}(\text{Bd}^\infty(\mathbf{A} \otimes \mathbf{G})) .$$

If \mathbf{A} was a (marked) pre- C^* -category, then by Lemma 6.12 we can simplify this to

$$\mathbf{A} \otimes_{\max} \mathbf{G} \cong \text{Compl}(\mathbf{A} \otimes \mathbf{G}) .$$

For \mathcal{C} in the list $\{C^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1^+\}$ we thus have defined a functor

$$- \otimes_{\max} - : \mathcal{C} \times \mathbf{Grpd}_1 \rightarrow \mathcal{C} .$$

6.13. EXAMPLE. We consider the C^* -category $\Delta_{C^* \mathbf{Cat}_1}^0$ associated to the C^* -algebra \mathbb{C} . For a groupoid \mathbf{G} the C^* -category $\Delta_{C^* \mathbf{Cat}_1}^0 \otimes_{\max} \mathbf{G}$ is the maximal groupoid C^* -category. In particular, if $\mathbf{G} = BH$ for a group H , then $\Delta_{C^* \mathbf{Cat}_1}^0 \otimes_{\max} BH$ is isomorphic to the C^* -category associated to the maximal group C^* -algebra $C_{\max}^*(H)$.

6.14. CONVENTION. In order to avoid a case-dependent notation we write \sharp for the tensor structures with groupoids \times , \otimes , or \otimes_{\max} in the respective cases. We will furthermore use the notation $\mathcal{F}un^?$, where $?$ is u , bd , or C^* in the respective cases. See Table 1.

Table 1:

case	\sharp	$?$
$* \mathbf{Cat}_1^{(+)}$	\times	u
$*_{\mathbb{C}} \mathbf{Cat}_1^{(+)}$	\otimes	u
$C_{\text{pre}}^* \mathbf{Cat}_1^{(+)}$	\otimes	bd
$C^* \mathbf{Cat}_1^{(+)}$	\otimes_{\max}	C^*

6.15. EXAMPLE. Let \mathcal{C} be in the list

$$\{ * \mathbf{Cat}_1, *_{\mathbb{C}} \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, * \mathbf{Cat}_1^+, *_{\mathbb{C}} \mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \}$$

and recall the morphism classifier object \mathbb{I} in \mathbf{Grpd}_1 from Definition 4.14. Let $f_0, f_1 : \mathbf{C} \rightarrow \mathbf{D}$ be two morphisms in \mathcal{C} . Then we have a bijective correspondence between (marked) unitary isomorphisms $u : f_0 \rightarrow f_1$ and functors $U : \mathbf{C} \sharp \mathbb{I} \rightarrow \mathbf{D}$ with $U \circ \iota_i = f_i$, where $\iota_i : \mathbf{C} \cong \mathbf{C} \sharp \Delta^0 \rightarrow \mathbf{C} \sharp \mathbb{I}$ for $i = 0, 1$ is induced by the objects 0 and 1 of \mathbb{I} . Given U , the (marked) unitary isomorphism u is obtained by $u = (U(\text{id}_c, 0 \rightarrow 1))_{c \in \mathbf{C}}$. Vice versa, given u , we can define U on morphisms by $U(\text{id}_c, 0 \rightarrow 1) = u_b$ and $U(a, \text{id}_0) := f_0(a)$ and compatibility with compositions and $*$.

Let \mathbf{G} be a groupoid and \mathcal{C} be in the list

$$\{ * \mathbf{Cat}_1, *_{\mathbb{C}} \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, * \mathbf{Cat}_1^+, *_{\mathbb{C}} \mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \} .$$

6.16. PROPOSITION. For \mathbf{C} and \mathbf{A} in \mathcal{C} we have a natural exponential law

$$\mathrm{Hom}_{\mathcal{C}}(\mathbf{C} \sharp \mathbf{G}, \mathbf{A}) \cong \mathrm{Hom}_{\mathcal{C}}(\mathbf{C}, \mathcal{F}un^?(\mathbf{G}, \mathbf{A})) .$$

PROOF. We start with giving the natural bijection in the case of $*$ -categories. Let Φ be in $\mathrm{Hom}_{*\mathbf{Cat}_1}(\mathbf{C} \times \mathbf{G}, \mathbf{A})$. The bijection identifies this morphism with a morphism Ψ in $\mathrm{Hom}_{*\mathbf{Cat}_1}(\mathbf{C}, \mathcal{F}un^u(\mathbf{G}, \mathbf{A}))$ given by

$$\Psi(c)(g) := \Phi(c, g) , \quad \Psi(c)(\phi) := \Phi(\mathrm{id}_c, \phi) , \quad \Psi(f) := (\Phi(f, \mathrm{id}_g)_{g \in \mathbf{G}}) .$$

Here c is an object of \mathbf{C} , g is an object of \mathbf{G} , ϕ is a morphism in \mathbf{G} , and f is a morphism in \mathbf{C} . Note that $\Psi(c)$ takes values in unitary functors since

$$\Psi(c)(\phi)^* = \Phi(\mathrm{id}_c, \phi)^* = \Phi(\mathrm{id}_c, \phi^*) = \Phi(\mathrm{id}_c, \phi^{-1}) = \Psi(c)(\phi)^{-1} .$$

Vice versa let the morphism Ψ be given. Then the bijection sends it to the morphism Φ given by

$$\Phi(c, g) := \Psi(c)(g) , \quad \Phi(f, \phi) := \Psi(c')(\phi) \circ \Psi(f)(g) ,$$

where $f : c \rightarrow c'$ is a morphism in \mathbf{C} and $\phi : g \rightarrow g'$ is a morphism in \mathbf{G} .

The same formulas work in the \mathbb{C} -linear case for morphisms of the form (f, ϕ) . The maps are then extended linearly. In the case of pre- C^* -categories we must check that Ψ takes values in the subfunctor $\mathcal{F}un^{\mathrm{bd}}(\mathbf{G}, \mathbf{A})$ of $\mathcal{F}un^u(\mathbf{G}, \mathbf{A})$. But this is clear since Ψ is a morphism between \mathbb{C} -linear categories, \mathbf{C} is a pre- C^* -category, and the universal property 2 of Bd^∞ .

We finally consider the case of C^* -categories. In this case we could cite [Del10]. Here is the argument. We first observe that for a C^* -category \mathbf{A} and pre- C^* -category \mathbf{C} we have a bijection

$$\mathrm{Hom}_{C^*_{\mathrm{pre}}\mathbf{Cat}_1}(\mathbf{C} \otimes \mathbf{G}, \mathcal{F}_-(\mathbf{A})) \cong \mathrm{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{C} \otimes_{\max} \mathbf{G}, \mathbf{A}) \quad (23)$$

by the universal property of the completion. We can use this bijection, the already verified bijection

$$\mathrm{Hom}_{C^*_{\mathrm{pre}}\mathbf{Cat}_1}(\mathbf{C} \otimes \mathbf{G}, \mathcal{F}_-(\mathbf{A})) \cong \mathrm{Hom}_{C^*_{\mathrm{pre}}\mathbf{Cat}_1}(\mathbf{C}, \mathcal{F}un^{\mathrm{bd}}(\mathbf{G}, \mathcal{F}_-(\mathbf{A}))) ,$$

and the completion map

$$\mathrm{Hom}_{C^*_{\mathrm{pre}}\mathbf{Cat}_1}(\mathbf{C}, \mathcal{F}un^{\mathrm{bd}}(\mathbf{G}, \mathcal{F}_-(\mathbf{A}))) \rightarrow \mathrm{Hom}_{C^*_{\mathrm{pre}}\mathbf{Cat}_1}(\mathbf{C}, \mathcal{F}_-(\mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A}))) \quad (24)$$

in order to produce a Ψ from a given Φ .

For the inverse we note that since \mathbf{A} is complete, the evaluation functors

$$e_g : \mathcal{F}un^{\mathrm{bd}}(\mathbf{G}, \mathcal{F}_-(\mathbf{A})) \rightarrow \mathbf{A}$$

for g in \mathbf{G} extend to the completion and provide functors

$$\bar{e}_g : \mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A}) \rightarrow \mathbf{A} .$$

Hence the formula which expresses Φ in terms of Ψ makes sense. It defines, by linear extension, an element in $\text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(\mathbf{C} \otimes \mathbf{G}, \mathcal{F}_-(\mathbf{A}))$ which gives the desired Φ in $\text{Hom}_{C^*\text{Cat}_1}(\mathbf{C} \otimes_{\text{max}} \mathbf{G}, \mathbf{A})$ using the isomorphism (23) above.

In the marked case we just observe the following. If Φ is a functor between marked categories, then Ψ takes values in marked functors, and vice versa, if Ψ has this property, then Φ preserves marked morphisms. ■

Let \mathbf{G} be a groupoid.

6.17. COROLLARY. *If \mathbf{A} is a (marked) C^* -category, then the completion morphism*

$$\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathcal{F}_-(\mathbf{A})) \rightarrow \mathcal{F}_-(\mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A}))$$

is an isomorphism and $\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})$ is a (marked) C^ -category.*

PROOF. In the proof of Proposition 6.16 we have actually shown that for every pre- C^* -category \mathbf{C} there is a natural isomorphism

$$\text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(\mathbf{C} \otimes_{\text{max}} \mathbf{G}, \mathcal{F}_-(\mathbf{A})) \cong \text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(\mathbf{C}, \mathcal{F}_-(\mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A}))) .$$

Furthermore we have a natural isomorphism

$$\text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(\mathbf{C} \otimes_{\text{max}} \mathbf{G}, \mathbf{A}) \cong \text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(\mathbf{C}, \mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})) .$$

By an inspection of the construction of these bijections we see that the resulting isomorphism

$$\text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(-, \mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A})) \cong \text{Hom}_{C^*_{\text{pre}}\text{Cat}_1}(-, \mathcal{F}_-(\mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A})))$$

of functors $C^*_{\text{pre}}\text{Cat}_1^{\text{op}} \rightarrow \mathbf{Set}$ is induced by the completion morphisms

$$\mathcal{F}un^{\text{bd}}(\mathbf{G}, \mathbf{A}) \rightarrow \mathcal{F}_-(\mathcal{F}un^{C^*}(\mathbf{G}, \mathbf{A})) .$$

By the Yoneda Lemma it is therefore an isomorphism, too. ■

In the following we introduce the fundamental groupoid functor $\Pi : \mathbf{sSet} \rightarrow \mathbf{Grpd}_1$.

6.18. DEFINITION. *The fundamental groupoid functor is defined as a left-adjoint of an adjunction*

$$\Pi : \mathbf{sSet} \rightleftarrows \mathbf{Grpd}_1 : \mathbf{N} ,$$

where \mathbf{N} takes the nerve of a groupoid.

Explicitly, the fundamental groupoid $\Pi(K)$ of a simplicial set K is the groupoid freely generated by the path category $P(K)$ of K . The category $P(K)$ in turn is given as follows:

1. The objects of $P(K)$ are the 0-simplices.
2. The morphisms of $P(K)$ are generated by the 1-simplices of K subject to the relation $g \circ f \sim h$ if there exists a 2-simplex σ in K with $d_2\sigma = f$, $d_0\sigma = g$ and $d_1\sigma = h$.

In the following definition we use the notation introduced in Convention 6.14. Let \mathcal{C} be a member of the list

$$\{^*\text{Cat}_1, {}^*\text{Cat}_1, C^*_{\text{pre}}\text{Cat}_1, C^*\text{Cat}_1, {}^*\text{Cat}_1^+, {}^*\text{Cat}_1^+, C^*_{\text{pre}}\text{Cat}_1^+, C^*\text{Cat}_1^+\} .$$

6.19. DEFINITION. We define the tensor and cotensor structure of \mathcal{C} with \mathbf{sSet} by

$$\mathcal{C} \times \mathbf{sSet} \rightarrow \mathcal{C}, \quad (\mathbf{A}, K) \mapsto \mathbf{A} \sharp K := \mathbf{A} \sharp \Pi(K)$$

and

$$\mathbf{sSet}^{\text{op}} \times \mathcal{C} \rightarrow \mathcal{C}, \quad (K, \mathbf{B}) \mapsto \mathbf{B}^K := \mathcal{F}un^?(\Pi(K), \mathbf{B}).$$

For objects \mathbf{A}, \mathbf{B} of \mathcal{C} we define the simplicial mapping space $\text{Map}(\mathbf{A}, \mathbf{B})$ in \mathbf{sSet} by

$$\text{Map}(\mathbf{A}, \mathbf{B})[-] := \text{Map}(\mathbf{A} \sharp \Delta^-, \mathbf{B}).$$

Then for every two objects \mathbf{A}, \mathbf{B} in \mathcal{C} and every simplicial set K , by Proposition 6.16 we have natural bijections

$$\text{Hom}_{\mathbf{sSet}}(K, \text{Map}(\mathbf{A}, \mathbf{B})) \cong \text{Hom}_{\mathcal{C}}(\mathbf{A} \sharp K, \mathbf{B}) \cong \text{Hom}_{\mathcal{C}}(\mathbf{A}, \mathbf{B}^K). \quad (25)$$

In this way we have defined a simplicial enrichment of \mathcal{C} .

7. The resolution

Let G be a group. In the present section we consider the functor categories from the arrow category \tilde{G} . These functor categories will serve as explicit fibrant resolutions later in Section 13.

A G -groupoid (or G -category) is a groupoid (or category) with a strict action of G , i.e., an object of $\mathbf{Fun}(BG, \mathbf{Grpd}_1)$ (or $\mathbf{Fun}(BG, \mathbf{Cat}_1)$).

7.1. DEFINITION. The arrow category \tilde{G} is the G -groupoid defined as follows:

1. The set of objects of \tilde{G} is the underlying set of G .
2. For every pair of objects g, h of \tilde{G} the set of morphisms $\text{Hom}_{\tilde{G}}(g, h)$ consists of one point which we will denote by $g \rightarrow h$. The composition of morphisms is defined in the only possible way.
3. The group G acts on the groupoid \tilde{G} by left multiplication.

For a G -category \mathbf{A} the functor category $\mathcal{F}un(\tilde{G}, \mathbf{A})$ is again a G -category. The group G acts on this functor category as follows. If $a : \tilde{G} \rightarrow \mathbf{A}$ is a functor, then we set $g(a) := g \circ a \circ g^{-1}$. The action on morphisms is similar. We interpret this construction as a functor

$$\mathcal{F}un(\tilde{G}, -) : \mathbf{Fun}(BG, \mathbf{Cat}_1) \rightarrow \mathbf{Fun}(BG, \mathbf{Cat}_1).$$

This construction extends to the various versions of (marked) $*$ -categories. For \mathcal{C} in the list

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}$$

we get a functor

$$\mathcal{F}un^?(\tilde{G}, -) : \mathbf{Fun}(BG, \mathcal{C}) \rightarrow \mathbf{Fun}(BG, \mathcal{C}),$$

see Convention 6.14 for notation.

7.2. LEMMA. *If \mathbf{A} is a (marked) pre- C^* -category, then the natural morphism*

$$\mathcal{F}un^{\text{bd}}(\tilde{G}, \mathbf{A}) \rightarrow \mathcal{F}un^u(\tilde{G}, \mathbf{A})$$

*is an isomorphism of (marked) \mathbb{C} -linear *-categories.*

PROOF. We must show that every morphism in $\mathcal{F}un^u(\tilde{G}, \mathbf{A})$ has a finite maximal norm. If a is a unitary functor from \tilde{G} to \mathbf{A} , then we have a unitary isomorphism

$$h_a := (a(1 \rightarrow g))_{g \in \tilde{G}} : \text{const}(a(1)) \rightarrow a ,$$

where $\text{const}(a(1))$ denotes the constant functor with value $a(1)$. Unitarity implies the norm estimates $\|h_a\|_{\max} \leq 1$ and $\|h_a^{-1}\|_{\max} \leq 1$, see Remark 4.8.

If $f : a \rightarrow a'$ is a morphism in $\mathcal{F}un^u(\tilde{G}, \mathbf{A})$, then we have the relation.

$$h_{a'} \circ \text{const}(f(1)) \circ h_a^{-1} = f .$$

This implies the inequality

$$\|f\|_{\max} \leq \|\text{const}(f(1))\|_{\max} \leq \|f(1)\|_{\max} .$$

■

Using Corollary 6.17 we conclude:

7.3. COROLLARY. *For a (marked) C^* -category \mathbf{A} the natural maps are isomorphisms*

$$\mathcal{F}un^{C^*}(\tilde{G}, \mathbf{A}) \xleftarrow{\cong} \mathcal{F}un^{\text{bd}}(\tilde{G}, \mathbf{A}) \xrightarrow{\cong} \mathcal{F}un^u(\tilde{G}, \mathbf{A})$$

*isomorphism of (marked) \mathbb{C} -linear *-categories.*

7.4. REMARK. In view of Corollary 7.3 and Lemma 7.2 we have an isomorphism of (marked) \mathbb{C} -linear *-categories.

$$\mathcal{F}un^?(\tilde{G}, \mathbf{A}) \cong \mathcal{F}un^u(\tilde{G}, \mathbf{A})$$

in all cases, see Convention 6.14 and Remark 6.2 for the usage of notation.

We have a G -equivariant version of the exponential law. Let \mathcal{C} be in the list

$$\{*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

7.5. PROPOSITION. *For \mathbf{A} and \mathbf{C} in $\mathbf{Fun}(BG, \mathcal{C})$ we have a natural isomorphism*

$$\text{Hom}_{\mathbf{Fun}(BG, \mathcal{C})}(\mathbf{C} \sharp \tilde{G}, \mathbf{A}) \cong \text{Hom}_{\mathbf{Fun}(BG, \mathcal{C})}(\mathbf{C}, \mathcal{F}un^?(\tilde{G}, \mathbf{A})) .$$

PROOF. This follows from an inspection of the proof of Proposition 6.16.

■

8. Completeness, cocompleteness, and local presentability

The first goal of this section is to show the following theorem.

8.1. THEOREM. *The categories in the list*

$$\{*\mathbf{Cat}_1, {}_c\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}_c\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}$$

are complete and cocomplete.

8.2. REMARK. The case of C^* -categories is due to [Del10]. For completeness of the presentation we reprove this case together with the others.

8.3. REMARK. In the proof we use that the categories of categories and marked categories \mathbf{Cat}_1 and \mathbf{Cat}_1^+ are complete and cocomplete. In particular we must understand colimits in \mathbf{Cat}_1 in some detail. Let $\mathbf{DirGraph}$ denote the category of directed graphs. Then we have an adjunction

$$\text{Free}_{\text{Cat}} : \mathbf{DirGraph} \rightleftarrows \mathbf{Cat}_1 : \mathcal{F}_\circ, \quad (26)$$

where \mathcal{F}_\circ sends a category to its underlying directed graph (i.e., forgets the composition), and Free_{Cat} sends a directed graph to the category freely generated by it. Colimits in $\mathbf{DirGraph}$ are formed by taking the colimits of the sets of vertices and edges separately.

If \mathbf{C} is a category, then the counit of the adjunction (26) provides a functor

$$\text{Free}_{\text{Cat}}(\mathcal{F}_\circ(\mathbf{C})) \rightarrow \mathbf{C}. \quad (27)$$

It is a bijection on objects. We $R(\mathbf{C})$ denote the equivalence relation on the morphisms of $\text{Free}_{\text{Cat}}(\mathcal{F}_\circ(\mathbf{C}))$ generated by the action of the functor (27) on morphisms. This relation is compatible with the category structure and the functor (27) induces an isomorphism

$$\text{Free}_{\text{Cat}}(\mathcal{F}_\circ(\mathbf{C}))/R(\mathbf{C}) \cong \mathbf{C}.$$

We now consider a diagram $\mathbf{A} : I \rightarrow \mathbf{Cat}_1$. Then we form the category $\text{Free}_{\text{Cat}}(\text{colim}_I \mathcal{F}_\circ(\mathbf{A}))$. On the morphisms of this category we consider the smallest equivalence relation R compatible with the category structure which contains the images of $R(\mathbf{A}(i))$ under the canonical maps

$$\text{Mor}(\text{Free}_{\text{Cat}}(\mathcal{F}_\circ(\mathbf{A}(i)))) \rightarrow \text{Mor}(\text{Free}_{\text{Cat}}(\text{colim}_I \mathcal{F}_\circ(\mathbf{A})))$$

for all i in I . Then one can check that

$$\text{Free}_{\text{Cat}}(\text{colim}_I \mathcal{F}_\circ(\mathbf{A}))/R \cong \text{colim}_I \mathbf{A}.$$

We will in particular need the following conclusion of this discussion: Every morphism in $\text{colim}_I \mathbf{A}$ is a finite composition of morphisms of the form $u_i(f_i)$ where, for i in I , $u_i : \mathbf{A}(i) \rightarrow \text{colim}_I \mathbf{A}$ is the canonical functor and f_i is in $\text{Mor}(\mathbf{A}(i))$.

PROOF. (of Theorem 8.1) We first show completeness. Our starting point is the fact that the 1-category \mathbf{Cat}_1 is complete. If $\mathbf{A} : I \rightarrow \mathbf{Cat}_1$ is a diagram of categories, then we have canonical isomorphisms of sets

$$\text{Ob}(\lim_I \mathbf{A}) \cong \lim_I \text{Ob}(\mathbf{A}) , \quad \text{Mor}(\lim_I \mathbf{A}) \cong \lim_I \text{Mor}(\mathbf{A}) . \tag{28}$$

Next we consider $^*\mathbf{Cat}_1$. Because of the adjunction (7) a limit in $^*\mathbf{Cat}_1$ (provided it exists) can be calculated as follows. One first calculates the limit in \mathbf{Cat}_1 (i.e., after forgetting the *-operation), and then restores the *-operation using the functoriality of the limit. Thus let $\mathbf{A} : I \rightarrow ^*\mathbf{Cat}_1$ be a diagram. We choose a category \mathbf{B} together with a morphism $\text{const}(\mathbf{B}) \rightarrow \mathcal{F}_*(\mathbf{A})$ which exhibits \mathbf{B} as a representative of the limit $\lim_I \mathcal{F}_*(\mathbf{A})$. Then we can take \mathbf{B}^{op} and $\text{const}(\mathbf{B})^{\text{op}} \rightarrow \mathcal{F}_*(\mathbf{A})^{\text{op}}$ in order to exhibit the limit $\lim_I \mathcal{F}_*(\mathbf{A})^{\text{op}}$. The transformation $* : \mathcal{F}_*(\mathbf{A}) \rightarrow \mathcal{F}_*(\mathbf{A})^{\text{op}}$ now determines a unique functor $* : \mathbf{B} \rightarrow \mathbf{B}^{\text{op}}$ such that

$$\begin{array}{ccc} \text{const}(\mathbf{B}) & \xrightarrow{\text{const}(*)} \text{const}(\mathbf{B}^{\text{op}}) & \xrightarrow{\cong} \text{const}(\mathbf{B})^{\text{op}} \\ \downarrow & & \downarrow \\ \mathcal{F}_*(\mathbf{A}) & \xrightarrow{*} & \mathcal{F}_*(\mathbf{A})^{\text{op}} \end{array}$$

commutes. This functor is the identity on objects and turns \mathbf{B} into a *-category. Together with the morphism of diagrams of *-categories $\text{const}(\mathbf{B}) \rightarrow \mathbf{A}$ it represents the limit $\lim_I \mathbf{A}$ in $^*\mathbf{Cat}_1$.

In order to deal with the marked case we observe that \mathbf{Cat}_1^+ is complete. If $\mathbf{A} : I \rightarrow \mathbf{Cat}_1^+$ is a diagram of marked categories, then in order to construct the marked category $\lim_I \mathbf{A}$, using the forgetful functor \mathcal{F}_+ from (13) we form the category $\lim_I \mathcal{F}_+(\mathbf{A})$ and mark the morphisms which are the elements of $\text{Mor}(\lim_I \mathcal{F}_+(\mathbf{A}))$ whose evaluation (use (28)) at all objects i of I are marked. We can now repeat the constructions above with \mathbf{A} and \mathbf{B} marked.

The same idea works for a diagram $\mathbf{A} : I \rightarrow ^*\mathbb{C}\mathbf{Cat}_1$ of \mathbb{C} -linear *-categories. Since the forgetful functor $\mathcal{F}_{\mathbb{C}}$ from (8) is the right-adjoint of the adjunction it preserves limits. Hence we have

$$\mathcal{F}_{\mathbb{C}}(\lim_I \mathbf{A}) \cong \lim_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})$$

if the limit on the left-hand side exists. The limit on the right-hand side is interpreted in $^*\mathbf{Cat}_1$ and exists as we have seen above. In the following we argue that limit on the left-hand side indeed exists. For a diagram $\mathbf{A} : I \rightarrow ^*\mathbb{C}\mathbf{Cat}_1$ we first define the object $\lim_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})$ in $^*\mathbf{Cat}_1$, then observe that the limit has an induced complex enrichment. We then observe that the canonical morphism

$$\text{const}(\lim_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})) \rightarrow \mathbf{A}$$

is compatible with the enrichment and exhibits the *-category $\lim_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})$ together with the enrichment as the limit $\lim_I \mathbf{A}$.

The same argument applies in the marked case.

Let \mathcal{F}_{pre} and Bd^∞ be the functors as in (12). For a diagram of pre- C^* -categories $\mathbf{A} : I \rightarrow C^*_{\text{pre}} \mathbf{Cat}_1$ we have an isomorphism

$$\lim_I \mathbf{A} \cong \text{Bd}^\infty(\lim_I \mathcal{F}_{\text{pre}}(\mathbf{A})) , \tag{29}$$

where the limit on the right-hand side is interpreted in ${}^*C \mathbf{Cat}_1$. This immediately follows from the adjunction (12) since \mathcal{F}_{pre} is a fully faithful inclusion. The same argument applies in the marked case.

Let \mathcal{F}_- be the forgetful functor from (10). Since it is the right-adjoint of an adjunction it is clear that it preserves limits. We consider a diagram of C^* -categories $\mathbf{A} : I \rightarrow C^* \mathbf{Cat}_1$. Then we have

$$\mathcal{F}_-(\lim_I \mathbf{A}) \simeq \lim_I \mathcal{F}_-(\mathbf{A})$$

if the limit $\lim_I \mathbf{A}$ exists. The limit on the right-hand side is interpreted in $C^*_{\text{pre}} \mathbf{Cat}_1$ and exists as seen above. We now argue that the limit on the left-hand side indeed exists. Every limit can be expressed as a finite combination of equalizers and products. It therefore suffices to show that $\lim_I \mathcal{F}_-(\mathbf{A})$ is a C^* -category in the case that I is a set or a finite category.

If I is a set, then the limit is represented by the bounded product

$$\lim_I \mathbf{A} \cong \prod_{i \in I}^{\text{bd}} \mathbf{A}(i) .$$

Indeed, this product has the universal property since morphisms between C^* -categories are norm-bounded by 1.

If I is finite, then we realize $\lim_I \mathbf{A}$ as a subcategory of $\prod_I \mathbf{A}(i)$ cut out by linear $*$ -invariant continuous equations. This is again C^* -category.

We now consider the marked case. If $\mathbf{A} : I \rightarrow C^* \mathbf{Cat}_1^+$ is a diagram of marked C^* -categories, then $\lim_I \mathcal{F}_-(\mathbf{A})$ is a marked pre- C^* -category. Above we have seen that it is also a marked C^* -category which then necessarily represents $\lim_I \mathbf{A}$.

This finishes the proof of completeness in all cases.

We now show cocompleteness.

We start with the cocompleteness of ${}^* \mathbf{Cat}_1$. In the argument we employ the fact that \mathbf{Cat}_1 is cocomplete, see Remark 8.3. Let $\mathbf{A} : I \rightarrow {}^* \mathbf{Cat}_1$ be a diagram. Then we choose a category \mathbf{B} together with a morphism $\mathcal{F}_*(\mathbf{A}) \rightarrow \text{const}(\mathbf{B})$ which exhibits \mathbf{B} as a colimit $\text{colim}_I \mathcal{F}_*(\mathbf{A})$, where \mathcal{F}_* is as in (7). Then we can take B^{op} and $\mathcal{F}_*(\mathbf{A})^{\text{op}} \rightarrow \text{const}(\mathbf{B})^{\text{op}}$ as a representative of the colimit $\text{colim}_I \mathcal{F}_*(\mathbf{A})^{\text{op}}$. Similarly as in the case of limits, the transformation $*$: $\mathcal{F}_*(\mathbf{A}) \rightarrow \mathcal{F}_*(\mathbf{A})^{\text{op}}$ now induces a functor $*$: $\mathbf{B} \rightarrow \mathbf{B}^{\text{op}}$ in the canonical way such that it is the identity on objects. Hence \mathbf{B} has the structure of a $*$ -category. The canonical morphism of diagrams of $*$ -categories $\mathbf{A} \rightarrow \text{const}(\mathbf{B})$ exhibits \mathbf{B} as the colimit $\text{colim}_I \mathbf{A}$.

In the marked case we use that we have already shown that ${}^* \mathbf{Cat}_1$ is cocomplete. If $\mathbf{A} : I \rightarrow {}^* \mathbf{Cat}_1^+$ is a diagram, then in order to construct the marked $*$ -category $\text{colim}_I \mathbf{A}$

we first form the *-category $\mathbf{B} := \text{colim}_I \mathcal{F}_+(\mathbf{A})$, where \mathcal{F}_+ is as in (13). In \mathbf{B} we mark all morphisms which are compositions of morphisms of the form $u_i(f_i)$ for i in I , a marked morphisms f_i in $\mathbf{A}(i)$, and where $u_i : \mathcal{F}_+(\mathbf{A}(i)) \rightarrow \text{colim}_I \mathcal{F}_+(\mathbf{A}) = \mathbf{B}$ is the canonical *-functor. Since u_i is a *-functor we have for a marked f_i that

$$u_i(f_i)^* = u_i(f_i^*) = u_i(f_i^{-1}) = u_i(f_i)^{-1} .$$

This implies that the marked morphisms in \mathbf{B} are unitary. By construction they are closed under composition so that \mathbf{B} is a marked *-category, and $\mathbf{A} \rightarrow \text{const}(\mathbf{B})$ is a morphism of diagrams of marked *-categories. It exhibits \mathbf{B} as the colimit $\text{colim}_i \mathbf{A}$ in ${}^*\mathbf{Cat}_1^+$.

In order to construct colimits in ${}^*\mathbb{C}\mathbf{Cat}_1$ we use the adjunction (8). If \mathbf{C} is a \mathbb{C} -linear *-category, then we have a natural exact sequence of \mathbb{C} -linear *-categories

$$0 \rightarrow R(\mathbf{C}) \rightarrow \text{Lin}_{\mathbb{C}}(\mathcal{F}_{\mathbb{C}}(\mathbf{C})) \rightarrow \mathbf{C} \rightarrow 0 ,$$

with the caveat that $R(\mathbf{C})$ is non-unital. The second map in this sequence is the counit of the adjunction (8). For a diagram $\mathbf{A} : I \rightarrow {}^*\mathbb{C}\mathbf{Cat}_1$ we now define \mathbf{B} as the quotient

$$0 \rightarrow \langle R(\mathbf{A}(i)) \mid i \in I \rangle \rightarrow \text{Lin}_{\mathbb{C}}(\text{colim}_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})) \rightarrow \mathbf{B} \rightarrow 0 ,$$

where $\langle R(\mathbf{A}(i)) \mid i \in I \rangle$ is the non-unital \mathbb{C} -linear *-subcategory of $\text{Lin}_{\mathbb{C}}(\text{colim}_I \mathcal{F}_{\mathbb{C}}(\mathbf{A}))$ generated as an ideal by the images of $R(\mathbf{A}(i))$ for all i in I . By construction for every i in I we have a canonical factorization

$$\begin{array}{ccc} \text{Lin}_{\mathbb{C}}(\mathcal{F}_{\mathbb{C}}(\mathbf{A}(i))) & \longrightarrow & \text{Lin}_{\mathbb{C}}(\text{colim}_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})) \\ \downarrow & & \downarrow \\ \mathbf{A}(i) & \cdots\cdots\cdots & \mathbf{B} \end{array}$$

In this way we get a morphism of diagrams $\mathbf{A} \rightarrow \text{const}(\mathbf{B})$. We now observe that this exhibits \mathbf{B} as the colimit $\text{colim}_I \mathbf{A}$ in ${}^*\mathbb{C}\mathbf{Cat}_1$. Indeed, let \mathbf{C} be in ${}^*\mathbb{C}\mathbf{Cat}_1$, and a morphism $\mathbf{A} \rightarrow \text{const}(\mathbf{C})$ of diagrams in ${}^*\mathbb{C}\mathbf{Cat}_1$ be given. Then we get a morphism

$$\text{Lin}_{\mathbb{C}}(\mathcal{F}_{\mathbb{C}}(\mathbf{A})) \rightarrow \text{const}(\mathbf{C})$$

in ${}^*\mathbb{C}\mathbf{Cat}$ from the counit of the adjunction (8). Hence we get a uniquely determined morphism

$$\text{Lin}_{\mathbb{C}}(\text{colim}_I \mathcal{F}_{\mathbb{C}}(\mathbf{A})) \rightarrow \mathbf{C}$$

since colimits in ${}^*\mathbf{Cat}_1$ exist and the left adjoint $\text{Lin}_{\mathbb{C}}$ preserves colimits. We finally see that this morphism, by definition of \mathbf{B} , uniquely factorizes over a morphism $\mathbf{B} \rightarrow \mathbf{C}$.

In order to deal with the marked case we use the marked version of the adjunction (8) and argue in a similar manner using that we have shown above that ${}^*\mathbf{Cat}_1^+$ is cocomplete.

We now consider colimits in $C_{\text{pre}}^* \mathbf{Cat}_1$. The inclusion functor \mathcal{F}_{pre} is the left-adjoint of the adjunction (12). Consequently it preserves colimits. It therefore suffices to show

that if $\mathbf{A} : I \rightarrow C^*_{\text{pre}} \mathbf{Cat}_1$ is a diagram of pre- C^* -categories, then the \mathbb{C} -linear $*$ -category $\text{colim}_I \mathcal{F}_{\text{pre}}(\mathbf{A})$ is in fact a pre- C^* -category. Let f be a morphism in $\text{colim}_I \mathcal{F}_{\text{pre}}(\mathbf{A})$. We must show that $\|f\|_{\text{max}}$ is finite. By the description of colimits in \mathbf{Cat}_1 given in Remark 8.3, and by the construction of colimits in $^* \mathbf{Cat}_1, {}^*_{\mathbb{C}} \mathbf{Cat}_1$ given above, the morphism f is equal to a finite linear combinations of finite compositions morphisms of the form $u_i(f_i)$, where f_i belongs to $\mathcal{F}_{\text{pre}}(\mathbf{A}(i))$ and $u_i : \mathcal{F}_{\text{pre}}(\mathbf{A}(i)) \rightarrow \text{colim}_I \mathcal{F}_{\text{pre}}(\mathbf{A})$ is the canonical morphism. Hence we can assume that $f = u_i(f_i)$. If ρ is a representation of $\text{colim}_I \mathcal{F}_{\text{pre}}(\mathbf{A})$ into a C^* -algebra B , then $\rho \circ u_i$ is a representation of $\mathcal{F}_{\text{pre}}(\mathbf{A}(i))$. It follows that $\|\rho(f)\|_B \leq \|\rho(u_i(f_i))\|_B \leq \|f(i)\|_{\text{max}}$.

The same argument applies in the marked case.

If $\mathbf{A} : I \rightarrow C^* \mathbf{Cat}_1^{(+)}$ is a diagram of (marked) C^* -categories, then we have an isomorphism

$$\text{colim}_I \mathbf{A} \cong \text{Compl}(\text{colim}_I \mathcal{F}_-(\mathbf{A})) , \tag{30}$$

where the colimit on the right-hand side is interpreted in $C^*_{\text{pre}} \mathbf{Cat}_1^{(+)}$ and exists as seen above. The isomorphism (30) follows immediately from the adjunction (10) since the forgetful functor \mathcal{F}_- is fully faithful. ■

The remainder of this section is devoted to Proposition 8.5 showing that some of the $*$ -categories considered in the present paper are locally presentable.

The following discussion serves as a preparation. Let \mathcal{C} be some category. Recall from [AR94, Sec. 0.6] that a generator of \mathcal{C} is a set of objects \mathcal{G} of \mathcal{C} such that for every two distinct morphisms $f, g : C \rightarrow D$ in \mathcal{C} there exists a morphism $h : G \rightarrow C$ for some G in \mathcal{G} such that $f \circ h \neq g \circ h$. The generator is strong if in addition for every object C of \mathcal{C} and proper subobject D of C there exists a morphism $h : G \rightarrow C$ for some G in \mathcal{G} which does not factor over D .

Let \mathcal{G} be a subset of objects of \mathcal{C} .

8.4. LEMMA. *If every object of \mathcal{C} is isomorphic to a colimit of a diagram in \mathcal{C} with values in \mathcal{G} , then \mathcal{G} is a strong generator of \mathcal{C} .*

PROOF. Let $f, g : C \rightarrow D$ be two distinct morphisms in \mathcal{C} . Let $B : I \rightarrow \mathcal{C}$ be a diagram with values in \mathcal{G} such that $C \cong \text{colim}_I B$. Then we have a bijection of sets

$$\varinjlim_{\text{Iop}} \text{Hom}_{\mathcal{C}}(B, D) \cong \text{Hom}_{\mathcal{C}}(C, D) .$$

Because of $f \neq g$ there exists i in I such that $f \circ h(i) \neq g \circ h(i)$, where $h(i) : B(i) \rightarrow \text{colim}_I B \cong C$ is the canonical map. Note that $B(i)$ belongs to \mathcal{G} by assumption.

Let now $\iota : D \rightarrow C$ be the inclusion of a proper subobject. We again consider a diagram $B : I \rightarrow \mathcal{C}$ with values in \mathcal{G} such that $C \cong \text{colim}_I B$.

We argue by contradiction and assume that every morphism $G \rightarrow C$ with G in \mathcal{G} factors over ι . Since D is a subobject this factorization is unique. The canonical morphism of I -diagrams $B \rightarrow \underline{C}$ therefore provides a morphism of I -diagrams $B \rightarrow \underline{D}$ and hence

a morphism $\pi : C \cong \operatorname{colim}_I B \rightarrow D$ by the universal property of the colimit. Since $B \rightarrow \underline{D} \xrightarrow{\iota} \underline{C}$ is the canonical morphism of I -diagrams for the presentation of C we conclude that $\iota \circ \pi = \operatorname{id}_C$. We now argue that also $\pi \circ \iota = \operatorname{id}_D$ and hence ι is an isomorphism. This is in conflict with the assumption that ι is the inclusion of a proper subobject, and hence we get the desired contradiction. We know that

$$\iota \circ (\pi \circ \iota) = (\iota \circ \pi) \circ \iota = \operatorname{id}_C \circ \iota = \iota .$$

Since also $\iota \circ \operatorname{id}_D = \iota$ and ι is a monomorphism, we conclude that $\operatorname{id}_D = \pi \circ \iota$ as required. ■

If \mathcal{G} satisfies the assumption of Lemma 8.4, then we say that \mathcal{G} strongly generates \mathcal{C} .

8.5. PROPOSITION. *If \mathcal{C} belongs to $\{*\mathbf{Cat}_1, {}^*\mathcal{C}\mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathcal{C}\mathbf{Cat}_1^+\}$, then the category \mathcal{C} is locally presentable.*

PROOF. The categories in question are cocomplete. By [AR94, Thm. 1.20] it suffices to show that they have a strong generator formed by κ -presentable objects for some regular cardinal κ . In order to exhibit a strong generator will use the criterion shown in Lemma 8.4.

We start with the case $*\mathbf{Cat}_1$. The following discussion is related with Remark 8.3. A directed $*$ -graph is a directed graph with an involution which preserves vertices and flips the direction of edges. We consider the category $*\mathbf{DirGraph}$ of directed $*$ -graphs and involution-preserving morphisms. Then we have an adjunction

$$\operatorname{Free}_{*Cat} : *\mathbf{DirGraph} \rightleftarrows *\mathbf{Cat}_1 : \mathcal{F}_\circ , \tag{31}$$

where \mathcal{F}_\circ forgets the category structure and retains the $*$ -operation. The left-adjoint $\operatorname{Free}_{*Cat}$ sends a directed $*$ -graph to the $*$ -category freely generated by it. The category $*\mathbf{DirGraph}$ is locally presentable. Indeed, it is cocomplete (as in the case of directed graphs, colimits are given by the colimits of the sets of vertices and edges, separately), and it is strongly generated by the objects in the list

$$\{pt, \mathcal{F}_\circ(\mathbf{1})\}$$

of compact directed $*$ -graphs. Note that pt is the directed $*$ -graph with one vertex and no edges, and the directed $*$ -graph $\mathcal{F}_\circ(\mathbf{1})$ has two vertices 0 and 1 and the edges $u : 0 \rightarrow 1$ and $u^* : 1 \rightarrow 0$. Given a $*$ -category \mathbf{A} we can consider the free $*$ -category

$$\mathbf{F}(A) := \operatorname{Free}_{*Cat}(\mathcal{F}_\circ(\mathbf{A}))$$

generated by the underlying directed $*$ -graph of \mathbf{A} . The counit of the adjunction (31) provides a canonical morphism

$$v_{\mathbf{A}} : \mathbf{F}(\mathbf{A}) \rightarrow \mathbf{A}$$

of $*$ -categories. We claim that $v_{\mathbf{A}}$ is an effective epimorphism, i.e, canonical coequalizer map

$$cv_{\mathbf{A}} : \operatorname{Coeq}(\mathbf{F}(\mathbf{A}) \times_{\mathbf{A}} \mathbf{F}(\mathbf{A}) \rightrightarrows \mathbf{F}(\mathbf{A})) \rightarrow \mathbf{A}$$

is an isomorphism in ${}^*\mathbf{Cat}_1$. We first observe that $v_{\mathbf{A}}$ induces a bijection on the level of objects. Consequently, the coequalizer map $cv_{\mathbf{A}}$ is a bijection on the level of objects, too. The morphisms of the coequalizer are given as a quotient of the morphisms in $\mathbf{F}(\mathbf{A})$ by the equivalence relation induced by $v_{\mathbf{A}}$ which is compatible with the $*$ -category structure. It is now clear that $cv_{\mathbf{A}}$ is also a bijection on morphisms.

We know that $\mathbf{F}(\mathbf{A})$ is isomorphic to a colimit of a diagram involving the generators

$$\{pt, \mathbf{F}(\mathbf{1})\} . \tag{32}$$

Note that the fibre product over \mathbf{A} is not a colimit. But we have a surjection

$$\mathbf{F}(\mathbf{F}(\mathbf{A}) \times_{\mathbf{A}} \mathbf{F}(\mathbf{A})) \rightarrow \mathbf{F}(\mathbf{A}) \times_{\mathbf{A}} \mathbf{F}(\mathbf{A})$$

and therefore an isomorphism

$$\text{Coeq}\left(\mathbf{F}(\mathbf{F}(\mathbf{A}) \times_{\mathbf{A}} \mathbf{F}(\mathbf{A})) \rightrightarrows \mathbf{F}(\mathbf{A})\right) \rightarrow \mathbf{A} .$$

The $*$ -category $\mathbf{F}(\mathbf{F}(\mathbf{A}) \times_{\mathbf{A}} \mathbf{F}(\mathbf{A}))$ is again a colimit of a diagram involving the generators in the list above. Hence \mathbf{A} itself is a colimit of a diagram built from the list (32). Since $\mathbf{F}(\mathbb{I})$ has two objects and countable morphism sets it is \aleph_1 -presentable. It follows that ${}^*\mathbf{Cat}_1$ is strongly generated by the list (32) of \aleph_1 -presentable objects.

In the case of ${}^*\mathbf{Cat}_1^+$ we argue similarly. We use the adjunction

$$\text{Free}_{*Cat^+} : {}^*\mathbf{DirGraph}^+ \rightleftarrows {}^*\mathbf{Cat}_1^+ : \mathcal{F}_\circ \tag{33}$$

and that the category of marked directed $*$ -graphs ${}^*\mathbf{DirGraph}^+$ is strongly generated by the list compact objects

$$\{pt, \mathcal{F}_\circ(\text{mi}(\mathbf{1}_{*Cat_1})), \mathcal{F}_\circ(\text{ma}(\mathbf{1}_{*Cat_1}))\} .$$

We can now repeat the argument. It follows that ${}^*\mathbf{Cat}_1^+$ is strongly generated by the list of \aleph_1 -presentable objects

$$\{pt, \mathbf{F}(\text{ma}(\mathbf{1}_{*Cat_1})), \mathbf{F}(\text{mi}(\mathbf{1}_{*Cat_1}))\} .$$

A similar argument applies in the \mathbb{C} -linear case. Here we use the adjunction (8) and the counit

$$v_{\mathbf{A}} : \mathbf{F}_{\mathbb{C}}(\mathbf{A}) := \text{Lin}_{\mathbb{C}}(\mathcal{F}_{\mathbb{C}}(\mathbf{A})) \rightarrow \mathbf{A} .$$

We again show that $v_{\mathbf{A}}$ is an effective coequalizer and get an isomorphism

$$\text{Coeq}\left(\mathbf{F}_{\mathbb{C}}(\mathbf{F}_{\mathbb{C}}(\mathbf{A}) \times_{\mathbf{A}} \mathbf{F}_{\mathbb{C}}(\mathbf{A})) \rightrightarrows \mathbf{F}_{\mathbb{C}}(\mathbf{A})\right) \rightarrow \mathbf{A} .$$

From the already verified case ${}^*\mathbf{Cat}_1$ and the fact that the left-adjoint $\text{Lin}_{\mathbb{C}}$ preserves colimits we conclude that for every \mathbf{B} in ${}^*\mathbb{C}\mathbf{Cat}_1$ the object $\mathbf{F}_{\mathbb{C}}(\mathbf{B})$ is a colimit of a diagram with values in the list

$$\{\text{Lin}_{\mathbb{C}}(pt), \text{Lin}_{\mathbb{C}}(\mathbf{F}(\mathbf{1}_{*Cat_1}))\} . \tag{34}$$

Since $\text{Lin}_{\mathbb{C}}(\mathbf{F}(\mathbf{1}_{*\mathbf{Cat}_1}))$ has two objects and countable-dimensional morphism spaces it is \aleph_1 -presentable. It follows that $*_{\mathbb{C}}\mathbf{Cat}_1$ is strongly generated by the list (34) of \aleph_1 -presentable objects.

Similarly, $*_{\mathbb{C}}\mathbf{Cat}_1^+$ is strongly generated by the list of \aleph_1 -presentable objects

$$\{\text{Lin}_{\mathbb{C}}(*) , \text{Lin}_{\mathbb{C}}(\mathbf{F}(\text{mi}(\mathbf{1}_{*\mathbf{Cat}_1}))) , \text{Lin}_{\mathbb{C}}(\mathbf{F}(\text{ma}(\mathbf{1}_{*\mathbf{Cat}_1})))\} .$$

■

8.6. REMARK. At the moment we do not have an argument that $C_{\text{pre}}^*\mathbf{Cat}_1$ or $C_{\text{pre}}^*\mathbf{Cat}_1^+$ are locally presentable.

Let \mathbf{A} be a pre- C^* -category. Then by the above there exists a functor $S : I \rightarrow *_{\mathbb{C}}\mathbf{Cat}_1$ such that $S(i)$ belongs to the list (34) for all i in I together with a transformation $u : S \rightarrow \text{const}(\mathcal{F}_{\text{pre}}(\mathbf{A}))$ such that its adjoint is an isomorphism $\text{colim}_I S \cong \mathcal{F}_{\text{pre}}(\mathbf{A})$. Then

$$\text{colim}_I \text{Bd}^{\infty}(S) \rightarrow \mathbf{A}$$

is a candidate for a presentation of \mathbf{A} .

8.7. PROPOSITION. *The categories $C^*\mathbf{Cat}_1$ and $C^*\mathbf{Cat}_1^+$ are locally presentable.*

PROOF. There is a set $*\mathbf{DirGraph}^{\text{fin}}$ of finite directed $*$ -graphs. For any directed $*$ -graph Q we can consider the \mathbb{C} -linear $*$ -category $\text{Lin}_{\mathbb{C}}(\text{Free}_{*\mathbf{Cat}}(Q))$ (see (31) for notation). We have a set of C^* -norms $N(Q)$ on $\text{Lin}_{\mathbb{C}}(\text{Free}_{*\mathbf{Cat}}(Q))$. For every norm $\| - \|$ in $N(Q)$ we form the C^* -category $\mathbf{A}(Q, \| - \|)$ by taking the closure.

We claim that the set of C^* -categories $\mathbf{A}(Q, \| - \|)$ as described above for all finite directed $*$ -graphs Q and norms $\| - \|$ in $N(Q)$ strongly generates $C^*\mathbf{Cat}_1$.

Let \mathbf{A} be a C^* -category. Any finite $*$ -invariant collection F of morphisms of \mathbf{A} defines a finite directed $*$ -graph $Q(F)$ by forgetting the composition. By the universal property of the $\text{Lin}_{\mathbb{C}} \circ \text{Free}_{*\mathbf{Cat}}$ -functor we have a canonical morphism

$$\text{Lin}_{\mathbb{C}}(\text{Free}_{*\mathbf{Cat}}(Q(F))) \rightarrow \mathbf{A}$$

which induces a norm $\| - \|_F$ in $N(Q(F))$. Then $\mathbf{A}(F) := \mathbf{A}(Q(F), \| - \|_F)$ is naturally isomorphic to a sub C^* -category of \mathbf{A} . The set of finite $*$ -invariant subsets of morphisms of \mathbf{A} is partially ordered by inclusion and filtered. If F is contained in a larger subset F' , then we clearly get a monomorphism $\mathbf{A}(F) \rightarrow \mathbf{A}(F')$ of C^* -categories. We claim that

$$\text{colim}_F \mathbf{A}(F) \cong \mathbf{A} .$$

To this end we verify the universal property of the colimit. Let \mathbf{B} be a C^* -category. We must produce a natural bijection

$$\text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}, \mathbf{B}) \cong \lim_F \text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}(F), \mathbf{B}) .$$

This bijection identifies a morphism Φ in $\text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}, \mathbf{B})$ with the morphism Ψ in $\lim_F \text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}(F), \mathbf{B})$. Given Φ we can find the system $\Psi = (\Psi_F)_F$ by $\Psi_F := \Phi|_{\mathbf{A}(F)}$. Vice versa, given $\Psi = (\Psi_F)_F$, then we define Φ as follows. Let a be an object of \mathbf{A} . Then we define $\Phi(a) := \Psi_{\{\text{id}_a\}}(a)$. For a morphism $(f : a \rightarrow a')$ in \mathbf{A} we consider any F such that $f \in F$. Then we define $\Phi(f) := \Psi_F(f)$. Note that $\Psi_F(f)$ is really a morphism from $\Phi(a)$ to $\Phi(a')$. Furthermore one checks that this definition is independent of the choice of F . Therefore we get maps Φ on the level of objects and morphisms. We now show that Φ is a morphism between \mathbb{C} -linear $*$ -categories. We discuss the compatibility with composition. Let f and g be composable morphisms. Then we choose F such that $f, g, g \circ f \in F$ and use that Ψ_F is a morphism of C^* -categories.

We now claim that the C^* -categories $\mathbf{A}(Q, \| - \|)$ for finite directed $*$ -graphs and norms $\| - \|$ in $N(Q)$ are \aleph_2 -compact. Let

$$B : I \rightarrow C^*\mathbf{Cat}_1, \quad i \mapsto B(i)$$

be an \aleph_2 -filtered diagram of C^* -categories and consider the natural map

$$V : \text{colim}_{i \in I} \text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}(Q, \| - \|), B(i)) \rightarrow \text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}(Q, \| - \|), \text{colim}_{i \in I} B(i)). \quad (35)$$

We must show that V is a bijection. We first discuss the surjectivity of V . Let Ψ belong to

$$\text{Hom}_{C^*\mathbf{Cat}_1}(\mathbf{A}(Q, \| - \|), \text{colim}_{i \in I} B(i)).$$

Since the directed $*$ -graph Q is finite there exists an element i in I such that $\Psi(Q) \subseteq B(i)$. For every j in $I_{\geq i}$ we get a morphisms of \mathbb{C} -linear $*$ -categories

$$\rho_j : \text{Lin}_{\mathbb{C}}(\text{Free}_{*Cat}(Q)) \rightarrow B(i) \rightarrow B(j).$$

This morphism induces a norm $\| - \|_j$ on $\text{Lin}_{\mathbb{C}}(\text{Free}_{*Cat}(Q))$ by $\|a\|_j := \|\rho_j(a)\|_{B(j)}$. If we can show that $\| - \|_k \leq \| - \|$ for some k in $I_{\geq i}$, then we get the desired factorization $\Phi : \mathbf{A}(Q, \| - \|) \rightarrow B(k)$ such that $V(\Phi) = \Psi$.

In order to find the element k we consider the set $N(Q)$ of norms on $\text{Lin}_{\mathbb{C}}(\text{Free}_{*Cat}(Q))$ as a partially ordered set. Then we have an order-preserving map

$$\ell : I_{\geq i} \rightarrow N(Q)^{\text{op}}, \quad j \mapsto \| - \|_j.$$

We now observe that the size of $N(Q)$ is bounded by \aleph_1 . A norm on $\text{Lin}_{\mathbb{C}}(\text{Free}_{*Cat}(Q))$ is determined by its restriction to the subcategory $\text{Lin}_{\mathbb{Q}}(\text{Free}_{*Cat}(Q))$. We then use that $\text{Lin}_{\mathbb{Q}}(\text{Free}_{*Cat}(Q))$ has countably many morphisms.

We let J be a subset of objects of $I_{\geq i}$ obtained by choosing a preimage under ℓ for every norm in the image $\ell(I_{\geq i})$. Then the size of J is bounded by \aleph_1 . Since I is \aleph_2 -filtered the subset J has an upper bound k in $I_{\geq i}$. Then by construction $\| - \|_k \leq \| - \|_j$ for all j in $I_{\geq i}$.

Since for every morphism a in $\text{Lin}_{\mathbb{C}}(\text{Free}_{*Cat}(Q))$ we have the inequality

$$\|a\|_k \leq \lim_{j \in I_{\geq i}} \|\rho_j(a)\|_{B(j)} = \|\Psi(a)\|_{\text{colim}_{j \in I} B(j)} \leq \|a\|$$

we have $\| - \|_k \leq \| - \|$ as desired.

We now consider injectivity of V in (35). Assume that Φ, Φ' in

$$\text{colim}_{i \in I} \text{Hom}_{C^*Cat_1}(\mathbf{A}(Q, \| - \|), B(i))$$

are such that $V(\Phi) = V(\Phi')$. We can assume that there is an element j in I such that Φ and Φ' are represented by morphisms Φ_j, Φ'_j in $\text{Hom}_{C^*Cat_1}(\mathbf{A}(Q, \| - \|), B(j))$. For every i in $I_{\geq j}$ we write Φ_i and Φ'_i for the morphisms obtained from Φ_j and Φ'_j by post-composition with $B(j) \rightarrow B(i)$. We must then show that there exists k in $I_{\geq j}$ such that $\Phi_k = \Phi'_k$.

Using that Q is finite, after increasing j if necessary, we can assume that Φ_j and Φ'_j coincide on objects. We furthermore write V_i for the composition of V with the canonical map

$$\text{Hom}_{C^*Cat_1}(\mathbf{A}(Q, \| - \|), B(i)) \rightarrow \text{colim}_{l \in I} \text{Hom}_{C^*Cat_1}(\mathbf{A}(Q, \| - \|), B(l)) .$$

For a morphism ϕ in $\mathbf{A}(Q, \| - \|)$ we have the equality

$$0 = \|V(\Phi(\phi)) - V(\Phi'(\phi))\| = \lim_{i \in I_{\geq j}} \|\Phi_i(\phi) - \Phi'_i(\phi)\| \tag{36}$$

(note that the difference makes sense since Φ_i and Φ'_i coincide on objects).

We now use that, by continuity, Φ_i and Φ'_i are uniquely determined by their restrictions along the functor

$$d : \text{Lin}_{\mathbb{Q}}(\text{Free}_{*Cat}(Q)) \rightarrow \mathbf{A}(Q, \| - \|) .$$

Because of (36), for every morphism ϕ in $\text{Lin}_{\mathbb{Q}}(\text{Free}_{*Cat}(Q))$ and positive real number r we can choose $i(\phi, r)$ in $I_{\geq j}$ such that

$$\|\Phi_{i(\phi, r)}(d(\phi)) - \Phi'_{i(\phi, r)}(d(\phi))\| \leq r .$$

Since the size of the set of morphisms ϕ and positive real numbers r is bounded by \aleph_1 and I is \aleph_2 -filtered there exists an element k in $I_{\geq j}$ which is greater than all the elements $i(\phi, r)$ chosen above. We conclude that $\|\Phi_k(d(\phi)) - \Phi'_k(d(\phi))\| = 0$ for all morphisms ϕ in $\text{Lin}_{\mathbb{Q}}(\text{Free}_{*Cat}(Q))$ and consequently $\Phi_k = \Phi'_k$. This finally implies that $\Phi = \Phi'$.

This finishes the proof of the Proposition 8.7 in the case of C^*Cat_1 . In the case of $C^*Cat_1^+$ we argue similarly with marked directed *-graphs and use the functor Free_{*Cat+} from (33) instead of Free_{*Cat} . ■

9. Verification of the model category axioms

Let \mathcal{C} be a member of the list

$$\{*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}. \quad (37)$$

In this section we state the main theorem on the model category structures again. We first recall the description of cofibrations, fibrations and weak equivalences.

9.1. DEFINITION.

1. A weak equivalence in \mathcal{C} is a (marked) unitary equivalence (see Definition 5.2).
2. A cofibration is a morphism in \mathcal{C} which is injective on objects.
3. A fibration is a morphism in \mathcal{C} which has the right-lifting property with respect to trivial cofibrations.

In condition 1 and below the word *marked* only applies to the four marked versions. In the marked case, by Lemma 5.3.1 a weak equivalence detects marked morphisms.

For the simplicial structure we refer to Definition 6.19.

9.2. THEOREM. *The structures described in Definition 9.1 and Definition 6.19 equip \mathcal{C} with a simplicial model category structure.*

If \mathcal{C} is a member of $\{\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, {}^*\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}$, then this model category structure is cofibrantly generated and combinatorial.*

9.3. REMARK. In the case of $\mathcal{C} = C^*\mathbf{Cat}_1$ a proof of the first part of the theorem has been given in [Del10].

9.4. REMARK. It is a lack of suitable morphism classifier objects in the pre- C^* -category cases, which prevents us to show cofibrant generation in these cases, see also Remark 4.7.

In the present section we show that the structures explained above determine a model category structure on \mathcal{C} . The simplicial axioms will be verified in Section 10. Finally, the additional assertions on cofibrant generation and combinatoriality are shown in Section 11

In the following we list the axioms (cf. [Hov99]) which we have to verify in order to show that \mathcal{C} with the structures given in Definition 9.1 is a model category:

1. ((co)completeness) Completeness and cocompleteness have been verified in Section 8.
2. (retracts) This is Proposition 9.12.
3. (2 out of 3) This is Lemma 9.11.
4. (lifting) This is Proposition 9.8 together with Corollary 9.9 and Proposition 9.10.
5. (factorization) This is shown in Proposition 9.13.

In Definition 9.1 we have characterized fibrations by the lifting property. In the following we explicitly define a set of morphisms called good morphisms for the moment. Later in Proposition 9.8 it will turn out that these are exactly the fibrations.

We consider a morphism $a : \mathbf{C} \rightarrow \mathbf{D}$ in a category \mathcal{C} belonging to the list (37).

9.5. DEFINITION. *The morphism a is called good⁶, if for every object d of \mathbf{D} and unitary (marked) morphism $u : a(c) \rightarrow d$ for some object c of \mathbf{C} there exists a unitary (marked) morphism $v : c \rightarrow c'$ such that $a(v) = u$.*

Here the word *marked* only applies in the marked cases.

Let Δ^0 in \mathcal{C} be the object classifier (Definition 4.2) and \mathbb{I} be the classifier of invertibles in \mathbf{Cat}_1 (Definition 4.14). Note that \mathbb{I} is a groupoid.

9.6. REMARK. In the unmarked case we have an isomorphism $\mathbf{1} \cong \Delta^0 \sharp \mathbb{I}$, where $\mathbf{1}$ is the unitary morphism classifier (Definition 4.10).

In the marked case we have $\mathbf{1}^+ \cong \Delta^0 \sharp \mathbb{I}$, where $\mathbf{1}^+$ (Definition 4.12) classifies the marked morphisms.

Let $\Delta^0 \rightarrow \Delta^0 \sharp \mathbb{I}$ classify the object 0. We consider a morphism $a : \mathbf{C} \rightarrow \mathbf{D}$ as above.

9.7. LEMMA. *The morphism a is good if and only if it has the right lifting property with respect to*

$$\Delta^0 \rightarrow \Delta^0 \sharp \mathbb{I} .$$

PROOF. In view of the universal properties of Δ^0 and $\Delta^0 \sharp \mathbb{I}$ this is just a reformulation of Definition 9.5. ■

9.8. PROPOSITION. *The good morphisms in \mathcal{C} have the right lifting property with respect to trivial cofibrations.*

PROOF. We consider a diagram

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{\alpha} & \mathbf{C} \\ \downarrow i & \nearrow \ell & \downarrow f \\ \mathbf{B} & \xrightarrow{\beta} & \mathbf{D} \end{array} \tag{38}$$

where f is good and i is a trivial cofibration. We can find a morphism $j : \mathbf{B} \rightarrow \mathbf{A}$ such that $j \circ i = \text{id}_{\mathbf{A}}$ and there is a unitary (marked) equivalence $u : i \circ j \rightarrow \text{id}_{\mathbf{B}}$ which in addition satisfies $u \circ i = \text{id}_i$.

On objects we define ℓ as follows: If b is an object of \mathbf{B} such that $b = i(a)$ for some object a of \mathbf{A} , then we set $\ell(b) := \alpha(a)$. This makes the upper triangle commute. If b is not in the image of i , then we get a (marked) unitary $\beta(u_b) : f(\alpha(j(b))) = \beta(i(j(b))) \rightarrow \beta(b)$. Using that f is good we choose a (marked) unitary $v : \alpha(j(b)) \rightarrow c$ such that $f(v) = \beta(u_b)$. We then set $\ell(b) := c$. This makes the lower triangle commute. ⁷

⁶The analog of this notion in category theory is called an isofibration. So we could call these morphisms unitary or marked isofibrations, but these names are longer.

⁷Note that the argument in [Del10, Lemma 4.10] contains a mistake at this point. With the definition given there the lower triangle would not commute on the level of objects

We now define the lift ℓ on a morphism $\phi : b \rightarrow b'$. We distinguish four cases:

1. If b and b' are in the image of i , then (since i is an equivalence) there exists a unique morphism ψ in \mathbf{A} such that $i(\psi) = \phi$ and we set

$$\ell(\phi) := \alpha(\psi) .$$

This again makes the upper triangle commute.

2. If $b = i(a)$ and b' is not in the image of i , then we let v' and c' be the choices as above made for b' . In this case we set

$$\ell(\phi) = v' \circ \alpha(j(\phi)) .$$

3. Similarly, if $b' = i(a')$ and b is not in the image of i , then we set

$$\ell(\phi) := \alpha(j(\phi)) \circ v^{-1} .$$

4. Finally, if both b and b' do not belong to the image, then we set

$$\ell(\phi) := v' \circ \alpha(j(\phi)) \circ v^{-1} .$$

One can check that then the lower triangle commutes and that this really defines a functor. One further checks (using that the morphisms v, v' are (marked) unitaries) that ℓ is a morphism of (marked) $*$ -categories. Finally, if \mathcal{C} is one of the \mathbb{C} -vector space enriched cases, then ℓ is a functor between (marked) \mathbb{C} -linear $*$ -categories. ■

9.9. COROLLARY. *The sets of good morphisms and the fibrations coincide.*

PROOF. Since $\Delta^0 \rightarrow \Delta^0 \sharp \mathbb{I}$ is a trivial cofibration, by Lemma 9.7 the fibrations are contained in the good morphisms. By Proposition 9.8 every good morphism is a fibration. ■

9.10. PROPOSITION. *The cofibrations in \mathcal{C} have the left-lifting property with respect to the good morphisms which are in addition weak equivalences.*

PROOF. We again consider a diagram (38). Since the map i is injective on objects and the morphism f is surjective on objects we can find a lift ℓ on the level objects. Let now b, b' be objects in \mathbf{B} . Since f is fully faithful (see Lemma 5.3) we have a bijection

$$\mathrm{Hom}_{\mathbf{C}}(\ell(b), \ell(b')) \xrightarrow{f_* \cong} \mathrm{Hom}_{\mathbf{D}}(\beta(b), \beta(b')) .$$


We can therefore define ℓ on $\mathrm{Hom}_{\mathbf{B}}(b, b')$ by

$$\mathrm{Hom}_{\mathbf{B}}(b, b') \xrightarrow{\beta} \mathrm{Hom}_{\mathbf{D}}(\beta(b), \beta(b')) \cong \mathrm{Hom}_{\mathbf{C}}(\ell(b), \ell(b')) .$$

The lower triangle commutes by construction. One can furthermore check that the upper triangle commutes. Finally one checks that this really defines a functor. Since f detects marked morphisms the functor ℓ preserves them. One now checks that the functor ℓ is a morphism between (marked) $*$ -categories. If \mathcal{C} is one of the \mathbb{C} -vector space enriched cases, then obviously ℓ is enriched, too. ■

9.11. LEMMA. *The weak equivalences in \mathcal{C} satisfy the two-out-of-three axiom.*

PROOF. It is clear that the composition of weak equivalences is a weak equivalence. Assume that $f : \mathbf{A} \rightarrow \mathbf{B}$ and $g : \mathbf{B} \rightarrow \mathbf{C}$ are morphisms such that f and $g \circ f$ are weak equivalences. Then we must show that g is a weak equivalence. Let $m : \mathbf{B} \rightarrow \mathbf{A}$ and $n : \mathbf{C} \rightarrow \mathbf{A}$ inverse functors and $u : m \circ f \rightarrow \text{id}_{\mathbf{A}}$ and $v : f \circ m \rightarrow \text{id}_{\mathbf{B}}$ and $x : n \circ g \circ f \rightarrow \text{id}_{\mathbf{A}}$ and $y : g \circ f \circ n \rightarrow \text{id}_{\mathbf{C}}$ the corresponding unitary (marked) isomorphisms.



Then we consider the functor $h := f \circ n : \mathbf{C} \rightarrow \mathbf{B}$. We have unitary (marked) isomorphisms

$$h \circ g = f \circ n \circ g \xrightarrow{v^{-1}} f \circ n \circ g \circ f \circ m \xrightarrow{x} f \circ m \xrightarrow{v} \text{id}_{\mathbf{B}} .$$

and

$$g \circ h = g \circ f \circ n \xrightarrow{y} \text{id}_{\mathbf{C}} .$$

■

9.12. PROPOSITION. *The cofibrations, fibrations and weak equivalences are closed under retracts.*

PROOF. Since fibrations maps are characterized by a right lifting property they are closed under retracts. Cofibrations are closed under retracts since a retract diagram of categories induces a retract diagram on the level of sets of objects, and injectivity of maps between sets is closed under retracts.

We finally consider weak equivalences (compare [Del10, Lemma 4.9]). Consider a diagram

$$\begin{array}{ccccc} \mathbf{A} & \xrightarrow{i} & \mathbf{A}' & \xrightarrow{p} & \mathbf{A} \\ \downarrow f & & \downarrow f' & & \downarrow f \\ \mathbf{B} & \xrightarrow{j} & \mathbf{B}' & \xrightarrow{q} & \mathbf{B} \end{array}$$

with $p \circ i = \text{id}_{\mathbf{A}}$ and $q \circ j = \text{id}_{\mathbf{B}}$, and where f' is a weak equivalence. Let $g' : \mathbf{B}' \rightarrow \mathbf{A}'$ be an inverse of f' up to unitary (marked) isomorphism. Then $p \circ g' \circ j : \mathbf{B} \rightarrow \mathbf{A}$ is an inverse of f up to unitary (marked) isomorphism. ■

9.13. PROPOSITION. *In the category \mathcal{C} we have functorial factorizations.*

PROOF. We use a functorial cylinder object in order to factorize a morphism as

$$\text{trivial fibration} \circ \text{cofibration} .$$

We use the notation Convention 6.14.

For a morphism $a : \mathbf{A} \rightarrow \mathbf{B}$ in \mathcal{C} we define the cylinder object $Z(a)$ as the push-out

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{a} & \mathbf{B} \\ \downarrow (1) & & \downarrow \beta \\ \mathbf{A} \# \mathbb{I} & \xrightarrow{a'} & Z(a) \end{array} ,$$

where (1) is induced by the inclusion of the object 1 in \mathbb{I} . We have a morphism $\text{pr} : \mathbb{I} \rightarrow \text{pt}$. Since $a \circ \text{pr} \circ (1) = \text{id}_{\mathbf{B}} \circ a$, using the universal property of the push-out we can extend the diagram to

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{a} & \mathbf{B} \\ \downarrow (1) & & \downarrow \beta \\ \mathbf{A} \# \mathbb{I} & \xrightarrow{a'} & Z(a) \end{array} \begin{array}{c} \xrightarrow{\text{id}_{\mathbf{B}}} \\ \xrightarrow{q} \\ \xrightarrow{a \circ \text{pr}} \end{array} \mathbf{B}$$

We finally extend the diagram as follows

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{a} & \mathbf{B} \\ \downarrow (1) & & \downarrow \beta \\ \mathbf{A} \# \mathbb{I} & \xrightarrow{a'} & Z(a) \end{array} \begin{array}{c} \xrightarrow{\text{id}_{\mathbf{B}}} \\ \xrightarrow{q} \\ \xrightarrow{a \circ \text{pr}} \end{array} \mathbf{B}$$

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{a} & \mathbf{B} \\ \downarrow (1) & & \downarrow \beta \\ \mathbf{A} \# \mathbb{I} & \xrightarrow{a'} & Z(a) \end{array} \begin{array}{c} \xrightarrow{\text{id}_{\mathbf{B}}} \\ \xrightarrow{q} \\ \xrightarrow{a \circ \text{pr}} \end{array} \mathbf{B}$$

$$\begin{array}{ccc} \mathbf{A} & \xrightarrow{a} & \mathbf{B} \\ \downarrow (1) & & \downarrow \beta \\ \mathbf{A} \# \mathbb{I} & \xrightarrow{a'} & Z(a) \end{array} \begin{array}{c} \xrightarrow{\text{id}_{\mathbf{B}}} \\ \xrightarrow{q} \\ \xrightarrow{a \circ \text{pr}} \end{array} \mathbf{B}$$

using that $a \circ \text{pr} \circ (0) = a$ and setting $j := a' \circ (0)$.

We claim that j is a cofibration and q is a trivial fibration. In order to see these properties it is useful to calculate an explicit model $\tilde{Z}(a)$ for $Z(a)$.

We define $\tilde{Z}(a)$ as follows:

1. $\text{Ob}(\tilde{Z}(a)) := \text{Ob}(\mathbf{A}) \sqcup \text{Ob}(\mathbf{B})$.
2. $\text{Hom}_{\tilde{Z}(a)}(x, y) := \begin{cases} \text{Hom}_{\mathbf{A}}(x, y) & x, y \in \mathbf{A} \\ \text{Hom}_{\mathbf{B}}(a(x), y) & x \in \mathbf{A}, y \in \mathbf{B} \\ \text{Hom}_{\mathbf{B}}(x, a(y)) & x \in \mathbf{B}, y \in \mathbf{A} \\ \text{Hom}_{\mathbf{B}}(x, y) & x, y \in \mathbf{B} \end{cases}$.

3. The composition is defined in the only possible way.
4. The *-operation is induced by the *-operations on **A** and **B** in the canonical way.
5. The \mathbb{C} enrichment of **A** and **B** induces an enrichment of $\tilde{Z}(a)$.
6. In the marked cases we mark all morphisms which are marked in **A** or **B**.

We have defined $\tilde{Z}(a)$ as an object of ${}^*\mathbf{Cat}_1^{(+)}$ or ${}^*\mathbb{C}\mathbf{Cat}_1^{(+)}$ in the \mathbb{C} -enriched cases. In the case of (marked) pre- C^* -categories, if we can identify $\tilde{Z}(a)$ with $Z(a)$ as a (marked) \mathbb{C} -linear *-category, then we can conclude that it is itself a (marked) pre- C^* -category. Here we use that the inclusion of (marked) pre- C^* -categories into (marked) \mathbb{C} -linear *-categories preserves colimits.

Furthermore we see by an inspection of the definition that $\tilde{Z}(a)$ is a (marked) C^* -category if **A** and **B** are (marked) C^* -categories.


We have a canonical morphism $\mathbf{B} \rightarrow \tilde{Z}(a)$. Furthermore we have a morphism $\mathbf{A}\sharp\mathbb{I} \rightarrow \tilde{Z}(a)$ given by

1. $(x, 0) \mapsto x$
2. $(x, 1) \mapsto a(x)$

for x an object of **A**, and which is fixed on morphisms by

1. $((f, \text{id}_0) : (x, 0) \rightarrow (y, 0)) \mapsto f$
2. $(\text{id}_x, (0 \rightarrow 1)) \rightarrow \text{id}_{a(x)}$

and the compatibility with composition and the *-operation. With these definitions the square in



commutes. Let now ϕ and ψ be given as indicated. Then we define a morphism $\tilde{Z}(a) \rightarrow \mathbf{D}$ on objects by

$$x \mapsto \begin{cases} \phi(x, 0) & x \in \mathbf{A} \\ \psi(y) & y \in \mathbf{B} \end{cases},$$

and on morphisms by

$$f \mapsto \begin{cases} \phi(f, \text{id}_0) & x, y \in \mathbf{A} \\ \psi(f) & \textit{else} \end{cases}$$

In fact this morphism is uniquely determined by the commutativity of the diagram. This implies that $\tilde{Z}(a)$ is an explicit model for the push-out and hence a model for $Z(a)$.

From now on we assume that $Z(a) = \tilde{Z}(a)$. In this model the morphism $q : Z(a) \rightarrow \mathbf{B}$ is given by

1. $q(x) = \begin{cases} a(x) & x \in \mathbf{A} \\ x & x \in \mathbf{B} \end{cases}$
2. $q(f : x \rightarrow y) = \begin{cases} a(f) & x, y \in \mathbf{A} \\ f & \text{else} \end{cases}$

It is surjective on objects. In order to see that q is a weak equivalence we define an inverse $p : \mathbf{B} \rightarrow Z(a)$ by

$$p(x) := x, \quad p(f) := f.$$

where both take values in the \mathbf{B} -component. Then $q \circ p = \text{id}_{\mathbf{B}}$. Furthermore, a (marked) unitary isomorphism $u : p \circ q \rightarrow \text{id}_{Z(a)}$ is given by $u_x = \text{id}_x$ for x in \mathbf{B} and $\text{id}_{a(x)}$ for x in \mathbf{A} . It follows that q is good and a weak equivalence, hence a trivial fibration.

The morphism $j : \mathbf{A} \rightarrow Z(a)$ is the canonical embedding and clearly a cofibration. We therefore have constructed a functorial factorization

$$(a : \mathbf{A} \rightarrow \mathbf{B}) \mapsto (\mathbf{A} \xrightarrow{j} Z(a) \xrightarrow{q} \mathbf{B}).$$

We will use a functorial path object to obtain a functorial factorization of morphisms as

$$\text{fibration} \circ \text{trivial cofibration}.$$

We again use the notation Convention 6.14.

For a morphism $a : \mathbf{A} \rightarrow \mathbf{B}$ we define $P(a)$ as the pull-back


$$\begin{array}{ccc} P(a) & \xrightarrow{\alpha} & \mathbf{A} \\ \downarrow a' & & \downarrow a \\ \mathcal{F}un^?(\mathbb{I}, \mathbf{B}) & \xrightarrow{(1)^*} & \mathbf{B} \end{array}.$$

Using the universal property of the pull-back we get an extension of the diagram to

$$\begin{array}{ccc} & & \mathbf{A} \\ & \text{const} \circ a & \downarrow \\ \mathbf{A} & \xrightarrow{j} & P(a) \xrightarrow{\alpha} \mathbf{A} \\ & & \downarrow a' \quad \downarrow a \\ & & \mathcal{F}un^?(\mathbb{I}, \mathbf{B}) \xrightarrow{(1)^*} \mathbf{B} \end{array}.$$

(Note: The diagram above is a simplified representation of the complex commutative diagram in the image, showing the relationships between \mathbf{A} , $P(a)$, \mathbf{A} , $\mathcal{F}un^?(\mathbb{I}, \mathbf{B})$, and \mathbf{B} with arrows $j, \alpha, a', a, (1)^*, \text{const} \circ a, \text{id}_{\mathbf{A}}$.)

We finally extend the diagram as follows



by setting $p := (0)^* \circ a'$

The morphism j is a cofibration since it is injective on objects because of $\alpha \circ j = \text{id}_A$.

We can describe $P(a)$ explicitly as the subcategory of $\mathcal{F}un^?(\mathbb{I}, \mathbf{B}) \times \mathbf{A}$ determined on objects (ϕ, x) by the condition $\phi(1) = a(x)$ and on morphisms by (u, f) by $u(1) = a(f)$. In this picture j is given by

$$j(x) := (\text{const}(a(x)), x) , \quad j(f) = (\text{const}(a(f)), f) .$$

Note that $\alpha \circ j = \text{id}_A$ by construction. We furthermore find a (marked) unitary isomorphism $\text{id}_{P(a)} \rightarrow j \circ \alpha$ by

$$u(\phi, x) := ((0 \mapsto \phi(0 \rightarrow 1), 1 \mapsto \text{id}_{a(x)}, \text{id}_x) : (\phi, x) \rightarrow (\text{const}(a(x)), x) .$$

This shows that j is a weak equivalence. Hence j is a trivial cofibration.

It remains to show that p is a fibration. By Corollary 9.9 it suffices to show that p is good. Let a (marked) unitary morphism $u : p(\phi, x) \rightarrow b$ be given. Then we take

$$c := ((0 \mapsto b, 1 \mapsto a(x); (0 \rightarrow 1) \mapsto u^{-1}), x)$$

and define $v : (\phi, x) \rightarrow c$ by $((0 \mapsto u, 1 \mapsto \text{id}_{a(x)}, \text{id}_x)$. Then $p(v) = u$. This shows that p is good. ■

10. The simplicial axioms

We assume that the category \mathcal{C} belongs to the list

$$\{ * \mathbf{Cat}_1, {}^*_{\mathcal{C}} \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, {}^* \mathbf{Cat}_1^+, {}^*_{\mathcal{C}} \mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+ \} .$$

In this section we verify that the model category structure on \mathcal{C} described in Definition 9.1 and with the simplicial structure introduced in Definition 6.19 is a simplicial model category [Hir03, Def. 9.1.6]. Note that the axiom M6 [Hir03, Def. 9.1.6] is satisfied, in

view of the bijections (25), by the construction of the simplicial structure Definition 6.19. So it remains to verify the axiom M7 [Hir03, Def. 9.1.6]. This follows from Proposition 10.4 showing the dual version of M7 (as stated in [Hir03, Def. 9.1.6]), and the validity of M6.

We closely follow the argument given in [Del10] for C^* -categories.

10.1. LEMMA. *For \mathbf{A} in \mathcal{C} the functor*

$$\mathbf{A}\sharp- : \mathbf{sSet} \rightarrow \mathcal{C}$$

preserves (trivial) cofibrations.

PROOF. Recall that this functor is defined in Definition 6.19 as the composition

$$\mathbf{sSet} \xrightarrow{\Pi} \mathbf{Grpd}_1 \xrightarrow{\mathbf{A}\sharp-} \mathcal{C}.$$

In the following proof it is useful not to drop Π from the notation. If $i : X \rightarrow Y$ is a cofibration of simplicial sets, then i is injective on zero simplices, and hence, by the explicit description of the functor Π given below the Definition 6.18, the morphism of groupoids $\Pi(i)$ is injective on objects. This implies that $\mathbf{A}\sharp\Pi(i)$ is injective on objects.

Assume now that i is in addition weak equivalence. Then $\Pi(i)$ is an equivalence of groupoids. Let $j : \Pi(Y) \rightarrow \Pi(X)$ be an inverse equivalence and $u : j \circ \Pi(i) \rightarrow \text{id}_{\Pi(X)}$ and $v : \Pi(i) \circ j \rightarrow \text{id}_{\Pi(Y)}$ be the corresponding isomorphisms. Then we get a (marked) unitary isomorphism

$$\mathbf{A}\sharp u : (\mathbf{A}\sharp j) \circ (\mathbf{A}\sharp\Pi(i)) \rightarrow \text{id}_{\mathbf{A}\sharp\Pi(X)}$$

by $(\mathbf{A}\sharp u)_{(a,x)} := (\text{id}_a, u_x)$. Similarly, we have a (marked) unitary isomorphism

$$\mathbf{A}\sharp v : (\mathbf{A}\sharp\Pi(i)) \circ (\mathbf{A}\sharp j) \rightarrow \text{id}_{\mathbf{A}\sharp\Pi(Y)}$$

given by $(\mathbf{A}\sharp v)_{(a,x)} := (\text{id}_a, v_x)$. ■

10.2. LEMMA. *For a groupoid \mathbf{G} the functor*

$$-\sharp\mathbf{G} : \mathcal{C} \rightarrow \mathcal{C}$$

preserves (trivial) cofibrations.

PROOF. If $a : \mathbf{A} \rightarrow \mathbf{B}$ is a cofibration, then it is injective on objects. But then $a\sharp\mathbf{G}$ is injective on objects and hence a cofibration. If a is in addition a (marked) unitary equivalence, then $a\sharp\mathbf{G}$ is a (marked) unitary equivalence, too. The argument is similar to the corresponding part of the argument in the proof of Lemma 10.1. ■

We consider a commutative square

$$\begin{array}{ccc}
 \mathbf{A} & \xrightarrow{i} & \mathbf{B} \\
 f \downarrow & & \downarrow g \\
 \mathbf{C} & \xrightarrow{j} & \mathbf{D}
 \end{array} \tag{39}$$

in \mathcal{C} .

10.3. LEMMA. *If (39) is a pushout and i is a trivial cofibration, then j is a trivial cofibration.*

PROOF. Since i is a trivial cofibration, there exists a morphism $i' : \mathbf{B} \rightarrow \mathbf{A}$ such that $i' \circ i = \text{id}_{\mathbf{A}}$ and a (marked) unitary isomorphism $u : i \circ i' \rightarrow \text{id}_{\mathbf{B}}$ satisfying $u \circ i = \text{id}_i$. By the universal property of the push-out, the morphism $f \circ i' : \mathbf{B} \rightarrow \mathbf{C}$ induces a morphism $j' : \mathbf{D} \rightarrow \mathbf{C}$ such that $j' \circ j = \text{id}_{\mathbf{C}}$. In particular, j is a cofibration.

The functor $- \# \mathbb{I} : \mathcal{C} \rightarrow \mathcal{C}$ is a left-adjoint by Proposition 6.16 and therefore preserves pushouts. Moreover, $g \circ u$ provides a (marked) unitary isomorphism $j \circ f \circ i' = g \circ i \circ i' \rightarrow g$.

Using Example 6.15 we consider the (marked) unitary isomorphism $g \circ u$ between functors from \mathbf{B} to \mathbf{D} as a morphism $\mathbf{B} \# \mathbb{I} \rightarrow \mathbf{D}$. Together with the morphism $\mathbf{C} \# \mathbb{I} \rightarrow \mathbf{D}$ corresponding id_j , by the universal property of the push-out diagram (39) $\# \mathbb{I}$ we obtain an induced morphism $\mathbf{D} \# \mathbb{I} \rightarrow \mathbf{D}$ which we can interpret as a (marked) unitary isomorphism $j \circ j' \rightarrow \text{id}_{\mathbf{D}}$. This proves that j is a weak equivalence. ■

We can now verify the simplicial axiom M7.

10.4. PROPOSITION. *Let $a : \mathbf{A} \rightarrow \mathbf{B}$ be a cofibration in \mathcal{C} and $i : X \rightarrow Y$ be a cofibration in \mathbf{sSet} . Then*

$$(\mathbf{A} \# Y) \sqcup_{\mathbf{A} \# X} (\mathbf{B} \# X) \rightarrow \mathbf{B} \# Y \tag{40}$$

is a cofibration. Moreover, if i or a is in addition a weak equivalence, then (40) is a weak equivalence.

PROOF. The set objects of the push-out one left-hand side of (40) is equal to the push-out of the object sets. We write objects in $\mathbf{A} \# X$ as pairs (α, x) .

Assume that the classes of (α, y) and (β, x) in the push-out go to the same object which is then (β, y) . Then $a(\alpha) = \beta$ and $i(x) = y$. This means that $(\alpha, y) = (\text{id}, i)(\alpha, x)$ and $(\beta, x) = (a, \text{id})(\alpha, x)$. Consequently, the classes of (α, y) and (β, x) in $(\mathbf{A} \# Y) \sqcup_{\mathbf{A} \# X} (\mathbf{B} \# X)$ coincide.

Assume now that the classes of (α, y) and (α', y') go to the same object which is necessarily $(a(\alpha), y)$. Then $\alpha = \alpha'$ and $y = y'$.

Similarly, if the classes of (β, x) and (β', x') go to the same object which is necessarily $(\beta, i(x))$, then $\beta = \beta'$ and $x = x'$.

This shows that the morphism marked by ? the extended diagram

$$\begin{array}{ccc}
 \mathbf{A}\#X & \xrightarrow{\quad\quad\quad} & \mathbf{A}\#Y \\
 \downarrow c & \nearrow & \downarrow e \\
 & (\mathbf{A}\#Y) \sqcup_{\mathbf{A}\#X} (\mathbf{B}\#X) & \\
 & \nearrow ? & \searrow \\
 \mathbf{B}\#X & \xrightarrow{\quad\quad\quad} & \mathbf{B}\#Y
 \end{array}$$

is injective on objects and hence a cofibration.

Assume that a is a weak equivalence. By Lemma 10.2 the map c is a trivial cofibration. By Lemma 10.3 the morphism d is again a trivial cofibration. Since (again by Lemma 10.2) the morphism e is a trivial cofibration it follows from the two-out-of-three property for weak equivalences verified in Lemma 9.11 that the morphism $?$ is a weak equivalence.

The case that i is a weak equivalence is similar using Lemma 10.1 for the horizontal arrows. ■

11. Cofibrant generation

Let \mathcal{C} be a member of the list

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

In this section we show that the model category structure on \mathcal{C} described in Definition 9.1 is cofibrantly generated. We adapt the arguments given in [Del10, Sec. 4.1].

Recall from Section 4 that Δ^0 denotes the object classifier object in \mathcal{C} , and that the groupoid \mathbb{I} denotes the isomorphism classifier object in \mathbf{Cat} . The morphism $\Delta^0 \rightarrow \Delta^0\#\mathbb{I}$ classifying the object 0 is a trivial cofibration since it is clearly injective on objects and a (marked) unitary equivalence. So by Lemma 9.7 and Corollary 9.9 we can take

$$J := \{\Delta^0 \rightarrow \Delta^0\#\mathbb{I}\}$$

as the set of generating trivial cofibrations.

We now define the set I of generating cofibrations. We must distinguish various cases and the set I will depend on the case:

In the following we describe the details. We first assume that \mathcal{C} belongs to the list $\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1\}$. Then we consider the cofibrations U, V, W defined as follows:

1. $U : \emptyset \rightarrow \Delta^0$.
2. We let $V : \Delta^0 \sqcup \Delta^0 \rightarrow \Delta^1$ classify the pair of objects $(0, 1)$ of the morphism classifier Δ^1 , see Definition 4.4.

Table 2: generating cofibrations

case	I
$*\mathbf{Cat}_1, {}^*\mathcal{C}\mathbf{Cat}_1$	$J \cup \{U, V, V^u, W, W^u\}$
$C^*\mathbf{Cat}_1$	$J \cup \{U, V^{\text{bd}}, V^u, W^{\text{bd}}, W^u\}$
$*\mathbf{Cat}_1^+, {}^*\mathcal{C}\mathbf{Cat}_1^+$	$J \cup \{U, V, V^+, W, W^+\}$
$C^*\mathbf{Cat}_1^+$	$J \cup \{U, V^{\text{bd}}, V^+, W^{\text{bd}}, W^+\}$

3. We define P by the push-out

$$\begin{array}{ccc} \Delta^0 \sqcup \Delta^0 & \xrightarrow{V} & \Delta^1 \\ \downarrow V & & \downarrow \\ \Delta^1 & \longrightarrow & P \end{array}$$

and let $W : P \rightarrow \Delta^1$ be the map induced by id_{Δ^1} and the universal property of the push-out.

4. We let $V^u : \Delta^0 \sqcup \Delta^0 \rightarrow \mathbf{1}$ classify the pair of objects $(0, 1)$ of the unitary morphism classifier $\mathbf{1}$, see Definition 4.10.

5. We define P^u by the push-out

$$\begin{array}{ccc} \Delta^0 \sqcup \Delta^0 & \xrightarrow{V^u} & \mathbf{1} \\ \downarrow V^u & & \downarrow \\ \mathbf{1} & \longrightarrow & P^u \end{array}$$

and let $W^u : P^u \rightarrow \mathbf{1}$ be the map induced by $\text{id}_{\mathbf{1}}$ and the universal property of the push-out.

We set

$$I := J \cup \{U, V, V^u, W, W^u\} .$$

We now assume that \mathcal{C} belongs to the list $\{*\mathbf{Cat}_1^+, {}^*\mathcal{C}\mathbf{Cat}_1^+\}$. Then we consider the following cofibrations:

1. We let $V^+ : \Delta^0 \sqcup \Delta^0 \rightarrow \mathbf{1}^+$ classify the pair of objects $(0, 1)$ of the marked morphism classifier $\mathbf{1}^+$, see Definition 4.12.

2. We define P^+ by the push-out

$$\begin{array}{ccc} \Delta^0 \sqcup \Delta^0 & \xrightarrow{V^+} & \mathbf{1}^+ \\ \downarrow V^+ & & \downarrow \\ \mathbf{1}^+ & \longrightarrow & P^+ \end{array}$$

and let $W^+ : P^+ \rightarrow \mathbf{1}^+$ be the map induced by $\text{id}_{\mathbf{1}^+}$ and the universal property of the push-out.

We then set

$$I := J \cup \{U, V, V^+, W, W^+\} .$$

We now consider the case that \mathcal{C} belongs to the list $\{C^*\mathbf{Cat}_1\}$. In this case we must replace the morphism classifier by the bounded morphism classifier, see Lemma 4.3 and Definition 4.6. We consider the following cofibrations:

1. We let $V^{\text{bd}} : \Delta^0 \sqcup \Delta^0 \rightarrow \Delta^{1,\text{bd}}$ classify the pair of objects $(0, 1)$ of the bounded morphism classifier $\Delta^{1,\text{bd}}$, see Definition 4.6.
2. We define P^{bd} as the push-out

$$\begin{array}{ccc} \Delta^0 \sqcup \Delta^0 & \xrightarrow{V^{\text{bd}}} & \Delta^{1,\text{bd}} \\ \downarrow V^{\text{bd}} & & \downarrow \\ \Delta^{1,\text{bd}} & \longrightarrow & P^{\text{bd}} \end{array}$$

and let $W^{\text{bd}} : P^{\text{bd}} \rightarrow \Delta^{1,\text{bd}}$ be the map induced by $\text{id}_{\Delta^{1,\text{bd}}}$ and the universal property of the push-out.

We set

$$I := J \cup \{U, V^{\text{bd}}, V^u, W^{\text{bd}}, W^u\} .$$

Finally, in the case that \mathcal{C} belongs to the list $\{C^*\mathbf{Cat}_1^+\}$, we set

$$I := J \cup \{U, V^{\text{bd}}, V^+, W^{\text{bd}}, W^+\} .$$

Let \mathcal{C} be a member of the list

$$\{*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

11.1. LEMMA. *The trivial fibrations in \mathcal{C} are exactly the morphisms which have the right-lifting property with respect to I .*

PROOF. In all cases, by Lemma 9.7 and Corollary 9.9, a morphism f has the right-lifting property with respect to J if and only if it is a fibration. So it remains to show that the right-lifting property of f with respect to the remaining morphisms in I is equivalent to the fact that f is a weak equivalence.

We first consider the case where \mathcal{C} is in $\{*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1\}$. By Lemma 5.3.2 it suffices to show that the right-lifting property of f with respect to $\{U, V, V^u, W, W^u\}$ is equivalent to the property that $\mathcal{F}_{\text{all}}(f)$ (see (14) for notation) and $\text{ma}(f)^+$ (see (13) and Remark (2.20)) are equivalences of categories. This follows from the following observations:

1. The right-lifting property of f with respect to U is equivalent to surjectivity of f on objects.

2. The right-lifting property of f with respect to V is equivalent to fullness of $\mathcal{F}_{\text{all}}(f)$.
3. The right-lifting property of f with respect to W is equivalent to faithfulness of $\mathcal{F}_{\text{all}}(f)$.
4. The right-lifting property of f with respect to V^u is equivalent to fullness of $\text{ma}(f)^+$.
5. The right-lifting property of f with respect to W^u is equivalent to faithfulness of $\text{ma}(f)^+$.

Indeed, these conditions imply that $\mathcal{F}_{\text{all}}(f)$ and $\text{ma}(f)^+$ are equivalence of categories. For the converse, if f is a fibration and $\mathcal{F}_{\text{all}}(f)$ is an equivalence of categories, then f is necessarily surjective on objects.

We next discuss the case where \mathcal{C} is in $\{C^*\mathbf{Cat}_1\}$. By Lemma 5.3.2 it suffices to show that the right-lifting property of f with respect to $\{U, V^{\text{bd}}, V^u, W^{\text{bd}}, W^u\}$ is equivalent to the property that $\mathcal{F}_{\text{all}}(f)$ and $\text{ma}(f)^+$ are equivalences of categories. This follows from the following observations:

1. The right-lifting property of f with respect to U is equivalent to surjectivity of f on objects.
2. The right-lifting property of f with respect to V^{bd} is equivalent to the surjectivity of f on the subspaces of the morphisms of maximal norm bounded by 1. Since a linear map between pre-normed vector spaces is surjective if it is so on vectors of norm bounded by 1 this implies that f is full.
3. The right-lifting property of f with respect to W^{bd} is equivalent to the injectivity of the restriction of f to the subspace of morphisms of norm bounded by 1. This implies that f is faithful.
4. The right-lifting property of f with respect to $\{V^u, W^u\}$ is equivalent to fully faithfulness of $\text{ma}(f)^+$.

We now consider the case that \mathcal{C} is in $\{^*\mathbf{Cat}_1^+, {}^*\mathcal{C}\mathbf{Cat}_1^+\}$. In view of Lemma 5.3.1 we must show that the right-lifting property of f with respect to $\{U, V, V^+, W, W^+\}$ is equivalent to the fact that $\mathcal{F}_{\text{all}}(f)$ and f^+ are equivalences of categories. We conclude by the following observations.

1. The right-lifting property of f with respect to $\{U, V, W\}$ is equivalent to the fact that $\mathcal{F}_{\text{all}}(f)$ is an equivalence of categories which is surjective on objects.
2. The right-lifting property of f with respect to V^+ is equivalent to fullness of f^+ .
3. The right-lifting property of f with respect to W^+ is equivalent to faithfulness of f^+ .

We finally consider the case that \mathcal{C} is in $\{C^*\mathbf{Cat}_1^+\}$. Again by Lemma 5.3.1 we must show that the right-lifting property of f with respect to $\{U, V^{bf}, V^+, W^{bd}, W^+\}$ is equivalent to the fact that $\mathcal{F}_{\text{all}}(f)$ and f^+ are equivalences of categories. This follows from the following two observations already made above:

1. The right-lifting property of f with respect to $\{U, V^{bd}, W^{bd}\}$ is equivalent to the fact that $\mathcal{F}_{\text{all}}(f)$ is an equivalence of categories which is surjective on objects.
2. The right-lifting property of f with respect to $\{U, V^+, W^+\}$ is equivalent to the fact that f^+ is an equivalence of categories which is surjective on objects.

■

Let κ be a regular cardinal. A partially ordered set I is called κ -filtered if every subset of cardinality $< \kappa$ has an upper bound. A κ -filtered⁸ diagram is a diagram indexed by a κ -filtered partially ordered set. An object A in a category \mathcal{C} is called κ -compact⁹ if the functor

$$\text{Hom}_{\mathcal{C}}(A, -) : \mathcal{C} \rightarrow \mathbf{Set}$$

preserves κ -filtered colimits. The object is called small if it is κ -compact for some regular cardinal κ .

Let \mathcal{C} be a member of the list

$$\{*\mathbf{Cat}_1, {}_{\mathbb{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}_{\mathbb{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

In the following lemma the classifier objects (and the objects P, P^+ derived from them) are associated to \mathcal{C} .

11.2. LEMMA.

1. The objects $\emptyset, \Delta^0, \mathbf{1}, P^u$ and $\mathbf{1}^+, P^+$ (in the marked cases) are compact (i.e. \aleph_0 -compact).
2. The objects Δ^1 and P (if defined) are \aleph_1 -compact.
3. The objects $\Delta^{1, \text{bd}}$ and P^{bd} (if defined) are κ -compact, where κ is a regular cardinal greater than the maximum of the dimensions of the morphism spaces $\text{Hom}_{\Delta^{1, \text{bd}}}(j, k)$ for j, k in $\{0, 1\}$.

PROOF. The assertions easily follow by an inspection of the descriptions of the explicit models for these classifier categories given in Section 4. The main observation for 1. is that the respective categories have finitely many objects and finite-dimensional morphism spaces. Similarly for 2. we use that Δ^1 and P have two objects and that their morphism spaces are countable or have countable dimension in the \mathbb{C} -linear cases. Finally for 3. we use that the categories have the two objects 0, 1. ■

⁸We follow the terminology of [Lur09]. In [AR94, Def. 1.13] the word κ -directed is used.

⁹We again follow the terminology of [Lur09]. In [AR94] the term κ -presented is used. A \aleph_0 -compact object is also called finitely presented, or just compact.

11.3. COROLLARY. *The model category \mathcal{C} is cofibrantly generated by finite sets of generating cofibrations and trivial cofibrations between small objects.*

Recall that cofibrantly generated simplicial model category which is in addition locally presentable is called combinatorial. It turns out that six of our eight examples are combinatorial.

11.4. COROLLARY. *The model category structures on the categories \mathcal{C} in the list*

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}$$

are combinatorial.

PROOF. Cofibrant generation is shown in Corollary 11.3. The simplicial structure is discussed in Section 10. Finally, local presentability is shown in Proposition 8.5 for the list $\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, \}$, and in Proposition 8.7 for the list $\{C^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1^+\}$. ■

12. The construction $\mathbf{A} \mapsto \hat{\mathbf{A}}^G$

Let G be a group. The category of G -objects in a category \mathcal{C} is defined as the functor category $\mathbf{Fun}(BG, \mathcal{C})$, where BG is as in Example 2.2.

We now assume that the category \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

As explained in Section 1.10 one of the purposes of the present paper is to calculate the object $\lim_{BG} \ell_{BG}(\mathbf{A})$ in \mathcal{C}_{∞} for \mathbf{A} in $\mathbf{Fun}(BG, \mathcal{C})$, and that calculation of the limit amounts more precisely to provide an object \mathbf{B} of \mathcal{C} and equivalence $\ell(\mathbf{B}) \simeq \lim_{BG} \ell_{BG}(\mathbf{A})$. In this section we define the candidate for \mathbf{B} which will be denoted by $\hat{\mathbf{A}}^G$. We refer to Theorem 13.7 for the justification and the actual calculation of the limit. The main point of the present section is the explicit description of $\hat{\mathbf{A}}^G$ provided in Remark 12.2.

We consider a G -object \mathbf{A} in \mathcal{C} and let $\mathcal{F}un^?(\tilde{G}, \mathbf{A})$ be as in Section 7.

12.1. DEFINITION. *We define the object $\hat{\mathbf{A}}^G$ in \mathcal{C} by*

$$\hat{\mathbf{A}}^G := \lim_{BG} \mathcal{F}un^?(\tilde{G}, \mathbf{A}) . \tag{41}$$

12.2. REMARK. In this remark we derive an explicit description of $\hat{\mathbf{A}}^G$. Note that the limit in (41) is interpreted in \mathcal{C} , and that the details of this interpretation depend on the case.

We first assume that \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+\} . \tag{42}$$

In all these cases the underlying category of $\hat{\mathbf{A}}^G$ is the category of G -invariant functors in $\mathcal{F}un^?(\tilde{G}, \mathbf{A})$. Hence an object a of $\hat{\mathbf{A}}^G$ associates to every object g of the G -groupoid \tilde{G} an object $a(g)$ in \mathbf{A} . Furthermore, for every pair of objects g, h in \tilde{G} we have a (marked) unitary morphism $a(g \rightarrow h) : a(g) \rightarrow a(h)$ in \mathbf{A} . The condition that the functor a is G -invariant means that for every group element k in G we have the equalities $a(kg) = k(a(g))$ and $a(kg \rightarrow kh) = k(a(g \rightarrow h))$. Therefore, the object a of $\hat{\mathbf{A}}^G$ is completely determined by the object $a(1)$ of \mathbf{A} and a collection of (marked) unitary morphisms $\rho(g) = a(1 \rightarrow g) : a(1) \rightarrow g(a(1))$ which satisfy the cocycle condition

$$g(\rho(h)) \circ \rho(g) = \rho(hg)$$

for all elements g, h of G . We will therefore write objects of $\hat{\mathbf{A}}^G$ as pairs (b, ρ) with b in \mathbf{A} and $\rho = (\rho(g))_{g \in G}$ a cocycle as above with $\rho(g) : b \rightarrow g(b)$ for all g in G .

Again by G -invariance, a morphism $a \rightarrow a'$ between objects of $\hat{\mathbf{A}}^G$ is determined by its restriction to $a(1)$ which necessarily intertwines the cocycles, i.e., which satisfies

$$\rho'(g) \circ f(1) = g(f(1)) \circ \rho(g)$$

for all elements g of G . In other words, a morphism $f : (b, \rho) \rightarrow (b', \rho')$ is a morphism $f : b \rightarrow b'$ in \mathbf{A} such that $\rho'(g) \circ f = f \circ \rho(g)$ for all g in G . We call such a morphism an intertwiner.

Consequently, the category $\hat{\mathbf{A}}^G$ is isomorphic to the category of pairs $(b, (\rho(g))_{g \in G})$ of an object b of \mathbf{A} and a cocycle ρ , and intertwiners.

The $*$ -operation on the category $\hat{\mathbf{A}}^G$ is induced by the $*$ -operation on \mathbf{A} . If \mathbf{A} was a \mathbb{C} -linear $*$ -category, then so is $\hat{\mathbf{A}}^G$. In the marked case, marked morphisms in $\hat{\mathbf{A}}^G$ are intertwiners which are marked morphisms in \mathbf{A} . This finishes the description of $\hat{\mathbf{A}}^G$ in the case that \mathcal{C} belongs to the list (42).

In order to calculate $\hat{\mathbf{A}}^G$ in the cases where \mathcal{C} belongs to the list

$$\{C_{\text{pre}}^* \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1^+\}$$

we use the adjunction $(\mathcal{F}_{\text{pre}}, \text{Bd}^\infty)$ given in Lemma 3.8.3. We have an isomorphism

$$\lim_{BG} \cong \text{Bd}^\infty \circ \lim_{BG} \circ \mathcal{F}_{\text{pre}} \tag{43}$$

of functors from $\mathbf{Fun}(BG, \mathcal{C})$ to \mathcal{C} , where the limit on the right-hand side is interpreted (marked) \mathbb{C} -linear $*$ -categories. Consequently we get an isomorphism

$$\hat{\mathbf{A}}^G \cong \text{Bd}^\infty(\widehat{\mathcal{F}_{\text{pre}}(\mathbf{A})}^G) . \tag{44}$$

In other words, if \mathbf{A} is a (marked) pre- C^* -category, then we can calculate $\hat{\mathbf{A}}^G$ by applying the $\widehat{(-)}^G$ -construction to \mathbf{A} considered as a (marked) \mathbb{C} -linear $*$ -category, and then applying the functor Bd^∞ .

Finally we assume that \mathcal{C} belongs to the list

$$\{C^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1^+\}.$$

In this case we use that the forgetful functor (see (10))

$$\mathcal{F}_- : C^*\mathbf{Cat}_1^{(+)} \rightarrow C_{\text{pre}}^*\mathbf{Cat}_1^{(+)}$$

preserves limits and reflects isomorphisms. We thus have

$$\hat{\mathbf{A}}^G \cong \widehat{\mathcal{F}_-(\mathbf{A})}^G \stackrel{(44)}{\cong} \text{Bd}^\infty(\mathcal{F}_{\text{pre}}(\widehat{\mathcal{F}_-(\mathbf{A})})^G) \stackrel{!}{\cong} \mathcal{F}_{\text{pre}}(\widehat{\mathcal{F}_-(\mathbf{A})})^G.$$

In order to justify the isomorphism marked by ! note that the (marked) \mathbb{C} -linear *-category $\mathcal{F}_{\text{pre}}(\widehat{\mathcal{F}_-(\mathbf{A})})^G$ is again a C^* -category since the space of morphisms from the object $(b, (\rho(g))_{g \in G})$ to the object $(b', (\rho'(g))_{g \in G})$ is a closed subset of $\text{Hom}_{\mathbf{A}}(b, b')$, and the C^* -property is induced. Here we use the description of C^* -categories given in Remark 2.15.

In other words, if \mathbf{A} is a (marked) C^* -category, then we can calculate $\hat{\mathbf{A}}^G$ by applying the $\widehat{(-)}^G$ -construction to \mathbf{A} considered as a marked \mathbb{C} -linear *-category, and then noting that the result is in fact a C^* -category.

13. Infinity-categorical G -fixed points

Let \mathcal{C} be a model category and I be a small category. For every i in I we have an evaluation functor $e_i : \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}$.

The following definition describes the weak equivalences, cofibrations, and fibrations of the injective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ provided it exists.

13.1. DEFINITION.

1. A weak equivalence in $\mathbf{Fun}(I, \mathcal{C})$ is a morphism f such that $e_i(f)$ is a weak equivalence in \mathcal{C} for every i in I .
2. A cofibration in $\mathbf{Fun}(I, \mathcal{C})$ is a morphism f such that $e_i(f)$ is a cofibration in \mathcal{C} for every i in I .
3. A fibration is a morphism in $\mathbf{Fun}(I, \mathcal{C})$ which has the right-lifting property with respect to trivial cofibrations.

Let \mathcal{C} belong to the list

$$\{*\mathbf{Cat}_1, {}_{\mathbb{C}}*\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}_{\mathbb{C}}*\mathbf{Cat}_1^+, C_{\text{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}$$

and I be a small category.

13.2. THEOREM. *The injective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ exists.*

PROOF. It is a non-trivial fact that the injective model category structure on a functor category $\mathbf{Fun}(I, \mathcal{C})$ exists provided that the model category structure on the target \mathcal{C} is combinatorial. The proof involves Smith's theorem, see e.g. [Bek00, Thm. 1.7], [Lur09, Sec. A.2.6]. A textbook reference of the fact stated precisely in the form we need is [Lur09, Prop. A.2.8.2]. So in view of the second assertion of Theorem 9.2 the assertion of the Theorem follows for \mathcal{C} in the list

$$\{*\mathbf{Cat}_1, *_{\mathcal{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathcal{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}.$$

It remains to discuss the case where \mathcal{C} belongs to the list

$$\{C_{\text{pre}}^*\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1^+\}.$$

In this case we employ a result Cisinski.¹⁰

13.3. PROPOSITION. *Let \mathcal{C} be a model category and assume in addition:*

1. *\mathcal{C} is right proper.*
2. *The class of cofibrations in \mathcal{C} is closed under small limits.*
3. *If i, g are composable morphisms in \mathcal{C} and $g \circ i$ is a cofibration, then also i is a cofibration.*

Then the injective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ exists.

PROOF. The proof is a slight modification of the one of [Cis03, Thm. 6.16]. In the following we indicate the necessary modifications using the notation of this reference.

- a) In the reference model categories are only required to admit finite limits and colimits. Therefore in the statement of [Cis03, Thm. 6.16] the existence of all small limits was included as an additional assumption. In the present paper a model category is complete and cocomplete by assumption so that this requirement is satisfied automatically.
- b) The assumption of right properness is copied from [Cis03, Thm. 6.16] and employed in the same way.
- c) In [Cis03, Thm. 6.16] it is assumed that the cofibrations are exactly the monomorphisms. Monomorphisms have a categorical characterization and are preserved by right-adjoints. This is used in the proof of [Cis03, Thm. 6.16] in order to ensure that the functors denoted by τ_{A*} preserve cofibrations. These functors are right Kan-extension functors. Using the pointwise formula for the values we observe that Condition 2. above is sufficient to ensure that τ_{A*} preserves cofibrations.

¹⁰I thank D. Ch. Cisinski for explaining this fact to me.

d) The conditions 1. and 3. are used in the discussion of the last diagram in the proof of [Cis03, Thm. 6.16] with the goal to show that the arrow i is a trivial cofibration. First of all, by 1. the upper (unnamed, let it call g for our purposes) horizontal arrow in the square is a weak equivalence, since the arrow $\tau_{A*}k$ is one and $\tau_{A*}q$ is a fibration. Using this it is shown that i is a weak equivalence. So far the argument is the same as is in the reference. Since $\tau_{A*}j = g \circ i$ is a cofibration (see c)), by 3. we can conclude that i is one. ■

We now show that we can apply Proposition 13.3 for \mathcal{C} in the list $\{C_{\text{pre}}^* \mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1^+\}$. For the first Assumption 1. we use the general fact that a model category \mathcal{C} is right-proper if all its objects a fibrant [Hir03, Cor. 13.1.3]. Since every (marked) pre- C^* -category is fibrant we can conclude that $C_{\text{pre}}^* \mathbf{Cat}_1^{(+)}$ is right-proper.

For Assumption 2. note that a limit of a diagram of injective maps of sets is injective. The action of a limit of a diagram of functors on objects is the limit of the diagram of maps induced on objects. Therefore a limit of cofibrations in $C_{\text{pre}}^* \mathbf{Cat}_1^{(+)}$ is again a cofibration.

Finally, if $i \circ g$ is a composition of morphisms in $C_{\text{pre}}^* \mathbf{Cat}_1^{(+)}$ which is a cofibration, then it is injective on objects. Consequently, i is injective on objects and hence a cofibration, too. Hence $C_{\text{pre}}^* \mathbf{Cat}_1^{(+)}$ satisfies Assumption 3. ■

Let \mathcal{C} belong to the list

$$\{*\mathbf{Cat}_1, {}_{\mathbb{C}}\mathbf{Cat}_1, C_{\text{pre}}^* \mathbf{Cat}_1, C^* \mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}_{\mathbb{C}}\mathbf{Cat}_1^+, C_{\text{pre}}^* \mathbf{Cat}_1^+, C^* \mathbf{Cat}_1^+\} .$$

We have a functor $\tilde{G} \rightarrow *$ in G -categories (see Section 7.1 for notation). It induces a transformation of functors (see Convention 6.14 for notation)

$$\text{id} \rightarrow \mathcal{F}un^?(\tilde{G}, -) : \mathbf{Fun}(BG, \mathcal{C}) \rightarrow \mathbf{Fun}(BG, \mathcal{C}) . \tag{45}$$

13.4. PROPOSITION. *The functor*

$$\mathcal{F}un^?(\tilde{G}, -) : \mathbf{Fun}(BG, \mathcal{C}) \rightarrow \mathbf{Fun}(BG, \mathcal{C})$$

together with the natural transformation (45) is a fibrant replacement functor with respect to the injective model structure on $\mathbf{Fun}(BG, \mathcal{C})$.

PROOF. We use Remark 7.4 stating that $\mathcal{F}un^?(\tilde{G}, \mathbf{A}) \cong \mathcal{F}un^u(\tilde{G}, \mathbf{A})$, where on the right-hand side we consider \mathbf{A} as a (marked) (\mathbb{C} -linear) $*$ -category. In this way we avoid a case-dependent discussion.

We must first show that for every object \mathbf{A} of \mathcal{C} the transformation (45) induces a weak equivalence

$$r : \mathbf{A} \rightarrow \mathcal{F}un^?(\tilde{G}, \mathbf{A})$$

in \mathcal{C} . To this end we must find an inverse up to (marked) unitary isomorphism of r on the level of underlying objects in \mathcal{C} . We define a (non G -equivariant) functor

$$e : \mathcal{F}un^?(\tilde{G}, \mathbf{A}) \rightarrow \mathbf{A} , \quad a \mapsto a(1) , \quad e(f : a \rightarrow a') := (f(1) : a(1) \rightarrow a'(1)) .$$

Then clearly $e \circ r = \text{id}_{\mathbf{A}}$. We furthermore have a (marked) unitary isomorphism $r \circ e \rightarrow \text{id}_{\mathcal{F}un^?(\tilde{G}, \mathbf{A})}$ given on a in $\mathcal{F}un^?(\tilde{G}, \mathbf{A})$ by the collection of (marked) unitary morphisms $(a(1 \rightarrow g) : a(1) \rightarrow a(g))_{g \in G}$.

In order to finish the proof we must show that $\mathcal{F}un^?(\tilde{G}, \mathbf{A})$ is fibrant. To this end we consider a square in $\mathbf{Fun}(BG, \mathcal{C})$:

$$\begin{array}{ccc} \mathbf{C} & \longrightarrow & \mathcal{F}un^?(\tilde{G}, \mathbf{A}) \\ \downarrow c & \nearrow & \downarrow \\ \mathbf{D} & \longrightarrow & * \end{array}$$

where $\mathbf{C} \rightarrow \mathbf{D}$ is a trivial cofibration in $\mathbf{Fun}(BG, \mathcal{C})$. We must show the existence of the diagonal lift.

We use the identification $\mathcal{F}un^?(BG, *) \simeq *$ (here $*$ denotes a final object in \mathcal{C}) and the exponential law Proposition 7.5 in order to rewrite the problem as

$$\begin{array}{ccc} \mathbf{C} \# \tilde{G} & \xrightarrow{\phi} & \mathbf{A} \\ \downarrow & \nearrow & \downarrow \\ \mathbf{D} \# \tilde{G} & \longrightarrow & * \end{array}$$

\tilde{d} (curved arrow from $\mathbf{D} \# \tilde{G}$ to $\mathbf{C} \# \tilde{G}$)

Since the underlying morphism of $c : \mathbf{C} \rightarrow \mathbf{D}$ is a trivial cofibration in \mathcal{C} it is injective on objects. We choose an inverse equivalence $d : \mathbf{D} \rightarrow \mathbf{C}$ (not necessarily G -invariant) up to (marked) unitary equivalence which is a precise inverse on the image of c . We can extend the composition

$$\mathbf{D} \xrightarrow{d} \mathbf{C} \rightarrow \mathbf{C} \times \{1\} \rightarrow \mathbf{C} \# \tilde{G}$$

uniquely to a G -invariant morphism

$$\tilde{d} : \mathbf{D} \# \tilde{G} \rightarrow \mathbf{C} \# \tilde{G} .$$

Indeed, we set

$$\tilde{d}(D, g) := (g(d(g^{-1}(D))), g) , \quad \tilde{d}(f : D \rightarrow D', g \rightarrow h) := gd(g^{-1}f)\#(g \rightarrow h) .$$

The desired diagonal can now be obtained as the composition $\phi \circ \tilde{d}$. ■

13.5. REMARK. Let (\mathcal{C}, W) be a relative category. Then we can consider the localization

$$\ell : \mathcal{C} \rightarrow \mathcal{C}_\infty := \mathcal{C}[W^{-1}] \tag{46}$$

in the realm of ∞ -categories, see (2). For a small category I we let

$$\ell_I : \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathbf{Fun}(I, \mathcal{C}_\infty) \tag{47}$$

denote the functor given by post-composition with ℓ in (46).

The content of the following proposition is well-known since it provides the basic justification that, in the case of limits, ∞ -categories and model categories yields equivalent homotopical constructions. But since we do not know a reference where it is stated in this ready-to-use form we will give a proof.

Let (\mathcal{C}, W) be a relative category and I be a small category.

13.6. PROPOSITION. Assume that (\mathcal{C}, W) extends to a simplicial model category with the following properties:

1. The injective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ exists.
2. All objects of \mathcal{C} are cofibrant.

Then for any fibrant replacement functor $r : \text{id}_{\mathbf{Fun}(I, \mathcal{C})} \rightarrow R$ in the injective model category structure of $\mathbf{Fun}(I, \mathcal{C})$ we have an equivalence of functors

$$\lim_I \circ \ell_I \simeq \ell \circ \lim_I \circ R : \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}_\infty .$$

PROOF. Since (\mathcal{C}, W) extends to a simplicial model category with weak equivalences W and with all objects cofibrant we can express the mapping spaces in \mathcal{C}_∞ in terms of the simplicial mapping spaces $\text{Map}_{\mathcal{C}}$ of \mathcal{C} . More precisely, if A is a cofibrant and A' is a fibrant object of \mathcal{C} , then by Remark 1.8 we have an equivalence of spaces

$$\text{Map}_{\mathcal{C}_\infty}(\ell(A), \ell(A')) \simeq \ell_{\mathbf{sSet}}(\text{Map}_{\mathcal{C}}(A, A')) . \tag{48}$$

We let W_I denote the weak equivalences in the injective model category structure on $\mathbf{Fun}(I, \mathcal{C})$. We then have the commuting diagram

$$\begin{array}{ccc} \mathbf{Fun}(I, \mathcal{C}) & \xrightarrow{\ell_I} & \mathbf{Fun}(I, \mathcal{C}_\infty) , \\ & \searrow \alpha & \nearrow \beta \\ & & \mathbf{Fun}(I, \mathcal{C})[W_I^{-1}] \end{array} \tag{49}$$

where the arrow β is induced by the universal property of the localization functor α , see (2). It is a crucial fact shown in [Cis19, Prop. 7.9.2]¹¹ that the functor β is an equivalence.

For A in \mathcal{C} and B in $\mathbf{Fun}(I, \mathcal{C})$ we then have the following chain of natural equivalences of spaces

$$\begin{aligned} \text{Map}_{\mathcal{C}_\infty}(\ell(A), \ell(\lim_I R(B))) &\stackrel{!}{\simeq} \ell_{\mathbf{sSet}}(\text{Map}_{\mathcal{C}}(A, \lim_I R(B))) \\ &\simeq \ell_{\mathbf{sSet}}(\text{Map}_{\mathbf{Fun}(I, \mathcal{C})}(\underline{A}, R(B))) \\ &\stackrel{!!}{\simeq} \text{Map}_{\mathbf{Fun}(I, \mathcal{C})[W_I^{-1}]}(\alpha(\underline{A}), \alpha(R(B))) \\ &\stackrel{\beta, \simeq}{\xrightarrow{\quad}} \text{Map}_{\mathbf{Fun}(I, \mathcal{C}_\infty)}(\ell_I(\underline{A}), \ell_I(R(B))) \\ &\stackrel{!!!, \simeq}{\xleftarrow{\quad}} \text{Map}_{\mathbf{Fun}(I, \mathcal{C}_\infty)}(\ell_I(\underline{A}), \ell_I(B)) \\ &\simeq \text{Map}_{\mathbf{Fun}(I, \mathcal{C}_\infty)}(\ell(A), \ell(B)) \\ &\simeq \text{Map}_{\mathcal{C}_\infty}(\ell(A), \lim_I \ell(B)) . \end{aligned}$$

¹¹Alternatively, if one in addition assumes that the model category structure on \mathcal{C} is combinatorial, then one could cite [Lur09, Section 4.2.4], or better, [Lur17, Cor. 1.3.4.26] for this fact.

For the equivalence marked by ! we use Assumption 2, that $\lim_I R(B)$ is fibrant, and (48). For the equivalence marked by !! we again use (48), but now for the functor category (note that all objects of the functor category are cofibrant in the injective model category structure, and that the latter has a simplicial extension), and that by assumption $R(B)$ is fibrant in the injective model category structure on $\mathbf{Fun}(I, \mathcal{C})$. Finally, for the equivalence marked by !!! we use that $\ell_I(r) : \ell_I \rightarrow \ell_I \circ R$ is an equivalence.

The natural equivalence

$$\mathrm{Map}_{\mathcal{C}_\infty}(\ell(A), \ell(\lim_I R(B))) \simeq \mathrm{Map}_{\mathcal{C}_\infty}(\ell(A), \lim_I \ell_I(B))$$

implies the asserted equivalence of functors. ■

If we assume in addition that (\mathcal{C}, W) extends to a combinatorial model category, then Prop. 13.6 is an immediate consequence of [Lur17, 1.3.4.23].

Note that the assumption 13.6.2 that the objects of the model category extension of (\mathcal{C}, W) are cofibrant comes in since we define \mathcal{C}_∞ as the localization of the whole category \mathcal{C} by the weak equivalences W . If not all objects are cofibrant, then the correct definition of the underlying ∞ -category \mathcal{C}_∞ of the model category would be $\mathcal{C}^c[W^{-1}]$ [Lur17, 1.3.4.15].

Below we will apply this proposition in the case $I = BG$.

Let \mathcal{C} be a member of the list

$$\{*\mathbf{Cat}_1, {}^*\mathbb{C}\mathbf{Cat}_1, C_{\mathrm{pre}}^*\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}^*\mathbb{C}\mathbf{Cat}_{1+}, C_{\mathrm{pre}}^*\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\}$$

and \mathbf{A} be an object of $\mathbf{Fun}(BG, \mathcal{C})$.

13.7. THEOREM. *We have an equivalence*

$$\lim_{BG} \ell_{BG}(\mathbf{A}) \simeq \ell(\hat{\mathbf{A}}^G) .$$

PROOF. We want to apply Proposition 13.6. By Theorem 1.4 the relative category (\mathcal{C}, W) extends to a simplicial model category. By Theorem 13.2 the injective model category structure on $\mathbf{Fun}(BG, \mathcal{C})$ exists. Finally, by an inspection of the definitions (Definition 9.1 for cofibrancy and, in addition, Proposition 9.5 and Corollary 9.9 for fibrancy), all objects of \mathcal{C} are cofibrant and fibrant.

We now apply Proposition 13.6 for the explicit version of R obtained in Proposition 13.4 and Definition 12.1 in order to get the equivalences

$$\lim_{BG} \ell_{BG}(\mathbf{A}) \simeq \ell(\lim_{BG}(\mathcal{F}un^?(\tilde{G}, \mathbf{A}))) \simeq \ell(\hat{\mathbf{A}}^G) .$$

■

13.8. **REMARK.** If \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, *_{\mathbb{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathbb{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} ,$$

then by Theorem 1.4 in combination with Remark 1.6 we could base the proof of Theorem 13.7 on the version of the proof of Proposition 13.6 which only uses [Lur09, Section 4.2.4], see the footnote in the proof of 13.6.

In the remaining two cases, where \mathcal{C} belongs to the list

$$\{C_{\text{pre}}^*\mathbf{Cat}_1, C_{\text{pre}}^*\mathbf{Cat}_1^+\} ,$$

we could then deduce the assertion of Theorem 13.7 as follows.

We use that Theorem 13.7 is true in the case of $*_{\mathbb{C}}\mathbf{Cat}_1$. We let

$$\mathcal{F}_{\text{pre}} : C_{\text{pre}}^*\mathbf{Cat}_1^{(+)} \rightarrow *_{\mathbb{C}}\mathbf{Cat}_1^{(+)} , \quad \mathcal{F}_{\text{pre}} : C_{\text{pre}}^*\mathbf{Cat}^{(+)} \rightarrow *_{\mathbb{C}}\mathbf{Cat}^{(+)}$$

denote the forgetful functors on the level of 1- and of ∞ -categories. They are inclusions of full subcategories fitting into adjunctions (18). We conclude that the ∞ -category $C_{\text{pre}}^*\mathbf{Cat}^{(+)}$ is complete and the limit of an I -diagram in $C_{\text{pre}}^*\mathbf{Cat}^{(+)}$ can be calculated by the formula

$$\lim_I \simeq \text{Bd}^\infty \circ \lim_I \circ \mathcal{F}_{\text{pre}} , \tag{50}$$

where the limit on the right-hand side is taken in $*_{\mathbb{C}}\mathbf{Cat}^{(+)}$.

For \mathbf{A} in $\mathbf{Fun}(BG, \mathcal{C})$ we then have the chain of equivalences

$$\begin{aligned} \lim_{BG} \ell_{BG}(\mathbf{A}) &\stackrel{(50)}{\simeq} \text{Bd}^\infty(\lim_{BG} \mathcal{F}_{\text{pre}}(\ell_{BG}(\mathbf{A}))) \\ &\stackrel{!}{\simeq} \text{Bd}^\infty(\lim_{BG} \ell_{BG}(\mathcal{F}_{\text{pre}}(\mathbf{A}))) \\ &\stackrel{\text{Thm.13.7}}{\simeq} \text{Bd}^\infty(\ell(\widehat{\mathcal{F}_{\text{pre}}(\mathbf{A})}^G)) \\ &\stackrel{!}{\simeq} \ell(\text{Bd}^\infty(\widehat{\mathcal{F}_{\text{pre}}(\mathbf{A})}^G)) \\ &\stackrel{(44)}{\simeq} \ell(\hat{\mathbf{A}}^G) . \end{aligned}$$

At the marked morphisms we use that \mathcal{F}_{pre} and Bd^∞ descend to the ∞ -categories since they preserve weak equivalences.

14. Infinity-categorical G -orbits

Let \mathcal{C} be a model category and I be a small category. In the following definition we describe the weak equivalences, cofibrations, and fibrations of the projective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ provided it exists. Recall, that for i in I we have the evaluation functor $e_i : \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}$.

14.1. DEFINITION.

1. A weak equivalence in $\mathbf{Fun}(I, \mathcal{C})$ is a morphism f such that $e_i(f)$ is a weak equivalence in \mathcal{C} for every i in I .
2. A fibration in $\mathbf{Fun}(I, \mathcal{C})$ is a morphism f such that $e_i(f)$ is a fibration in \mathcal{C} for every i in I .
3. A cofibration is a morphism in $\mathbf{Fun}(I, \mathcal{C})$ which has the left-lifting property with respect to trivial fibrations.

It is known (see e.g. [Hir03, Thm. 11.6.1]) that the projective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ exists if the model category structure on \mathcal{C} is cofibrantly generated.

14.2. REMARK. This remark is similar to Remark 13.5. Let (\mathcal{C}, W) be relative category and I be a small category. As before we let $\ell : \mathcal{C} \rightarrow \mathcal{C}_\infty := \mathcal{C}[W^{-1}]$ be the localization and $\ell_I : \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathbf{Fun}(I, \mathcal{C}_\infty)$ be the induced functor. For an object C in $\mathbf{Fun}(I, \mathcal{C})$ we want to calculate the colimit $\operatorname{colim}_I \ell_I(C)$ in \mathcal{C}_∞ using model categorical methods. The following proposition is the analog of Proposition 13.6. Its content is well-known, but we do not have a reference where it is stated in this ready-to-use form.

14.3. PROPOSITION. *Assume that (\mathcal{C}, W) extends to a combinatorial model category. Then for any cofibrant replacement functor $l : L \rightarrow \operatorname{id}_{\mathbf{Fun}(I, \mathcal{C})}$ in the projective model category structure of $\mathbf{Fun}(I, \mathcal{C})$ we have an equivalence of functors*

$$\operatorname{colim}_I \circ \ell_I \simeq \ell \circ \operatorname{colim}_I \circ L : \mathbf{Fun}(I, \mathcal{C}) \rightarrow \mathcal{C}_\infty .$$

PROOF. We shall sketch a proof which is completely analogous to the proof of Proposition 13.6. Since a combinatorial model category structure is in particular cofibrantly generated the projective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ exists. It is again combinatorial [Lur09, Prop. A.2.8.2].

We again have the commuting diagram (49) where the arrow β is an equivalence. For a fibrant object B in \mathcal{C} and C in $\mathbf{Fun}(I, \mathcal{C})$ we then have the following chain of natural equivalences of spaces

$$\begin{aligned} \operatorname{Map}_{\mathcal{C}_\infty}(\ell(\operatorname{colim}_I L(C)), \ell(B)) &\stackrel{!}{\simeq} \ell_{\mathbf{sSet}}(\operatorname{Map}_{\mathcal{C}}(\operatorname{colim}_I L(C), B)) \\ &\simeq \ell_{\mathbf{sSet}}(\operatorname{Map}_{\mathbf{Fun}(I, \mathcal{C})}(L(C), \underline{B})) \\ &\stackrel{!!}{\simeq} \operatorname{Map}_{\mathbf{Fun}(I, \mathcal{C})[W_I^{-1}]}(\alpha(L(C)), \alpha(\underline{B})) \\ &\stackrel{\beta, \simeq}{\xrightarrow{}} \operatorname{Map}_{\mathbf{Fun}(I, \mathcal{C}_\infty)}(\ell_I(L(C)), \ell_I(\underline{B})) \\ &\stackrel{!!!, \simeq}{\longleftarrow} \operatorname{Map}_{\mathbf{Fun}(I, \mathcal{C}_\infty)}(\ell_I(C), \ell_I(\underline{B})) \\ &\simeq \operatorname{Map}_{\mathbf{Fun}(I, \mathcal{C}_\infty)}(\ell_I(C), \underline{\ell(B)}) \\ &\simeq \operatorname{Map}_{\mathcal{C}_\infty}(\operatorname{colim}_I \ell_I(C), \ell(B)) . \end{aligned}$$

with the same justifications of the equivalences as in the proof of Proposition 13.6. For the equivalences marked by ! and !! we use that the model category structures on \mathcal{C} and the functor category are combinatorial so that we still can apply Remark 1.8 in order to justify the equivalence 48 (note that in the projective model category structure on $\mathbf{Fun}(I, \mathcal{C})$ we can not expect that all objects are cofibrant), where we now have to use the existence of functorial factorizations and [Lur17, Rem. 1.3.4.16]. Note that $\text{colim}_I L(C)$ is cofibrant in \mathcal{C} .

The natural equivalence

$$\text{Map}_{\mathcal{C}_\infty}(\ell(\text{colim}_I L(C)), \ell(B)) \simeq \text{Map}_{\mathcal{C}_\infty}(\text{colim}_I \ell_I(C), \ell(B))$$

for all fibrant B implies the asserted equivalence of functors. ■

Alternatively, the assertion of Prop. 14.3 is an immediate consequence of [Lur17, 1.3.4.24].

We now assume that \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, {}_c\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, {}_c\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

14.4. REMARK. We must exclude the pre- C^* -category cases since we do not know that the corresponding model categories are combinatorial.

The relative category (\mathcal{C}, W) extends to a combinatorial model category (Theorem 1.4 and Remark 1.6) in which all objects are cofibrant and fibrant. Consequently the projective model category structure on $\mathbf{Fun}(BG, \mathcal{C})$ exists and Proposition 14.3 applies to (\mathcal{C}, W) .

Recall the Definition 7.1 of the arrow category \tilde{G} of G . Furthermore recall the Convention 6.14 concerning the usage of \sharp . We consider the functor

$$L := -\sharp\tilde{G} : \mathbf{Fun}(BG, \mathcal{C}) \rightarrow \mathbf{Fun}(BG, \mathcal{C})$$

together with the transformation $L \rightarrow \text{id}_{\mathbf{Fun}(BG, \mathcal{C})}$ induced by the morphism of G -groupoids $\tilde{G} \rightarrow *$.

14.5. LEMMA. *The functor L together with the transformation $L \rightarrow \text{id}_{\mathbf{Fun}(BG, \mathcal{C})}$ is a cofibrant replacement functor for the projective model category structure on $\mathbf{Fun}(BG, \mathcal{C})$.*

PROOF. Since $\tilde{G} \rightarrow \text{id}$ is an (non-equivariant) equivalence of groupoids and for every object \mathbf{D} in $\mathbf{Fun}(BG, \mathcal{C})$ the functor $\mathbf{D}\sharp-$ from groupoids to \mathcal{C} preserves (marked) unitary equivalences (see the proof of Lemma 10.1), the morphism $\mathbf{D}\sharp\tilde{G} \rightarrow \mathbf{D}$ is a weak equivalence. We must show that $L(\mathbf{D})$ is cofibrant. To this end we consider the lifting problem

$$\begin{array}{ccc} \emptyset & \longrightarrow & \mathbf{A} \\ \downarrow & \nearrow c & \downarrow f \\ \mathbf{D}\sharp\tilde{G} & \xrightarrow{u} & \mathbf{B} \end{array}$$

where f is a trivial fibration in \mathcal{C} . Since f is surjective on objects we can find an inverse equivalence (possibly non-equivariant) $g : \mathbf{B} \rightarrow \mathbf{A}$ for f such that $f \circ g = \text{id}_{\mathbf{B}}$. The map $\mathbf{D}\sharp\{1\} \xrightarrow{u|\mathbf{D}\sharp\{1\}} \mathbf{B} \xrightarrow{g} \mathbf{A}$ can uniquely be extended to an equivariant morphism c which is the desired lift. ■

If \mathbf{C} is an object of \mathcal{C} , then by $\underline{\mathbf{C}}$ we denote the object of $\mathbf{Fun}(BG, \mathcal{C})$ given by \mathbf{C} with the trivial action of G .

We assume that \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, *_{\mathbb{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathbb{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

Let \mathbf{C} be an object of \mathcal{C} . Note that BG is a groupoid.

14.6. THEOREM. *We have an equivalence*

$$\text{colim}_{BG} \ell_{BG}(\underline{\mathbf{C}}) \simeq \ell(\mathbf{C}\sharp BG) .$$

PROOF. By Proposition 14.3 and Lemma 14.5 we have an equivalence

$$\text{colim}_{BG} \ell_{BG}(\underline{\mathbf{C}}) \simeq \ell(\text{colim}_{BG} \underline{\mathbf{C}}\sharp \tilde{G}) . \tag{51}$$

So it remains to calculate the colimit $\text{colim}_{BG} \underline{\mathbf{C}}\sharp \tilde{G}$ in \mathcal{C} . To this end will show that the functor $\mathbf{C}\sharp - : \mathbf{Grpd}_1 \rightarrow \mathcal{C}$ commutes with colimits and calculate that $\text{colim}_{BG} \tilde{G} \cong BG$.

Let \mathbf{A} be an object of \mathcal{C} . For a second object \mathbf{D} in \mathcal{C} we let $\mathbf{Func}_{\mathcal{C}}(\mathbf{A}, \mathbf{D})_+$ denote the subgroupoid of the functor category $\mathbf{Func}_{\mathcal{C}}(\mathbf{A}, \mathbf{D})$ of all functors and (marked) unitary isomorphisms.

14.7. LEMMA. *We have an adjunction*

$$\mathbf{A}\sharp - : \mathbf{Grpd}_1 \rightleftarrows \mathcal{C} : \mathbf{Func}_{\mathcal{C}}(\mathbf{A}, -)_+ .$$

PROOF. For \mathbf{D} in \mathcal{C} and \mathbf{G} in \mathbf{Grpd}_1 we construct a natural bijection

$$\text{Hom}_{\mathcal{C}}(\mathbf{A}\sharp \mathbf{G}, \mathbf{D}) \cong \text{Hom}_{\mathbf{Grpd}_1}(\mathbf{G}, \mathbf{Func}_{\mathcal{C}}(\mathbf{A}, \mathbf{D})_+) .$$

This bijection sends a morphism Φ in $\text{Hom}_{\mathcal{C}}(\mathbf{A}\sharp \mathbf{G}, \mathbf{D})$ to Ψ in $\text{Hom}_{\mathbf{Grpd}_1}(\mathbf{G}, \mathbf{Func}_{\mathcal{C}}(\mathbf{A}, \mathbf{D})_+)$. Let Φ be given. We let g, h denote objects of \mathbf{G} and $\phi : g \rightarrow h$ a be morphism. Then we define Ψ by

$$\Psi(g)(a) := \Phi(a, g) , \quad \Psi(g)(f) := \Phi(f, \text{id}_g) , \quad \Psi(\phi) := (\Phi(\text{id}_a, \phi))_{a \in \mathbf{A}} .$$

Here a is an object of \mathbf{A} and f is a morphism in \mathbf{A} . Observe that $\Psi(\phi)$ is a unitary isomorphism since Φ is compatible with the involution and ϕ is a unitary isomorphism. In the marked case, if f is marked, then $\Psi(g)(f)$ is marked since (f, id_g) is marked in $\mathbf{A}\sharp \mathbf{G}$ and Φ preserves marked morphisms. Furthermore, $\Psi(\phi)$ is implemented by marked isomorphisms.

Vice versa, let Ψ be given. Then we define

$$\Phi(a, g) := \Psi(g)(a) , \quad \Phi(f, \phi) := \Psi(\phi)_a \circ \Psi(g)(f) .$$

This formula determines Φ on the generators of the morphisms. It can be extended by linearity (in the \mathbb{C} -linear cases) and continuity (in the C^* -cases). ■

14.8. COROLLARY. *We have an adjunction*

$$\mathbf{C}\sharp- : \mathbf{Fun}(BG, \mathbf{Grpd}_1) \rightleftarrows \mathbf{Fun}(BG, \mathcal{C}) : \mathbf{Fun}_{\mathbf{Fun}(BG, \mathcal{C})}(\underline{\mathbf{C}}, -)_+ .$$

Since $\mathbf{C}\sharp-$ is a left-adjoint functor it commutes with colimits. Consequently we have an isomorphism

$$\operatorname{colim}_{BG}(\underline{\mathbf{C}}\sharp\tilde{G}) \cong \mathbf{C}\sharp(\operatorname{colim}_{BG}\tilde{G}) . \tag{52}$$

14.9. LEMMA. *We have an isomorphism $\operatorname{colim}_{BG}\tilde{G} \cong BG$.*

PROOF. We check the universal property of the colimit. Let \mathbf{K} be any groupoid. Then we have a natural bijection

$$\operatorname{Hom}_{\mathbf{Fun}(BG, \mathbf{Grpd}_1)}(\tilde{G}, \underline{\mathbf{K}}) \cong \operatorname{Hom}_{\mathbf{Grpd}_1}(BG, \mathbf{K}) .$$

This bijection sends Φ in $\operatorname{Hom}_{\mathbf{Fun}(BG, \mathbf{Grpd}_1)}(\tilde{G}, \underline{\mathbf{K}})$ to the morphism Ψ in $\operatorname{Hom}_{\mathbf{Grpd}_1}(BG, \mathbf{K})$ given by

$$\Psi(*) := \Phi(1) , \quad \Psi(g) := \Phi(1 \rightarrow g) .$$

If Ψ is given, then we define Φ by

$$\Phi(g) := \Psi(*) , \quad \Phi(g \rightarrow h) := \Psi(g^{-1}h) .$$

■

The assertion of Theorem 14.6 now follows from a combination of the relations (51), (52), and Lemma 14.9. ■

In the following we discuss an application of Theorem 14.6 to the calculation of the values of an induction functor J^G (see Definition 14.10) from \mathcal{C} to functors from the orbit category $\mathbf{Orb}(G)$ of G to \mathcal{C}_∞ .

The objects of $\mathbf{Orb}(G)$ are the transitive G -sets, and its morphisms are equivariant maps. We can consider the underlying set of G as a transitive G -set with respect to the right action. One can then identify $\operatorname{End}_{\mathbf{Orb}(G)}(G)$ with the group G acting by left-multiplication. We therefore get a fully faithful functor

$$j : BG \rightarrow \mathbf{Orb}(G)$$

which sends the unique object of BG to the transitive G -set G , and which identifies the group $\operatorname{End}_{BG}(pt)$ (given by G) with the group $\operatorname{End}_{\mathbf{Orb}(G)}(G)$ as described above.

If \mathcal{C}_∞ is a presentable ∞ -category (or sufficiently cocomplete), then we get an adjunction

$$j_! : \mathbf{Fun}(BG, \mathcal{C}_\infty) \rightleftarrows \mathbf{Fun}(\mathbf{Orb}(G), \mathcal{C}_\infty) : j^* , \tag{53}$$

where j^* is the restriction functor along j .

We now assume that \mathcal{C} belongs to the list

$$\{*\mathbf{Cat}_1, *_{\mathbb{C}}\mathbf{Cat}_1, C^*\mathbf{Cat}_1, *\mathbf{Cat}_1^+, *_{\mathbb{C}}\mathbf{Cat}_1^+, C^*\mathbf{Cat}_1^+\} .$$

Then the corresponding ∞ -category \mathcal{C}_∞ is presentable by Corollary 1.9 so that (53) applies.

14.10. DEFINITION. We define the induction functor J^G as the composition

$$J^G : \mathcal{C} \xrightarrow{(-)} \mathbf{Fun}(BG, \mathcal{C}) \xrightarrow{\ell_{BG}} \mathbf{Fun}(BG, \mathcal{C}_\infty) \xrightarrow{j_!} \mathbf{Fun}(\mathbf{Orb}(G), \mathcal{C}_\infty) . \tag{54}$$

For a subgroup H of G we consider $H \backslash G$ with the action of G by right multiplication as an object of $\mathbf{Orb}(G)$.

14.11. PROPOSITION. We have an equivalence

$$J^G(\mathbf{C})(H \backslash G) \simeq \ell(\mathbf{C} \sharp BH) .$$

PROOF. The functor $j_!$ is a left Kan-extension functor. The point-wise formula for the left Kan-extension gives an equivalence

$$J^G(\mathbf{C})(H \backslash G) \simeq j_!(\ell_{BG}(\mathbf{C}))(H \backslash G) \simeq \operatorname{colim}_{BG/(H \backslash G)} \ell_{BG}(\mathbf{C}) .$$

The functor $BH \rightarrow BG/(H \backslash G)$ which sends the object pt to the projection $G \rightarrow H \backslash G$ and the element h of $H = \operatorname{End}_{BH}(pt)$ to the morphism in $BG/(H \backslash G)$ given by left-multiplication by h is an equivalence of categories. Consequently, we get an equivalence

$$\operatorname{colim}_{BG/(H \backslash G)} \ell_{BG}(\mathbf{C}) \simeq \operatorname{colim}_{BH} \ell_{BH}(\mathbf{C}) \stackrel{Thm.14.6}{\simeq} \ell(\mathbf{C} \sharp BH) . \tag{55}$$

■

In the following examples we apply Proposition 14.11 to the construction of equivariant K -theory functors. Let \mathbf{S} be a stable ∞ -category, e.g., the category of spectra. A Bredon-type G -equivariant \mathbf{S} -valued homology theory is determined by a functor

$$\mathbf{Orb}(G) \rightarrow \mathbf{S} .$$

(see e.g. [DL98]). If $K : \mathcal{C}_\infty \rightarrow \mathbf{S}$ is some functor and we fix an object \mathbf{C} in \mathcal{C} , then we can define such a functor by precomposing with the induction functor. We set

$$K_{\mathbf{C}}^G := K \circ J^G(\mathbf{C}) : \mathbf{Orb}(G) \rightarrow \mathbf{S} .$$

By Proposition 14.11 the values of this functor are given by

$$K_{\mathbf{C}}^G(H \backslash G) \simeq K(\ell(\mathbf{C} \sharp BH)) . \tag{56}$$

14.12. EXAMPLE. We let $\mathcal{C} = {}^*\mathbf{Cat}_1$ and $K : {}^*\mathbf{Cat}_1 \rightarrow \mathbf{Sp}$ be the algebraic K -theory functor. The latter is defined as the composition

$$K : {}^*\mathbf{Cat} \xrightarrow{\mathcal{F}} \mathbf{preAdd} \xrightarrow{(-)_{\oplus}} \mathbf{Add} \xrightarrow{K^{alg}} \mathbf{Sp} . \tag{57}$$

Here **preAdd** and **Add** are the ∞ -categories of preadditive and additive categories obtained from the corresponding 1-categories by inverting the exact equivalences. The forgetful functor \mathcal{F} takes the underlying preadditive category of a \mathbb{C} -linear $*$ -category, $(-)_\oplus$ is the additive completion functor (the left-adjoint to the inclusion **Add** \rightarrow **preAdd**), and K^{alg} is the K -theory functor for additive categories. We refer to [BEKW] for further details.

We now fix the object classifier object Δ^0 of $*_{\mathbb{C}}\mathbf{Cat}$ (given by the \mathbb{C} -linear $*$ -category associated to the $*$ -algebra \mathbb{C}). Then we get the functor

$$K_{\Delta^0}^G : \mathbf{Orb}(G) \rightarrow \mathbf{Sp} .$$

Let $K_{ring}^{alg} : \mathbf{Ring} \rightarrow \mathbf{Sp}$ be the algebraic K -theory functor for rings given in terms of K^{alg} as the composition

$$K_{ring}^{alg} : \mathbf{Ring} \rightarrow \mathbf{preAdd} \xrightarrow{(-)_\oplus} \mathbf{Add} \xrightarrow{K^{alg}} \mathbf{Sp} , \tag{58}$$

where the first functor interprets a ring as a pre-additive category with one object. Then we see that $K_{\Delta^0}^G$ has the values

$$K_{\Delta^0}^G(H \setminus G) \simeq K_{ring}^{alg}(\mathbb{C}[H]) .$$

Indeed,

$$K_{\Delta^0}^G(H \setminus G) \stackrel{(56)}{\simeq} K(\ell(\Delta^0 \sharp BH)) \stackrel{Ex.6.13}{\simeq} K(\ell(\mathbb{C}[H])) \stackrel{(57),(58)}{\simeq} K_{ring}^{alg}(\mathbb{C}[H]) .$$

We see that $K_{\Delta^0}^G$ provides a categorical construction of a functor which can be compared with the usual equivariant algebraic K -theory functor as considered e.g. in [DL98, Sec. 2]¹².

14.13. EXAMPLE. We let $\mathcal{C} = C^*\mathbf{Cat}_1$ and $K_1^{top} : C^*\mathbf{Cat}_1 \rightarrow \mathbf{Sp}$ be the topological K -theory functor for C^* -categories. We refer [BE16, Sec. 7.5] for a construction of such a functor as a composition

$$K_1^{top} : C^*\mathbf{Cat}_1 \xrightarrow{A^f} C^*\mathbf{Alg} \xrightarrow{K_{C^*\mathbf{Alg}}^{top}} \mathbf{Sp} ,$$

where A^f is the functor which associates to a C^* -category the free C^* -algebra [Joa03] generated by it, and $K_{C^*\mathbf{Alg}}^{top}$ is the usual topological K -theory functor for C^* -algebras. The subscript 1 indicates that the functor is defined on the 1-category of C^* -categories. In particular, by [BE16, Cor. 7.44] the functor K_1^{top} sends unitary equivalences of C^* -categories to equivalences of spectra and therefore has an essentially unique factorization K^{top} as in

$$\begin{array}{ccc} C^*\mathbf{Cat}_1 & \xrightarrow{K_1^{top}} & \mathbf{Sp} \\ & \searrow \ell & \nearrow K^{top} \\ & & C^*\mathbf{Cat} \end{array} .$$

¹²Note that we have only discussed the values, not the action of the functor on morphisms.

We again fix the object classifier object Δ^0 in $C^*\mathbf{Cat}_1$ and consider the functor

$$K_{\Delta^0}^{top,G} : \mathbf{Orb}(G) \rightarrow \mathbf{Sp} .$$

We then see that $K_{\Delta^0}^{top,G}$ has the values

$$K_{\Delta^0}^{top,G}(H \setminus G) \simeq K_{C^*\mathbf{Alg}}^{top}(C)(C_{\max}^*(H)) .$$

Indeed,

$$K_{\Delta^0}^{top,G}(H \setminus G) \stackrel{(56)}{\simeq} K^{top}(\ell(\Delta^0 \# BH)) \stackrel{Ex.6.13}{\simeq} K^{top}(\ell(C_{\max}^*(H))) \simeq K_{C^*\mathbf{Alg}}^{top}(C_{\max}^*(H)) ,$$

where the last equivalence follows from an inspection of the definition of the K -theory functor K^{top} . We again see that $K_{\Delta^0}^{top,G}$ provides a categorical construction of a functor which can be compared with the topological K -theory functors as considered in [DL98, Sec. 2]. But note that our functor involves the maximal group C^* -algebra, while the functor constructed in [DL98] involves the reduced group C^* -algebra.

References

- [AR94] J. Adámek and J. Rosický. *Locally Presentable and Accessible Categories*. London Math. Soc. Lecture Notes Series 189, 1994.
- [BE16] U. Bunke and A. Engel. Homotopy theory with bornological coarse spaces. <https://arxiv.org/abs/1607.03657>, 2016.
- [Bek00] T. Beke. Sheafifiable homotopy model categories. *Math. Proc. Cambridge Philos. Soc.*, 129(3):447–475, 2000.
- [BEKW] U. Bunke, A. Engel, D. Kasprowski, and Ch. Winges. Homotopy theory with marked additive categories. <https://arxiv.org/abs/1811.04007>.
- [Cis03] D.-Ch. Cisinski. Images directes cohomologiques dans les catégories de modèles. *Annales Mathématiques Blaise Pascal*, pages 195–244, 2003.
- [Cis19] D.-Ch Cisinski. *Higher categories and homotopical algebra*, volume 180 of *Cambridge studies in advanced mathematics*. Cambridge University Press, 2019. Available online under <http://www.mathematik.uni-regensburg.de/cisinski/CatLR.pdf>.
- [Del10] I. Dell’Ambrogio. The unitary symmetric monoidal model category of small C^* -categories. *Homology, Homotopy and Applications, Vol. 14 (2012), No. 2, pp.101-127*, 04 2010.
- [DK80a] W. G. Dwyer and D. M. Kan. Function complexes in homotopical algebra. *Topology*, 19(4):427–440, 1980.

- [DK80b] W.G. Dwyer and D.M. Kan. Simplicial localizations of categories. *J. of Pure and Applied Algebra*, 17:267–284, 1980.
- [DL98] J. F. Davis and W. Lück. Spaces over a Category and Assembly Maps in Isomorphism Conjectures in K - and L -Theory. *K-Theory*, 15:201–252, 1998.
- [Dug01] D. Dugger. Combinatorial model categories have presentations. *Adv. Math.*, 164(1):177–201, 2001.
- [Hin16] V. Hinich. Dwyer-Kan localization revisited. *Homology Homotopy Appl.*, 18(1):27–48, 2016.
- [Hir03] Ph. S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [Hov99] M. Hovey. *Model Categories*. Mathematical Surveys and Monographs vol. 63. Amer. Math. Soc., 1999.
- [Joa03] M. Joachim. K -homology of C^* -categories and symmetric spectra representing K -homology. *Math. Ann.*, 327:641–670, 2003.
- [Joy10] A. Joyal. Model structure on \dagger -categories. <https://ncatlab.org/nlab/show/dagger+category#ModelStructure>, 2010.
- [Lur09] J. Lurie. *Higher topos theory*, volume 170 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2009.
- [Lur17] J. Lurie. Higher algebra. <http://www.math.harvard.edu/~lurie/papers/HA.pdf>, 2017.

Fakultät für Mathematik, Universität Regensburg, 93040 Regensburg, GERMANY
Email: ulrich.bunke@mathematik.uni-regensburg.de

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS L^AT_EX₂ε is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

T_EXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT T_EX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Gabriella Böhm, Wigner Research Centre for Physics: bohm.gabriella@wigner.mta.hu

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Dirk Hoffman, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

F. William Lawvere, State University of New York at Buffalo: wlawvere@buffalo.edu

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Robert Paré, Dalhousie University: pare@mathstat.dal.ca

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be

R. J. Wood, Dalhousie University: rjwood@mathstat.dal.ca