

PROPS FOR INVOLUTIVE MONOIDS AND INVOLUTIVE BIMONOIDS

DANIEL GRAVES

ABSTRACT. The category of involutive non-commutative sets encodes the structure of an involution compatible with a (co)associative (co)multiplication. We prove that the category of involutive bimonoids in a symmetric monoidal category is equivalent to the category of algebras over a PROP constructed from the category of involutive non-commutative sets.

Introduction

The categorification of algebras over a unital commutative ring k to algebras over a PROP was first introduced by Markl in order to study the deformation theory of algebras [Mar96]. In that paper he defined PROPs, in terms of generators and relations, whose categories of algebras are equivalent to the category of associative algebras, the category of commutative algebras and the category of bialgebras over k [Mar96, Examples 2.5, 2.6 and 2.7].

Pirashvili [Pir02] gave an explicit description of a PROP that categorified associative algebras, commutative algebras and bialgebras in the category of vector spaces over a field. This PROP is constructed from the category of non-commutative sets, introduced by Feigin and Tsygan [FT87, A10], using the generalized Quillen Q -construction of Fiedorowicz and Loday [FL91, 2.5]. An alternative approach, using distributive laws for PROPs, was given by Lack [Lac04, Section 5]. In this setting, Pirashvili's PROP is described as a composite constructed from the PRO of finite ordinals and its opposite category and the result is shown to be more general, holding for bimonoids in a symmetric monoidal category.

In this paper we combine both of these methods. We introduce the PROP of *involutive non-commutative sets*, denoted $\mathcal{IF}(as)$. Using the machinery of [Lac04] we describe a composite PROP constructed from $\mathcal{IF}(as)$ whose algebras in a symmetric monoidal category are the involutive bimonoids.

The paper is organized as follows. In Section 1 we recall the definition of involutive bimonoid in a symmetric monoidal category. In Section 2 we recall the definitions of PRO and PROP, together with some examples. In Section 3 we define the PROP of involutive

Received by the editors 2020-06-09 and, in final form, 2020-09-18.

Transmitted by Ross Street. Published on 2020-09-22.

2020 Mathematics Subject Classification: 16T10, 18M05, 18M85.

Key words and phrases: involutive non-commutative sets, bimonoid, bialgebras, PROP, symmetric monoidal categories.

© Daniel Graves, 2020. Permission to copy for private use granted.

non-commutative sets, $\mathcal{IF}(as)$, and prove that the category of algebras of $\mathcal{IF}(as)$ in a symmetric monoidal category \mathbf{C} is equivalent to the category of involutive monoids in \mathbf{C} . In Section 4 we construct a double category from $\mathcal{IF}(as)$ whose bimorphisms encode the compatibility conditions for an involutive bimonoid. In Section 5 we construct a composite PROP, in the sense of [Lac04, Section 4], from the PROP $\mathcal{IF}(as)$ and its opposite category. This can be seen as a composite of the PROPs for involutive monoids and involutive comonoids described in Section 3, where the compatibility of the two is encoded in the double category described in Section 4. We prove that the category of algebras of this composite PROP in a symmetric monoidal category \mathbf{C} is equivalent to the category of involutive bimonoids in \mathbf{C} .

Acknowledgements

I would like to thank Sarah Whitehouse for all her guidance and encouragement. I am grateful to James Brotherston, James Cranch and the anonymous referees for their helpful suggestions.

1. Involutive monoids and involutive bimonoids

1.1. DEFINITION. A monoid M in a symmetric monoidal category \mathbf{C} is called involutive if it comes equipped with a monoid morphism $j: M \rightarrow M^{op}$ satisfying $j^2 = id_M$. We denote the category of involutive monoids and involution-preserving morphisms by $\mathbf{IMon}(\mathbf{C})$.

We denote the category of involutive comonoids in \mathbf{C} by $\mathbf{IComon}(\mathbf{C}) = \mathbf{IMon}(\mathbf{C}^{op})^{op}$.

A bimonoid B in \mathbf{C} is said to be involutive if it comes equipped with a bimonoid morphism $j: B \rightarrow B^{op, cop}$ such that $j^2 = id_B$. We denote the category of involutive bimonoids in \mathbf{C} by $\mathbf{IBimon}(\mathbf{C})$.

2. PROs and PROPs

2.1. DEFINITION. For $n \geq 1$ we define \underline{n} to be the set $\{1, \dots, n\}$. We define $\underline{0} = \emptyset$.

2.2. DEFINITION. A PRO \mathbb{T} is a strict monoidal category whose objects are the sets \underline{n} for $n \geq 0$ and whose tensor product is given by the disjoint union. For a monoidal category \mathbf{C} , an algebra of \mathbb{T} in \mathbf{C} is a strict monoidal functor $\mathbb{T} \rightarrow \mathbf{C}$.

2.3. DEFINITION. A PROP \mathbf{P} is a symmetric strict monoidal category whose objects are the sets \underline{n} for $n \geq 0$ with tensor product given by the disjoint union. For a symmetric monoidal category \mathbf{C} , a \mathbf{P} -algebra in \mathbf{C} is a symmetric strict monoidal functor $\mathbf{P} \rightarrow \mathbf{C}$. We denote the category of \mathbf{P} -algebras in \mathbf{C} and natural transformations by $\mathbf{Alg}(\mathbf{P}, \mathbf{C})$.

2.4. EXAMPLE. We denote by \mathbb{D} the PRO of finite ordinals and order-preserving maps as in [Lac04, 2.2]. For a strict monoidal category \mathbf{C} , an algebra of \mathbb{D} in \mathbf{C} is a monoid in \mathbf{C} , see [ML98, VII 5].

2.5. DEFINITION. Let $C_2 = \langle t \mid t^2 = 1 \rangle$. Let \mathbb{C}_2 be the PRO such that $\text{Hom}_{\mathbb{C}_2}(\underline{n}, \underline{m})$ is empty if $n \neq m$ and $\text{Hom}_{\mathbb{C}_2}(\underline{n}, \underline{n}) = C_2^n$. The disjoint union of morphisms corresponds to the product of group elements.

2.6. EXAMPLE. Following [Lac04, 2.4], let \mathbb{P} denote the PRO of finite sets and bijections.

2.7. REMARK. An equivalent definition of a PROP is as a PRO \mathbb{T} with a map of PROs $\mathbb{P} \rightarrow \mathbb{T}$. Therefore \mathbb{P} is a PROP. Given a PROP \mathbf{P} we will denote its underlying PRO by \mathbf{P}_0 .

2.8. EXAMPLE. Following [Lac04, 5.1, 5.2], we denote by \mathbb{F} the PROP of finite sets and finite set maps. For a symmetric monoidal category \mathbf{C} , the category $\mathbf{Alg}(\mathbb{F}, \mathbf{C})$ is equivalent to the category of commutative monoids in \mathbf{C} and $\mathbf{Alg}(\mathbb{F}^{op}, \mathbf{C})$ is equivalent to the category of cocommutative comonoids in \mathbf{C} .

We can form new PROs and PROPs via the notion of a *distributive law* as defined in [Lac04, Section 3] and [RW02, Section 2]. In particular we will form a composite PROP from \mathbb{D} , \mathbb{P} and \mathbb{C}_2 whose structure is that of the hyperoctahedral category defined by Fiedorowicz and Loday [FL91, Section 3]. We begin by recalling the definition of the hyperoctahedral groups.

2.9. DEFINITION. For $n \geq 1$, the hyperoctahedral group H_n is defined to be the semi-direct product $C_2^n \rtimes \Sigma_n$ where Σ_n acts on C_2^n by permuting the factors.

2.10. EXAMPLE. A pair (x, f) where $f \in \text{Hom}_{\mathbb{F}}(\underline{n}, \underline{m})$ and $x \in \text{Hom}_{\mathbb{C}_2}(\underline{m}, \underline{m})$ determines a unique pair (f, x') where f has remained unchanged and, if $x = (g_1, \dots, g_m)$, $x' = (g_{f(1)}, \dots, g_{f(n)}) \in \text{Hom}_{\mathbb{C}_2}(\underline{n}, \underline{n})$. A straightforward check of the relations in [RW02, 2.4] and [Lac04, 3.7] shows that this defines a distributive law $\mathbb{C}_2 \otimes \mathbb{F} \rightarrow \mathbb{F} \otimes \mathbb{C}_2$ which is compatible with the monoidal structures of \mathbb{F} and \mathbb{C}_2 .

By [Lac04, Theorem 3.8], $\mathbb{F} \otimes \mathbb{C}_2$ is a PRO such that morphisms in $\text{Hom}_{\mathbb{F} \otimes \mathbb{C}_2}(\underline{n}, \underline{m})$ can be written uniquely as pairs (f, x) with $x \in \text{Hom}_{\mathbb{C}_2}(\underline{n}, \underline{n})$ and $f \in \text{Hom}_{\mathbb{F}}(\underline{n}, \underline{m})$ with composition determined by the distributive law. In fact, $\mathbb{F} \otimes \mathbb{C}_2$ has a canonical PROP structure induced from the PROP structure on \mathbb{F} .

2.11. EXAMPLE. A pair (x, σ) where $\sigma \in \text{Hom}_{\mathbb{P}}(\underline{n}, \underline{n})$ and $x \in \text{Hom}_{\mathbb{C}_2}(\underline{n}, \underline{n})$ determines a unique pair (σ, x') where σ has remained unchanged and, if $x = (g_1, \dots, g_n)$, $x' = (g_{\sigma(1)}, \dots, g_{\sigma(n)})$.

Similarly to the previous example, this is a distributive law compatible with the monoidal structure of \mathbb{P} and \mathbb{C}_2 and we have a PRO $\mathbb{P} \otimes \mathbb{C}_2$.

We observe that $\text{Hom}_{\mathbb{P} \otimes \mathbb{C}_2}(\underline{n}, \underline{n})$ is isomorphic to H_n , the n^{th} hyperoctahedral group as defined in Definition 2.9. We therefore write $\mathbb{H} = \mathbb{P} \otimes \mathbb{C}_2$. We note that \mathbb{H} has a canonical PROP structure induced from \mathbb{P} . We refer to \mathbb{H} as the PROP of hyperoctahedral groups.

2.12. **EXAMPLE.** Given a pair (g, φ) where $\varphi \in \text{Hom}_{\mathbb{D}}(\underline{n}, \underline{m})$ and $g \in \text{Hom}_{\mathbb{H}}(\underline{m}, \underline{m})$ there is a unique pair $(g_*(\varphi), \varphi^*(g))$ where $\varphi^*(g) \in \text{Hom}_{\mathbb{H}}(\underline{n}, \underline{n})$ and $g_*(\varphi) \in \text{Hom}_{\mathbb{D}}(\underline{n}, \underline{m})$, as constructed in [FL91, 3.1]. The fact that these assignments satisfy the relations of [RW02, 2.4] follows from the fact that they satisfy the relations of a crossed simplicial group given in [FL91, 1.6] and a routine check shows that they respect the monoidal structures of \mathbb{D} and \mathbb{H} .

We therefore have a PRO $\mathbb{D} \otimes \mathbb{H}$ defined similarly to the examples above. In fact, $\mathbb{D} \otimes \mathbb{H}$ has a canonical PROP structure induced from \mathbb{H} .

2.13. **PROPOSITION.** *Let \mathbf{C} be a symmetric monoidal category. There is an equivalence of categories*

$$\mathbf{Alg}(\mathbb{D} \otimes \mathbb{H}, \mathbf{C}) \simeq \mathbf{IMon}(\mathbf{C}).$$

PROOF. By [Lac04, 3.10], a $(\mathbb{D} \otimes \mathbb{H})_0$ -algebra structure on an object M of \mathbf{C} consists of a \mathbb{D} -algebra structure and a \mathbb{H} -algebra structure subject to a compatibility condition. A \mathbb{D} -algebra structure is a monoid structure. A \mathbb{H} -algebra is an object M together with a morphism $j: M \rightarrow M$ satisfying $j^2 = id_M$ and, for each element $g \in H_n$, a morphism $M^{\otimes n} \rightarrow M^{\otimes n}$ given by applying j to the tensor factors according to the element of C_2^n followed by an isomorphism determined by the element of Σ_n . Arguing analogously to [Lac04, 5.5] a $(\mathbb{D} \otimes \mathbb{H})_0$ -algebra structure is a $\mathbb{D} \otimes \mathbb{H}$ -algebra structure if and only if the only isomorphisms $M^{\otimes n} \rightarrow M^{\otimes n}$ are those induced from the symmetry isomorphisms. The compatibility condition is precisely the condition requiring j to be an involution compatible with the monoid structure. Finally, a morphism in \mathbf{C} is a map of involutive monoids if and only if it respects the \mathbb{D} -algebra structure and the \mathbb{H} -algebra structure. By [Lac04, 3.12], this is true if and only if it respects the $\mathbb{D} \otimes \mathbb{H}$ -algebra structure. ■

2.14. **REMARK.** This result tells us that the PROP governing the structure of an involutive monoid can be thought of as a composite of PROPs governing the structure of a monoid and the structure of an involution respectively. We will give an explicit description of this category, where the technicalities of distributive laws are distilled into data on the preimages of set maps in Section 3.

We have chosen to emphasize the connection between involutive monoids and the category associated to the hyperoctahedral crossed simplicial group. One advantage of this approach is that the distributive laws employed are already well-known, being the composition in hyperoctahedral groups and the hyperoctahedral category. It is also an interesting new application of the hyperoctahedral crossed simplicial group: the other known applications are found in the field of equivariant stable homotopy theory!

An alternative method of proof would be to begin with the composite PROP $\mathbb{D} \otimes \mathbb{P}$ of [Lac04, 3.14], define a distributive law between this and the PRO \mathbb{C}_2 and to analyse the resulting composite.

2.15. **COROLLARY.** *There is an equivalence of categories*

$$\mathbf{Alg}((\mathbb{D} \otimes \mathbb{H})^{op}, \mathbf{C}) \simeq \mathbf{IComon}(\mathbf{C}).$$

PROOF. We observe that

$$\mathbf{IComon}(\mathbf{C}) = \mathbf{IMon}(\mathbf{C}^{op})^{op} \simeq \mathbf{Alg}((\mathbb{D} \otimes \mathbb{H}), \mathbf{C}^{op})^{op} = \mathbf{Alg}((\mathbb{D} \otimes \mathbb{H})^{op}, \mathbf{C})$$

as required. ■

2.16. PROPOSITION. *Let \mathbf{C} be a symmetric monoidal category. There is an equivalence of categories between $\mathbf{Alg}(\mathbb{F} \otimes \mathbb{C}_2, \mathbf{C})$ and the category of involutive commutative monoids in \mathbf{C} . The category $\mathbf{Alg}((\mathbb{F} \otimes \mathbb{C}_2)^{op}, \mathbf{C})$ is equivalent to the category of involutive comonoids in \mathbf{C} .*

PROOF. The proof of the first equivalence is similar to Proposition 2.13. We note that an \mathbb{F} -algebra structure is a commutative monoid structure. A \mathbb{C}_2 -algebra structure consists of an object M in \mathbf{C} together with a morphism $j: M \rightarrow M$ satisfying $j^2 = id_M$ and for each element of C_2^n a morphism $M^{\otimes n} \rightarrow M^{\otimes n}$ defined by applying j to the tensor factors according to the element of C_2^n . An $(\mathbb{F} \otimes \mathbb{C}_2)_0$ -structure is a $\mathbb{F} \otimes \mathbb{C}_2$ -structure if and only if the only isomorphisms $M^{\otimes n} \rightarrow M^{\otimes n}$ are those induced from the symmetry isomorphisms. The compatibility condition in this case is the condition that requires j to be an involution compatible with a commutative monoid structure. Finally we note that a morphism in \mathbf{C} is a map of involutive commutative monoids if and only if it preserves both the \mathbb{F} -algebra structure and the \mathbb{C}_2 -algebra structure. The second equivalence follows a similar argument to Corollary 2.15. ■


3. The PROPs $\mathcal{IF}(as)$ and \mathcal{IF}

In the previous section we described a PROP for involutive monoids as a composite. In this section we provide an explicit description of this PROP, called the category of *involutive non-commutative sets*. A variant of this category first appeared in the author’s thesis [Gra19, Part V]. This category takes the technicalities of the composition of pairs defined via a distributive law and presents it as simple structure on the preimages of maps of finite sets. We shall also see, in Section 4, that we can construct a double category from the category of involutive non-commutative sets whose bimorphisms encode the structure of an involutive bimonoid.

The PROP of involutive, non-commutative sets, $\mathcal{IF}(as)$ will have as objects the sets \underline{n} of Definition 2.1 for $n \geq 0$. An element $f \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$ will be a map of sets such that the preimage of each singleton $i \in \underline{m}$ is a totally ordered set such that each element comes adorned with a superscript label from the group $C_2 = \langle t \mid t^2 = 1 \rangle$. Note that for $m \geq 1$, the set $\text{Hom}_{\mathcal{IF}(as)}(\underline{0}, \underline{m})$ will be the singleton set consisting of the unique set map $\emptyset \rightarrow \underline{m}$ and $\text{Hom}_{\mathcal{IF}(as)}(\underline{m}, \underline{0})$ will be the empty set.

3.1. REMARK. Henceforth we will say that a morphism in $\mathcal{IF}(as)$ is a map of sets together with a *labelled, ordered set* for each preimage. In particular, note that we will use *preimage* to mean preimage of a singleton.

3.2. EXAMPLE. Let $f \in \text{Hom}_{\mathcal{IF}(as)}(\underline{5}, \underline{4})$ have underlying map of sets



with the following labelled, ordered sets as preimages:

$$f^{-1}(1) = \{2^1\}, \quad f^{-1}(2) = \{1^t\}, \quad f^{-1}(3) = \{4^t < 5^1\} \text{ and } f^{-1}(4) = \{3^t\}.$$

We will denote composition in $\mathcal{IF}(as)$ by \bullet in order to distinguish from the composition of maps of sets. In particular, we use \circ for two morphisms in $\mathcal{IF}(as)$ if we are referring to the composite of the underlying maps of sets. In order to ease notation we have chosen not to introduce notation for the forgetful functor $\mathcal{IF}(as) \rightarrow \mathbf{Set}$.

Let $f_1 \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$ and $f_2 \in \text{Hom}_{\mathcal{IF}(as)}(\underline{m}, \underline{l})$. In order to define the composite $f_2 \bullet f_1 \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{l})$ we must provide a map of sets and describe the labelled total orderings on each of the preimages.

As a map of sets, $f_2 \bullet f_1$ is the composite of the underlying map of sets $f_2 \circ f_1$. In order to specify a labelled, ordered set for the preimage of each singleton in \underline{l} under the composite we first make a definition.

3.3. DEFINITION. We define an action of C_2 , which will be denoted by a superscript, on finite, ordered sets with C_2 -labels by

$$\{j_1^{\alpha_{j_1}} < \dots < j_r^{\alpha_{j_r}}\}^t = \{j_r^{t\alpha_{j_r}} < \dots < j_1^{t\alpha_{j_1}}\}.$$

That is, we invert the ordering and multiply each label by $t \in C_2$.

3.4. DEFINITION. Let $f_1 \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$ and $f_2 \in \text{Hom}_{\mathcal{IF}(as)}(\underline{m}, \underline{l})$. We define $f_2 \bullet f_1 \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{l})$ to have underlying map of sets $f_2 \circ f_1$. We define the labelled totally ordered set $(f_2 \bullet f_1)^{-1}(i)$ to be the ordered disjoint union of labelled, ordered sets

$$\coprod_{j^{\alpha_j} \in f_2^{-1}(i)} f_1^{-1}(j)^{\alpha_j}.$$

3.5. DEFINITION. The PROP of involutive, non-commutative sets, $\mathcal{IF}(as)$, has as objects the sets \underline{n} of Definition 2.1 for $n \geq 0$. An element of $\text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$ is a map of sets with a total ordering on each preimage such that each element of the domain comes adorned with a superscript label from the group C_2 . Composition of morphisms is as defined in Definition 3.4. The symmetry isomorphisms are given by block permutations.

3.6. REMARK. For each $m \geq 1$, the set $\text{Hom}_{\mathcal{IF}(as)}(\underline{0}, \underline{m})$ is the singleton set consisting of the unique set map $\emptyset \rightarrow \underline{m}$ and $\text{Hom}_{\mathcal{IF}(as)}(\underline{m}, \underline{0})$ is the empty set.

3.7. **REMARK.** Recall the PROP of non-commutative sets, $\mathcal{F}(as)$, from [Pir02, Section 3]. That is, the category whose objects are the sets \underline{n} for $n \geq 0$ and whose morphisms are maps of sets with a total ordering on the preimage of each singleton in the codomain. We observe that $\mathcal{F}(as)$ is isomorphic to the subcategory of $\mathcal{IF}(as)$ which contains only the morphisms for which every label is $1 \in C_2$.

3.8. **DEFINITION.** We define the fundamental morphisms m , u and i of $\mathcal{IF}(as)$ as follows.

- Let $m \in \text{Hom}_{\mathcal{IF}(as)}(\underline{2}, \underline{1})$ be defined by $m^{-1}(1) = \{1^1 < 2^1\}$,
- let u be the unique morphism in $\text{Hom}_{\mathcal{IF}(as)}(\underline{0}, \underline{1})$ and
- let $i \in \text{Hom}_{\mathcal{IF}(as)}(\underline{1}, \underline{1})$ be defined by $i^{-1}(1) = \{1^t\}$.

3.9. **REMARK.** The morphism m will encode the multiplication and comultiplication in a bimonoid, the morphism u will encode the unit and counit and the morphism i will encode the involution.

3.10. **REMARK.** We note that $\mathcal{IF}(as)$ contains the morphisms of the PRO of finite ordinals \mathbb{D} . These are the order-preserving maps of sets with the canonical total ordering on each preimage with each label being $1 \in C_2$. Furthermore, $\mathcal{IF}(as)$ contains the morphisms of the PROP of hyperoctahedral groups \mathbb{H} . These are the bijections in $\mathcal{IF}(as)$.

3.11. **PROPOSITION.** *There is an isomorphism of PROPs $\mathcal{IF}(as) \cong \mathbb{D} \otimes \mathbb{H}$.*

PROOF. Consider the data of a morphism $f \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$. The total ordering data on preimages determines a unique bijection of the set \underline{n} with a C_2 -label for each singleton preimage determined by the labelling data. That is, the preimage data determines a unique bijection $g \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{n})$. There is then a unique morphism $\varphi \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$ such that φ is order-preserving, with the canonical total ordering on each preimage, every label is $1 \in C_2$ and $f = \varphi \bullet g$. In other words, any morphism in $\mathcal{IF}(as)$ can be written uniquely as a composite of a morphism in \mathbb{H} followed by a morphism in \mathbb{D} .

For composable morphisms f_1 and f_2 in $\mathcal{IF}(as)$ write $f_2 \bullet f_1 = \varphi_2 \bullet g_2 \bullet \varphi_1 \bullet g_1$. A straightforward check shows that the composite $g_2 \bullet \varphi_1$ in $\mathcal{IF}(as)$ is equal to the composite $g_{2\star}(\varphi_1) \bullet \varphi_1^*(g_2)$, where $g_{2\star}(\varphi_1) \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{m})$ is an order-preserving map and $\varphi_1^*(g_2) \in \text{Hom}_{\mathcal{IF}(as)}(\underline{n}, \underline{n})$ is a bijection, both maps being determined using the structure of the hyperoctahedral crossed simplicial group described in [FL91, 3.1]. It follows that there is an isomorphism of PROPs $\mathcal{IF}(as) \cong \mathbb{D} \otimes \mathbb{H}$ given by sending a morphism $f = \varphi \bullet g$ in $\mathcal{IF}(as)$ to the pair (φ, g) . ■

3.12. **COROLLARY.** *Let \mathbf{C} be a symmetric monoidal category. There are equivalences of categories*

$$\text{Alg}(\mathcal{IF}(as), \mathbf{C}) \simeq \text{IMon}(\mathbf{C}) \quad \text{and} \quad \text{Alg}(\mathcal{IF}(as)^{op}, \mathbf{C}) \simeq \text{IComon}(\mathbf{C}).$$

PROOF. This follows from Proposition 3.11, Proposition 2.13 and Corollary 2.15. ■

3.13. DEFINITION. Let \mathcal{IF} be the category whose objects are the sets \underline{n} of Definition 2.1 for $n \geq 0$. A morphism in \mathcal{IF} is a map of sets such that the elements of the preimage of each singleton in the codomain come adorned with a label from C_2 . Composition is given by composition of set maps and multiplication of labels.

3.14. PROPOSITION. There is an isomorphism of PROPs $\mathcal{IF} \cong \mathbb{F} \otimes C_2$.

PROOF. The method of proof is similar to Proposition 3.11. ■

3.15. COROLLARY. Let \mathbf{C} be a symmetric monoidal category. The categories $\mathbf{Alg}(\mathcal{IF}, \mathbf{C})$ and $\mathbf{Alg}(\mathcal{IF}^{op}, \mathbf{C})$ are equivalent to the category of involutive monoids in \mathbf{C} and the category of involutive cocommutative comonoids in \mathbf{C} respectively.

PROOF. This follows from Proposition 2.16 and Proposition 3.14. ■

4. Double categories

We construct a double category from $\mathcal{IF}(as)$. The bimorphisms of this double category precisely encode the structure of an involutive bimonoid in a symmetric monoidal category. This double category also possesses extra structure: it satisfies the star condition of [FL91, 2.3]. This extra structure will be used in Section 5 to construct a PROP which governs the structure of an involutive bimonoid. We also construct a double category from \mathcal{IF} and two double categories that combine the structure of $\mathcal{IF}(as)$ and \mathcal{IF} which will encode commutativity and cocommutativity.

Recall from [FL91, Section 2.1] that a small *double category* \mathbf{D} consists of a set of objects, a set of horizontal morphisms, a set of vertical morphisms and a set of bimorphisms subject to natural composition identities.

4.1. DEFINITION. The double category $\mathcal{IF}(as)_2$ has as objects the objects of $\mathcal{IF}(as)$. Furthermore, the sets of horizontal and vertical morphisms in $\mathcal{IF}(as)_2$ are both equal to the set of all morphisms in $\mathcal{IF}(as)$. A bimorphism in $\mathcal{IF}(as)_2$ is a not necessarily commutative square

$$\begin{array}{ccc} \underline{n} & \xrightarrow{f_1} & \underline{p} \\ \varphi_1 \downarrow & & \downarrow \varphi \\ \underline{m} & \xrightarrow{f} & \underline{q} \end{array}$$

of morphisms in $\mathcal{IF}(as)$ such that

- the underlying diagram of finite sets is a pullback square,
- for all $x \in \underline{m}$ the map $\varphi_1^{-1}(x) \rightarrow \varphi^{-1}(f(x))$ induced by f_1 is an isomorphism of labelled, ordered sets and
- for all $y \in \underline{p}$ the map $f_1^{-1}(y) \rightarrow f^{-1}(\varphi(y))$ induced by φ_1 is an isomorphism of labelled ordered sets.

4.2. **REMARK.** The composition laws of a double category can be verified using the fact that the composite of pullback squares is itself a pullback square and using the composition rule for morphisms in $\mathcal{IF}(as)$ described in Definition 3.4.

4.3. **DEFINITION.** Let B_1, B_2, B_3, B_4 and J denote the bimorphisms

$$\begin{array}{ccccc}
 \underline{4} & \xrightarrow{m^{\Pi_2}} & \underline{2} & & \underline{0} & \xrightarrow{id_0} & \underline{0} & & \underline{0} & \xrightarrow{id_0} & \underline{0} & & \underline{0} & \xrightarrow{u^2} & \underline{2} & & \underline{1} & \xrightarrow{id_1} & \underline{1} \\
 m^{\Pi_2} \bullet \tau_{2,3} \downarrow & & \downarrow m & & id_0 \downarrow & & \downarrow u & & u^2 \downarrow & & \downarrow u & & id_0 \downarrow & & \downarrow m & & id_1 \downarrow & & \downarrow i \\
 \underline{2} & \xrightarrow{m} & \underline{1} & & \underline{0} & \xrightarrow{u} & \underline{1} & & \underline{2} & \xrightarrow{m} & \underline{1} & & \underline{0} & \xrightarrow{u} & \underline{1} & & \underline{1} & \xrightarrow{i} & \underline{1}
 \end{array}$$

respectively in $\mathcal{IF}(as)_2$. Here $\tau_{2,3}$ is the transposition $(2\ 3)$ with the label $1 \in C_2$ for each preimage. We call B_1, B_2, B_3, B_4 and J the fundamental bimorphisms of $\mathcal{IF}(as)_2$.

4.4. **REMARK.** The fundamental bimorphisms encode the compatibility conditions of an involutive bimonoid. The notation is chosen such that the bimorphisms B_1 to B_4 encode the compatibility conditions of a bimonoid and J encodes the compatibility condition of an involution.

4.5. **DEFINITION.** The double category \mathcal{IF}_2 is defined similarly to $\mathcal{IF}(as)_2$; the objects are those of \mathcal{IF} , the sets of horizontal and vertical morphisms are the set of morphisms in \mathcal{IF} and the bimorphisms are defined similarly to the bimorphisms of $\mathcal{IF}(as)_2$.

4.6. **DEFINITION.** The double category \mathcal{V} has as objects the objects of $\mathcal{IF}(as)$. The set of vertical morphisms is the set of morphisms in $\mathcal{IF}(as)$. The set of horizontal morphisms is the set of morphisms in \mathcal{IF} . The bimorphisms are defined similarly to those of $\mathcal{IF}(as)_2$ except that the horizontal morphisms are now in \mathcal{IF} .

The double category \mathcal{H} is defined similarly; the set of horizontal morphisms is the set of morphisms in $\mathcal{IF}(as)$, the set of vertical morphisms is the set of morphisms in \mathcal{IF} and the bimorphisms are defined similarly to those of $\mathcal{IF}(as)_2$ except that the vertical morphisms are in \mathcal{IF} .

4.7. **REMARK.** Recall from [FL91, 2.3] that a double category \mathbf{D} is said to satisfy the *star condition* if a horizontal morphism and a vertical morphism with the same codomain determine a unique bimorphism in \mathbf{D} .

Let $\mathbf{D} = \mathcal{IF}(as)_2, \mathcal{IF}_2, \mathcal{V}$ or \mathcal{H} . Given a horizontal morphism $f: \underline{m} \rightarrow \underline{q}$ and a vertical morphism $\varphi: \underline{p} \rightarrow \underline{q}$ in \mathbf{D} we determine a unique bimorphism by first taking the pullback of the underlying maps of sets. The resulting maps have a unique lift to the category \mathbf{D} where the preimage data is induced from f and φ using the conditions on bimorphisms. Therefore these four double categories satisfy the star condition.

5. Involutive bimonoids

We construct composite PROPs, in the sense of [Lac04, Section 4], from the PROPs $\mathcal{IF}(as), \mathcal{IF}$ and their opposites. We prove that the category of algebras of the composite

PROP constructed from $\mathcal{IF}(as)$ and its opposite in a symmetric monoidal category \mathbf{C} is equivalent to the category of involutive bimonoids in \mathbf{C} .

5.1. PROPOSITION. *There exist composite PROPs*

$$\mathcal{IF} \otimes_{\mathbb{P}} \mathcal{IF}^{op}, \quad \mathcal{IF} \otimes_{\mathbb{P}} \mathcal{IF}(as)^{op}, \quad \mathcal{IF}(as) \otimes_{\mathbb{P}} \mathcal{IF}^{op} \quad \text{and} \quad \mathcal{IF}(as) \otimes_{\mathbb{P}} \mathcal{IF}(as)^{op}.$$

PROOF. We provide the details for the case $\mathcal{IF} \otimes_{\mathbb{P}} \mathcal{IF}^{op}$. The others are similar.

A pair (φ, f) where $f \in \text{Hom}_{\mathcal{IF}}(\underline{m}, \underline{q})$ and $\varphi \in \text{Hom}_{\mathcal{IF}^{op}}(\underline{q}, \underline{p})$ can be written as a diagram

$$\begin{array}{ccc} & & \underline{p} \\ & & \downarrow \varphi \\ \underline{m} & \xrightarrow{f} & \underline{q} \end{array}$$

in \mathcal{IF} . By the star condition for the double category \mathcal{IF}_2 , there exist unique morphisms φ_1 and f_1 in \mathcal{IF} forming a bimorphism

$$\begin{array}{ccc} \underline{n} & \xrightarrow{f_1} & \underline{p} \\ \varphi_1 \downarrow & & \downarrow \varphi \\ \underline{m} & \xrightarrow{f} & \underline{q} \end{array}$$

in \mathcal{IF}_2 .

We observe that the assignment $(\varphi, f) \mapsto (f_1, \varphi_1)$ defines a distributive law of PROs $\mathcal{IF}^{op} \otimes \mathcal{IF} \rightarrow \mathcal{IF} \otimes \mathcal{IF}^{op}$, in the sense of [Lac04, 3.6]. The star condition, together with the composition rule for bimorphisms, ensures that the equations for a distributive law are satisfied and compatibility with the monoidal structure follows from the compatibility of the star condition with the disjoint union.

Furthermore, since both \mathcal{IF} and \mathcal{IF}^{op} are PROPs, we observe that $\mathcal{IF} \otimes \mathcal{IF}^{op}$ has a PROP structure.

A morphism in $\mathcal{IF} \otimes \mathcal{IF}^{op}$ from \underline{n} to \underline{m} can be written as a span

$$\underline{n} \xleftarrow{\varphi} \underline{p} \xrightarrow{f} \underline{m}.$$

Two spans

$$\underline{n} \xleftarrow{\varphi} \underline{p} \xrightarrow{f} \underline{m} \quad \text{and} \quad \underline{n} \xleftarrow{\varphi_1} \underline{p} \xrightarrow{f_1} \underline{m}$$

are said to be *equivalent* if there exists a bijection $h: \underline{p} \rightarrow \underline{p}$ in \mathcal{IF} such that $\varphi_1 \circ h = \varphi$ and $f_1 \circ h = f$.

It follows from [Lac04, Theorem 4.6] that $\mathcal{IF} \otimes_{\mathbb{P}} \mathcal{IF}^{op}$, that is the category obtained from $\mathcal{IF} \otimes \mathcal{IF}^{op}$ by identifying equivalent spans, is a composite PROP defined via a distributive law induced from the one defined for PROs.

The remaining three cases are similar, making use of the star condition from the double categories \mathcal{V} , \mathcal{H} and $\mathcal{IF}(as)_2$ respectively. ■

5.2. DEFINITION. For ease of notation, let $Q = \mathcal{IF}(as) \otimes_{\mathbb{P}} \mathcal{IF}(as)^{op}$. Let $Q_{\mathcal{V}} = \mathcal{IF} \otimes_{\mathbb{P}} \mathcal{IF}(as)^{op}$. Let $Q_{\mathcal{H}} = \mathcal{IF}^{op} \otimes_{\mathbb{P}} \mathcal{IF}(as)$. Let $Q_{\mathcal{IF}} = \mathcal{IF} \otimes_{\mathbb{P}} \mathcal{IF}^{op}$.

5.3. THEOREM. Let \mathbf{C} be a symmetric monoidal category. There is an equivalence of categories

$$\mathbf{Alg}(Q, \mathbf{C}) \simeq \mathbf{IBimon}(\mathbf{C}).$$

PROOF. By [Lac04, Proposition 4.7], an algebra for Q in \mathbf{C} consists of an object M with an $\mathcal{IF}(as)$ -algebra structure and an $\mathcal{IF}(as)^{op}$ -algebra structure subject to compatibility conditions. An $\mathcal{IF}(as)$ -algebra structure is an involutive monoid structure and an $\mathcal{IF}(as)^{op}$ -algebra structure is an involutive comonoid structure. Let F be the $\mathcal{IF}(as)$ -algebra and let G be the $\mathcal{IF}(as)^{op}$ -algebra. The compatibility condition requires that for every bimorphism

$$\begin{array}{ccc} \underline{n} & \xrightarrow{f_1} & \underline{p} \\ \varphi_1 \downarrow & & \downarrow \varphi \\ \underline{m} & \xrightarrow{f} & \underline{q} \end{array}$$

in the double category $\mathcal{IF}(as)_2$, the diagram

$$\begin{array}{ccc} M^{\otimes n} & \xrightarrow{F(f_1)} & M^{\otimes p} \\ G(\varphi_1) \uparrow & & \uparrow G(\varphi) \\ M^{\otimes m} & \xrightarrow{F(f)} & M^{\otimes q} \end{array}$$

commutes. Arguing analogously to [Lac04, 5.3], it suffices to have commutativity for the fundamental bimorphisms of Definition 4.3. These are precisely the conditions requiring M to be an involutive bimonoid. Finally we observe that a morphism in \mathbf{C} is a morphism of involutive bimonoids if and only if it preserves the $\mathcal{IF}(as)$ -algebra structure and the $\mathcal{IF}(as)^{op}$ -algebra structure. By [Lac04, 4.8] this is true if and only if it preserves the Q -algebra structure. ■

5.4. REMARK. The theorem tells us that the PROP governing the structure of involutive bimonoids is a composite of the PROPs for involutive monoids and involutive comonoids where the compatibility conditions are precisely the fundamental bimorphisms of the category $\mathcal{IF}(as)_2$ and the distributive law is determined by the star condition. We have chosen this method of proof as we believe that the double category $\mathcal{IF}(as)_2$ most neatly encapsulates the structure required to construct the PROP Q .

An alternative method of proof would be to define a distributive law between the PROP given in [Lac04, 5.9] and the PRO \mathbf{C}_2 and analyse the resulting composite. The technical details of such a proof are quite similar to those we have used.

5.5. THEOREM. *Let \mathbf{C} be a symmetric monoidal category.*

1. *The category $\mathbf{Alg}(Q_{\mathcal{V}}, \mathbf{C})$ is equivalent to the category of involutive commutative bimonoids in \mathbf{C} .*
2. *The category $\mathbf{Alg}(Q_{\mathcal{H}}, \mathbf{C})$ is equivalent to the category of involutive cocommutative bimonoids in \mathbf{C} .*
3. *The category $\mathbf{Alg}(Q_{\mathcal{IF}}, \mathbf{C})$ is equivalent to the category of involutive, commutative, cocommutative bimonoids in \mathbf{C} .*

PROOF. These equivalences are proved similarly to Theorem 5.3. ■

References

- [FL91] Zbigniew Fiedorowicz and Jean-Louis Loday. Crossed simplicial groups and their associated homology. *Trans. Amer. Math. Soc.*, 326(1):57–87, 1991.
- [FT87] B. L. Feĭgin and B. L. Tsygan. Additive K -theory. In *K-theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 67–209. Springer, Berlin, 1987.
- [Gra19] Daniel Graves. *Functor Homology for Augmented Algebras*, 2019. PhD Thesis. University of Sheffield. Available at <http://etheses.whiterose.ac.uk/26243/>.
- [Lac04] Stephen Lack. Composing PROPS. *Theory Appl. Categ.*, 13:No. 9, 147–163, 2004.
- [Mar96] Martin Markl. Cotangent cohomology of a category and deformations. *J. Pure Appl. Algebra*, 113(2):195–218, 1996.
- [ML98] Saunders Mac Lane. *Categories for the working mathematician*, volume 5 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1998.
- [Pir02] Teimuraz Pirashvili. On the PROP corresponding to bialgebras. *Cah. Topol. Géom. Différ. Catég.*, 43(3):221–239, 2002.
- [RW02] Robert Rosebrugh and R. J. Wood. Distributive laws and factorization. *J. Pure Appl. Algebra*, 175:327–353, 2002.

School of Mathematics and Statistics, University of Sheffield, Sheffield, S3 7RH, UK
 Email: dan.graves92@gmail.com

This article may be accessed at <http://www.tac.mta.ca/tac/>

THEORY AND APPLICATIONS OF CATEGORIES will disseminate articles that significantly advance the study of categorical algebra or methods, or that make significant new contributions to mathematical science using categorical methods. The scope of the journal includes: all areas of pure category theory, including higher dimensional categories; applications of category theory to algebra, geometry and topology and other areas of mathematics; applications of category theory to computer science, physics and other mathematical sciences; contributions to scientific knowledge that make use of categorical methods. Articles appearing in the journal have been carefully and critically refereed under the responsibility of members of the Editorial Board. Only papers judged to be both significant and excellent are accepted for publication.

SUBSCRIPTION INFORMATION Individual subscribers receive abstracts of articles by e-mail as they are published. To subscribe, send e-mail to tac@mta.ca including a full name and postal address. Full text of the journal is freely available at <http://www.tac.mta.ca/tac/>.

INFORMATION FOR AUTHORS \LaTeX 2e is required. Articles may be submitted in PDF by email directly to a Transmitting Editor following the author instructions at <http://www.tac.mta.ca/tac/authinfo.html>.

MANAGING EDITOR. Geoff Cruttwell, Mount Allison University: gcruttwell@mta.ca

TEXNICAL EDITOR. Michael Barr, McGill University: michael.barr@mcgill.ca

ASSISTANT TEX EDITOR. Gavin Seal, Ecole Polytechnique Fédérale de Lausanne: gavin_seal@fastmail.fm

TRANSMITTING EDITORS.

Clemens Berger, Université de Nice-Sophia Antipolis: cberger@math.unice.fr

Julie Bergner, University of Virginia: jeb2md@virginia.edu

Richard Blute, Université d' Ottawa: rblute@uottawa.ca

Gabriella Böhm, Wigner Research Centre for Physics: bohm.gabriella@wigner.mta.hu

Valeria de Paiva, Nuance Communications Inc: valeria.depaiva@gmail.com

Richard Garner, Macquarie University: richard.garner@mq.edu.au

Ezra Getzler, Northwestern University: getzler@northwestern.edu

Kathryn Hess, Ecole Polytechnique Fédérale de Lausanne: kathryn.hess@epfl.ch

Dirk Hofmann, Universidade de Aveiro: dirk@ua.pt

Pieter Hofstra, Université d' Ottawa: phofstra@uottawa.ca

Anders Kock, University of Aarhus: kock@math.au.dk

Joachim Kock, Universitat Autònoma de Barcelona: kock@mat.uab.cat

Stephen Lack, Macquarie University: steve.lack@mq.edu.au

Tom Leinster, University of Edinburgh: Tom.Leinster@ed.ac.uk

Matias Menni, Conicet and Universidad Nacional de La Plata, Argentina: matias.menni@gmail.com

Ieke Moerdijk, Utrecht University: i.moerdijk@uu.nl

Susan Niefield, Union College: niefiels@union.edu

Kate Ponto, University of Kentucky: kate.ponto@uky.edu

Robert Rosebrugh, Mount Allison University: rrosebrugh@mta.ca

Jiri Rosicky, Masaryk University: rosicky@math.muni.cz

Giuseppe Rosolini, Università di Genova: rosolini@disi.unige.it

Alex Simpson, University of Ljubljana: Alex.Simpson@fmf.uni-lj.si

James Stasheff, University of North Carolina: jds@math.upenn.edu

Ross Street, Macquarie University: ross.street@mq.edu.au

Tim Van der Linden, Université catholique de Louvain: tim.vanderlinden@uclouvain.be