



Update on 2009 Fraser Sockeye: Context, Forecasts & Returns

*Roundtable presentation
Vancouver September 11, 2009*




This talk will cover

1. Environmental context: increased variability & uncertainty
2. Preparation of pre-season forecasts
3. In-season management & 2009 Fraser sockeye returns


Part 1: Environmental context

A warming world


AOGCM Projections of Surface Temperatures



Impacts on salmon: Higher temperatures in streams, lakes and rivers



Fraser River Critical Year Comparison




By the end of the century, returning salmon will experience conditions in most years that are more severe than any year during the 1961-1990 base period.

Impacts in the open ocean – reduction in habitat for sockeye in the North Pacific?

*Projected southern limit of
sockeye salmon under a
doubling of CO2 climate
change scenario*

*1995 southern limit of
sockeye salmon distribution*





What can we expect?

- Changes will not be “smooth”
 - El Nino/La Nina and decadal climate variability overlay global warming. There will be periods (last year was one) where the ocean will cool rather than warm
- Our ability to predict future status of fish stocks will likely decrease as we move out of the historical range of our observations
 - Prediction of impacts has proven to be difficult at the management unit level e.g. Fraser River sockeye stocks do not all show a clear, coherent response to ocean conditions
- New issues and challenges
 - Hypoxia, acidification, disease


Part 2: Preparation of pre-season forecasts




Fraser Basin Watershed



Building a forecast



Role of Probability



Forecasts are presented as slices through a cumulative probability distribution (probability of exceeding a specified run size).





Probability of Exceeding Specified Run Sizes

Sockeye run timing group	Mean Run Size			
	2009 cycle	10%	50%	90%
Early Stuart	797,000	645,000	255,000	107,000
Early Summer	(316,000)	2,284,000	739,000	264,000
Summer	11,111,000	31,813,000	8,677,000	2,858,000
Late	(946,000)	2,875,000	907,000	327,000
Total	13,170,000	37,617,000	10,578,000	3,556,000

10

S. Grant & A. Cass (DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2009/022)

Forecasts have high uncertainty





Consider indicators of ocean conditions

	(BROOD YEAR) OCEAN ENTRY YEAR (RETURN YEAR)	(1996) 1998 (2000)	(1997) 1999 (2001)	(1998) 2000 (2002)	(1999) 2001 (2003)	(2000) 2002 (2004)	(2001) 2003 (2005)	(2002) 2004 (2006)	(2003) 2005 (2007)	(2004) 2006 (2008)	(2005) 2007 (2009)
Chilko Marine Survival		G	Y	G	G	R	Y	G	R	Y	R
Ocean Indices											
1 PDO (Jan-March average)		R	G	G	R	G	R	R	R	Y	Y
2 ALPI		R	G	Y	R	R	R	R	Y	G	G
Physical Conditions											
3 SST (Entrance Island)		R	G	G	G	G	R	R	R	Y	Y
4 SST (Pine Island)		R	G	G	G	Y	R	R	R	Y	G
5 Upwelling index (48°N)		G	G	R	Y	G	R	Y	R	Y	G
6 Spring transition timing (48°N)		G	G	Y	Y	G	Y	Y	R	Y	Y
Biological Conditions											
7 Southern Copepods (SVI)		R	G	Y	G	G	R	Y	R	R	G
8 Boreal Shelf Copepods (SVI)		R	G	G	Y	G	Y	R	R	R	G
9 Southern Copepods (NVI)		R	G	G	G	Y	R	Y	R	R	G
10 Boreal Shelf Copepods (NVI)		Y	G	G	R	G	R	R	R	Y	G

Part 3: In-season management &




Fraser River sockeye salmon returns



Test fisheries & acoustic monitoring to update estimates in-season



Graphs from www.psc.org




Each data point is a different year

Separate relationships for each week of fishing

Assume normal distribution for run timing


Operational Models to guide in-season decision-making

- A numerical model predicts Fraser River flow and temperature conditions
- These predictions drive a biological effects model to estimate in-river mortality
- If required, harvest levels are reduced to compensate for excessive predicted in-river mortality (i.e. management adjustment)



What happened in 2009?

DFO took immediate measures
to curtail harvests



Fraser Panel news release (Aug 7, 2009):

"current estimates of Summer-run sockeye run size are considerably below their 90% probability level forecast of 2,858,000 fish and are less than those needed to provide harvestable surpluses"

Fraser River Sockeye Catch Estimates

Commercial	-	
Recreational	-	
FN FSC – Marine	10,020	} 10-year average of 782,000
FN FSC – Fraser	49,240	
<i>Canadian Total</i>	59,260	
Test Fishing	33,146	
U.S.	3,400	
TOTAL	95,806	

Catches as of 08-Sep-2009

No First Nation group met their full FSC target. DFO took actions to provide access to all where possible.

Fraser Sockeye Escapement


Management Group	Escapement goal	Projected Escapement*	Actual Escapement For 2009 Cycle (1953-2005)
Early Stuart	85,000	55,370	222,000
Early Summer	175,000	98,720	100,000
Summer	520,000	466,980	1,850,000
Birkenhead	48,000	56,400	50,000
L.Lates	320,000	92,563	89,000
Total	1,148,000	770,033	2,311,000

* Escapement based upon in-season run size minus projected en-route mortality


Given 2009 management actions, harvest amounts to only 6% of the total return.

Chilko Lake sockeye survival

Freshwater Survival



Marine Survival



Preliminary 2009 return suggests lowest marine survival on record

Poor marine survival for 2007 return (2005 sea entry)

Preliminary 2009 salmon returns coastwide

Chinook salmon

Taku, Stikine, Skeena, Fraser (spring age 4): well below average

Fraser (spring / summer age 5): near expectation & below average

Fraser (summer age 4): near expectation & above average

Coho salmon

Taku, Skeena, WCVI, Columbia: average to well above average

Sockeye salmon

Taku, Stikine, Nass, Skeena, Smith, Fraser: well below average

Barkley Sound, Columbia: above expectation & below average

Pink salmon

Taku: below average

Skeena, Johnstone St, St of Georgia, Fraser, Puget Sound: above expectations; records in some areas